

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/355142359>

# Decision-Making Application Based on Aggregations of Complex Fuzzy Hypersoft Set and Development of Interval-Valued Complex Fuzzy Hypersoft Set

Article · October 2021

DOI: 10.5281/zenodo.5553532

CITATIONS

0

READS

13

5 authors, including:



**Atiqe Ur Rahman**

University of Management and Technology (Pakistan)

37 PUBLICATIONS 59 CITATIONS

[SEE PROFILE](#)



**Muhammad Saeed**

University of Management and Technology (Pakistan)

133 PUBLICATIONS 723 CITATIONS

[SEE PROFILE](#)



**Arooj Khalid**

University of Management and Technology (Pakistan)

1 PUBLICATION 0 CITATIONS

[SEE PROFILE](#)



**Muhammad Rayees Ahmad**

University of Management and Technology (Pakistan)

16 PUBLICATIONS 138 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Call for Papers, Scientific Inquiry and Review (SIR), Volume 2, Issue 2 [View project](#)



GENERALIZATION OF TOPSIS FROM SOFT SET TO FUZZY SOFT SETS IN DECISION MAKING PROBLEM [View project](#)



## Decision-Making Application Based on Aggregations of Complex Fuzzy Hypersoft Set and Development of Interval-Valued Complex Fuzzy Hypersoft Set

Atiqe Ur Rahman<sup>1,\*</sup>, Muhammad Saeed<sup>2</sup>, Arooj Khalid<sup>3</sup>, Muhammad Rayees Ahmad<sup>4</sup> and Sehrish Ayaz<sup>5</sup>

<sup>1,2,3,4,5</sup> Department of Mathematics, University of Management and Technology, Lahore, Pakistan ;  
aurkhhb@gmail.com; muhammad.saeed@umt.edu.pk; ; f2019109013@umt.edu.pk;  
rayeesmalik.ravian@gmail.com; f2019349001@umt.edu.pk

\*Correspondence: aurkhhb@gmail.com

**Abstract.** Hypersoft set, an extension of soft set, deals with disjoint attribute-valued sets corresponding to distinct attributes. In this study, the innovation of complex fuzzy hypersoft set (CFH-set) is conferred, which can tackle with uncertainties and vagueness that lie in the data by taking into account the amplitude and phase terms of the complex numbers at the same time. This model establishes a gluing framework of the fuzzy set and hypersoft set characterized in the complex plane. This structure is more flexible and useful as it consents a broad range of values for membership function by expanding them to the unit circle in a complex plane through the characterization of the fuzzy hypersoft set to consider the periodic nature of the information and the attributes can further be classified into attribute-values sets for vivid understanding. With the characterization of its some fundamental properties and operations, aggregations of complex fuzzy hypersoft set: matrix, cardinal set, cardinal matrix of cardinal set, aggregation operator/set and matrix of aggregation set, are conceptualized along with application in decision-making. Moreover, complex interval-valued fuzzy hypersoft set is developed and some of its fundamentals i.e. subset, equal sets, null set, absolute set etc. and theoretic operations i.e. compliment, union, intersection etc. are investigated.

**Keywords:** Complex fuzzy sets (CF-Sets), soft set, hypersoft set and complex fuzzy hypersoft set.

### 1. Introduction

The concept of complex fuzzy set theory (CFS-Theory) [1] is an extension of fuzzy set theory (FS-Theory) [2], which uses complex-valued state for the membership of its elements. FS-Theory and CFS-Theory have some kind of complexities which restrain them to solve problem involving uncertainty professionally. The reason for these hurdles is, possibly, the

inadequacy of the parametrization tool. It demands a mathematical tool free of all such impediments to tackle such issues. This scantiness is resolved with the development of soft set theory (SS-Theory) [7] which is a new parameterized family of subsets of the universe of discourse. The researchers [8]- [17] studied and investigated some elementary properties, operations, laws and hybrids of SS-Theory with applications in decision making. The gluing concept of NS-Theory and SS-Theory, is studied in [18] to make the NS-Theory adequate with parameterized tool. In many real life situations, distinct attributes are further partitioned in disjoint attribute-valued sets but existing SS-Theory is insufficient for dealing with such kind of attribute-valued sets. Hypersoft set theory (HS-Theory) [19] is developed to make the SST in line with attribute-valued sets to tackle real life scenarios. HS-Theory is an extension of SS-Theory as it transforms the single argument function into a multi-argument function. Certain elementary properties, aggregation operations, laws, relations and functions of HS-Theory, are investigated by [20]- [22] for proper understanding and further utilization in different fields. The applications of HS-Theory in decision making is studied by [23]- [27] and the intermingling study of HS-Theory with complex sets, convex and concave sets is studied by [28, 29]. Deli [30] characterized hybrid set structures under uncertainly parameterized hypersoft sets with theory and applications. Gayen et al. [31] analyzed some essential aspects of plithogenic hypersoft algebraic structures. They also investigated the notions and basic properties of plithogenic hypersoft subgroups ie plithogenic fuzzy hypersoft subgroup, plithogenic intuitionistic fuzzy hypersoft subgroup, plithogenic neutrosophic hypersoft subgroup. Saeed et al. [32, 33] discussed decision making techniques for neutrosophic hypersoft mapping and complex multi-fuzzy hypersoft set. Rahman et al. [34–36] studied decision making applications based on neutrosophic parameterized hypersoft Set, fuzzy parameterized hypersoft set and rough hypersoft set. Ihsan et al. [37] investigated hypersoft expert set with application in decision making for the best selection of product.

### 1.1. *Motivation*

In order to address the limitation of fuzzy soft set for dealing with periodic nature of data, Thirunavukarasu et al. [38] developed the theory of complex fuzzy soft set and discussed its some fundamentals along with applications. Kumar et al. [39] extended the work of Thirunavukarasu et al. to complex intuitionistic fuzzy soft sets and calculated its distance measures and entropies. Selvachandran et al. [40] investigated interval-valued complex fuzzy soft set with application. Abd et al. [41] discussed the fundamentals, properties and application of complex generalised fuzzy soft sets. These existing models employed single set of attributes for dealing uncertainties under fuzzy set-like environments but there are many situations when each attribute is required to be further partitioned into its attribute-valued set.

---

Atiqe Ur Rahman, Muhammad Saeed, Arooj Khalid, Muhammad Rayees Ahmad and Sehrish Ayaz,  
Decision-Making Application Based on Aggregations of Complex Fuzzy Hypersoft Set and  
Development of Interval-Valued Complex Fuzzy Hypersoft Set

These existing structures has limitation regarding the consideration of such attribute-valued sets. Inspiring from the above literature, the decision system of complex fuzzy hypersoft set is developed with the help of the characterization of its aggregation operations and fundamental theory of interval-valued complex fuzzy hypersoft set is investigated. The proposed structure complex fuzzy hypersoft set (CFH-set) and interval-valued complex hypersoft set (IV-CFHS) are more flexible and useful as they

- (i) generalize the existing structures of complex fuzzy soft set.
- (ii) permit a broad range of values for membership function by expanding them to the unit circle in a complex plane.
- (iii) consider the periodic nature of the information through the phase-terms.
- (iv) classify distinct attributes into corresponding attribute-values sets for vivid understanding.

## 1.2. Organization of Paper

The rest of the paper is organized as: section 2 reviews the notions of fuzzy set, soft set, complex fuzzy set and relevant definitions used in the proposed work. Section 3, presents the decision system of complex fuzzy hypersoft set based on its some decisive aggregation operations along with application in decision-making. Section 4, investigates the fundamental theory of interval-valued complex fuzzy hypersoft set. Lastly, paper is summarized with future directions.

## 2. Preliminaries

Here some existing fundamental concepts regarding fuzzy set, fuzzy soft set and fuzzy hypersoft set are presented along with their structures with complex fuzzy set from literature. Throughout the paper,  $\mathbb{U}$ ,  $P(\mathbb{U})$ ,  $F(\mathbb{U})$ ,  $C(\mathbb{U})$  and  $C_h(\mathbb{U})$  will present universe of discourse, power set of  $\mathbb{U}$ , collection of fuzzy sets, collection of complex fuzzy sets on soft sets and collection of complex fuzzy sets on hypersoft sets respectively.

### Definition 2.1. [2]

Suppose a universal set  $\mathbb{U}$  and a fuzzy set  $X \subseteq \mathbb{U}$ . The set  $X$  will be written as  $X = \{(x, \alpha_X(x)) | x \in \mathbb{U}\}$  such that

$$\alpha_X : \mathbb{U} \rightarrow [0, 1]$$

where  $\alpha_X(x)$  describes the membership percentage of  $x \in X$ .

### Definition 2.2. [1]

A complex fuzzy set  $\mathbb{C}_f$  is of the form

$$\mathbb{C}_f = \{(\epsilon, \mu_{\mathbb{C}_f}(\epsilon)) : \epsilon \in \mathbb{U}\} = \{(\epsilon, r_{\mathbb{C}_f}(\epsilon)e^{i\omega_{\mathbb{C}_f}(\epsilon)}) : \epsilon \in \mathbb{U}\}.$$

where  $\mu_{\mathbb{C}_f}(\epsilon)$  is a membership function of  $\mathbb{C}_f$  with  $r_{\mathbb{C}_f}(\epsilon) \in [0, 1]$  and  $\omega_{\mathbb{C}_f}(\epsilon) \in (0, 2\pi]$  as amplitude and phase terms respectively and  $i = \sqrt{-1}$ .

Buckley [3] and Zhang et al. [4] presented fuzzy complex number in different way. However, according to [5]- [6], both amplitude and phase terms are captured by fuzzy sets.

**Definition 2.3.** [7]

A *soft set*  $\mathfrak{S}$  over  $\mathbb{U}$ , is defined as

$$\mathfrak{S} = \{(\epsilon, f_{\mathfrak{S}}(\epsilon)) : \epsilon \in E_1\}$$

where  $f_{\mathfrak{S}} : E_1 \rightarrow P(\mathbb{U})$ . and  $E_1 \subseteq E$  (set of parameters).

**Definition 2.4.** [9]

A *fuzzy soft set* (FS-set)  $\Gamma_{E_1}$  on  $\mathbb{U}$ , is defined as

$$\Gamma_{E_1} = \{(\epsilon, \gamma_{E_1}(\epsilon)) : \epsilon \in E_1, \gamma_{E_1}(\epsilon) \in F(\mathbb{U})\}$$

where  $\gamma_{E_1} : E_1 \rightarrow F(\mathbb{U})$  such that  $\gamma_{E_1}(\epsilon) = \emptyset$  if  $\epsilon \notin E_1$ , and for all  $\epsilon \in E_1$ ,

$$\gamma_{E_1}(\epsilon) = \left\{ \mu_{\gamma_{E_1}(\epsilon)}(v)/v : v \in \mathbb{U}, \mu_{\gamma_{E_1}(\epsilon)}(v) \in [0, 1] \right\}$$

is a fuzzy set over  $\mathbb{U}$ . Also  $\gamma_{E_1}$  is the approximate function of  $\Gamma_{E_1}$  and the value  $\gamma_A(x)$  is a fuzzy set called  $\epsilon$ -element of FS-set. Note that if  $\gamma_{E_1}(\epsilon) = \emptyset$ , then  $(\epsilon, \gamma_{E_1}(\epsilon)) \notin \Gamma_{E_1}$ .

**Definition 2.5.** [38]

A *complex fuzzy soft set* (CFS-set)  $\chi_{E_1}$  over  $\mathbb{U}$ , is defined as

$$\chi_{E_1} = \{(\epsilon, \psi_{E_1}(\epsilon)) : \epsilon \in E_1, \psi_{E_1}(\epsilon) \in C(\mathbb{U})\}.$$

where  $\psi_{E_1} : E_1 \rightarrow C(\mathbb{U})$  such that  $\psi_{E_1}(\epsilon) = \emptyset$  if  $\epsilon \notin E_1$  and it is complex fuzzy approximate function of CFS-set  $\chi_{E_1}$  and its value  $\psi_{E_1}(\epsilon)$  is called  $\epsilon$ -member of CFS-set  $\chi_{E_1}$  for all  $\epsilon \in E_1$ . Operations of CF-sets and CFS-sets were defined in [1] and [38] respectively.

**Definition 2.6.** [19]

The pair  $(H, G)$  is called a *hypersoft set* over  $\mathbb{U}$ , where  $G$  is the cartesian product of  $n$  disjoint sets  $H_1, H_2, H_3, \dots, H_n$  having attribute values of  $n$  distinct attributes  $h_1, h_2, h_3, \dots, h_n$  respectively and  $H : G \rightarrow P(\mathbb{U})$ .

**Definition 2.7.** [19]

A hypersoft set over a fuzzy universe of discourse is called *fuzzy hypersoft set*.

For more definitions and operations of hypersoft set, see [20]- [22]

### 2.1. Complex Fuzzy Hypersoft Set

The following subsections 2.1 and 2.2 are reviewed from [28].

**Definition 2.8.** Let  $A_1, A_2, A_3, \dots, A_n$  are disjoint sets having attribute values of  $n$  distinct attributes  $a_1, a_2, a_3, \dots, a_n$  respectively for  $n \geq 1$ ,  $G = A_1 \times A_2 \times A_3 \times \dots \times A_n$  and  $\psi(\underline{x})$  be a CF-set over  $\mathbb{U}$  for all  $\underline{\epsilon} = (d_1, d_2, d_3, \dots, d_n) \in G$ . Then, *complex fuzzy hypersoft set* (CFH-set)  $\chi_G$  over  $\mathbb{U}$  is defined as

$$\chi_G = \{(\underline{\epsilon}, \psi(\underline{\epsilon})) : \underline{\epsilon} \in G, \psi(\underline{\epsilon}) \in C(\mathbb{U})\}$$

where

$$\psi : G \rightarrow C(\mathbb{U}), \quad \psi(\underline{\epsilon}) = \emptyset \text{ if } \underline{\epsilon} \notin G.$$

is a CF-approximate function of  $\chi_G$  and its value  $\psi(\underline{\epsilon})$  is called  $\underline{\epsilon}$ -member of CFH-set  $\forall \underline{\epsilon} \in G$ .

**Example 2.9.** Suppose a Department Promotion Committee (DPC) wants to observe(evaluate) the characteristics of some teachers by some defined indicators for departmental promotion. For this purpose, consider a set of teachers as a universe of discourse  $\mathbb{U} = \{t_1, t_2, t_3, t_4\}$ . The attributes of the teachers under consideration are the set  $E = \{A_1, A_2, A_3\}$ , where

$A_1$  = Total experience in years =  $\{3, < 10\} = \{e_{11}, e_{12}\}$

$A_2$  = Total no. of publications =  $\{10, 10 <\} = \{e_{21}, e_{22}\}$

$A_3$  = Performance Evaluation Report (PER) remarks =  $\{eligible, not\ eligible\} = \{e_{31}, e_{32}\}$

and

$$G = A_1 \times A_2 \times A_3 = \left\{ \begin{array}{l} (e_{11}, e_{21}, e_{31}), (e_{11}, e_{21}, e_{32}), (e_{11}, e_{22}, e_{31}), \\ (e_{11}, e_{22}, e_{32}), (e_{12}, e_{21}, e_{31}), (e_{12}, e_{21}, e_{32}), \\ (e_{12}, e_{22}, e_{31}), (e_{12}, e_{22}, e_{32}) \end{array} \right\} = \{e_1, e_2, e_3, \dots, e_8\}$$

Complex fuzzy set  $\psi_G(e_1), \psi_G(e_2), \dots, \psi_G(e_8)$  are defined as,

$$\begin{aligned} \psi_G(e_1) &= \left\{ \frac{0.4e^{i0.5\pi}}{t_1}, \frac{0.8e^{i0.6\pi}}{t_2}, \frac{0.8e^{i0.8\pi}}{t_3}, \frac{1.0e^{i0.75\pi}}{t_4} \right\}, \\ \psi_G(e_2) &= \left\{ \frac{0.6e^{i0.7\pi}}{t_1}, \frac{0.9e^{i0.9\pi}}{t_2}, \frac{0.7e^{i0.9\pi}}{t_3}, \frac{0.75e^{i0.95\pi}}{t_4} \right\}, \\ \psi_G(e_3) &= \left\{ \frac{0.5e^{i0.6\pi}}{t_1}, \frac{0.8e^{i0.9\pi}}{t_2}, \frac{0.6e^{i0.9\pi}}{t_3}, \frac{0.65e^{i0.95\pi}}{t_4} \right\}, \\ \psi_G(e_4) &= \left\{ \frac{0.3e^{i0.7\pi}}{t_1}, \frac{0.7e^{i0.9\pi}}{t_2}, \frac{0.5e^{i0.9\pi}}{t_3}, \frac{0.75e^{i0.65\pi}}{t_4} \right\}, \\ \psi_G(e_5) &= \left\{ \frac{0.2e^{i0.5\pi}}{t_1}, \frac{0.3e^{i0.8\pi}}{t_2}, \frac{0.8e^{i0.7\pi}}{t_3}, \frac{0.45e^{i0.65\pi}}{t_4} \right\}, \\ \psi_G(e_6) &= \left\{ \frac{0.5e^{i0.9\pi}}{t_1}, \frac{0.3e^{i0.9\pi}}{t_2}, \frac{0.7e^{i0.8\pi}}{t_3}, \frac{0.85e^{i0.95\pi}}{t_4} \right\}, \\ \psi_G(e_7) &= \left\{ \frac{0.6e^{i0.9\pi}}{t_1}, \frac{0.9e^{i0.6\pi}}{t_2}, \frac{0.5e^{i0.6\pi}}{t_3}, \frac{0.85e^{i0.75\pi}}{t_4} \right\}, \end{aligned}$$

and

$$\psi_G(e_8) = \left\{ \frac{0.8e^{i0.9\pi}}{t_1}, \frac{0.8e^{i0.8\pi}}{t_2}, \frac{0.6e^{i0.8\pi}}{t_3}, \frac{0.65e^{i0.85\pi}}{t_4} \right\}$$

then CFH-set  $\chi_G$  is written by,

$$\chi_G = \left\{ \begin{array}{l} (e_1, \frac{0.4e^{i0.5\pi}}{t_1}, \frac{0.8e^{i0.6\pi}}{t_2}, \frac{0.8e^{i0.8\pi}}{t_3}, \frac{1.0e^{i0.75\pi}}{t_4}), (e_2, \frac{0.6e^{i0.7\pi}}{t_1}, \frac{0.9e^{i0.9\pi}}{t_2}, \frac{0.7e^{i0.9\pi}}{t_3}, \frac{0.75e^{i0.95\pi}}{t_4}), \\ (e_3, \frac{0.5e^{i0.6\pi}}{t_1}, \frac{0.8e^{i0.9\pi}}{t_2}, \frac{0.6e^{i0.9\pi}}{t_3}, \frac{0.65e^{i0.95\pi}}{t_4}), (e_4, \frac{0.3e^{i0.7\pi}}{t_1}, \frac{0.7e^{i0.9\pi}}{t_2}, \frac{0.5e^{i0.9\pi}}{t_3}, \frac{0.75e^{i0.65\pi}}{t_4}), \\ (e_5, \frac{0.2e^{i0.5\pi}}{t_1}, \frac{0.3e^{i0.8\pi}}{t_2}, \frac{0.8e^{i0.7\pi}}{t_3}, \frac{0.45e^{i0.65\pi}}{t_4}), (e_6, \frac{0.5e^{i0.9\pi}}{t_1}, \frac{0.3e^{i0.9\pi}}{t_2}, \frac{0.7e^{i0.8\pi}}{t_3}, \frac{0.85e^{i0.95\pi}}{t_4}), \\ (e_7, \frac{0.6e^{i0.9\pi}}{t_1}, \frac{0.9e^{i0.6\pi}}{t_2}, \frac{0.5e^{i0.6\pi}}{t_3}, \frac{0.85e^{i0.75\pi}}{t_4}), (e_8, \frac{0.8e^{i0.9\pi}}{t_1}, \frac{0.8e^{i0.8\pi}}{t_2}, \frac{0.6e^{i0.8\pi}}{t_3}, \frac{0.65e^{i0.85\pi}}{t_4}) \end{array} \right\}$$

**Definition 2.10.** Let  $\chi_{G_1} = (\psi_1, G_1)$  and  $\chi_{G_2} = (\psi_2, G_2)$  be two CFH-sets over the same  $\mathbb{U}$ .

The set  $\chi_{G_1} = (\psi_1, G_1)$  is said to be the *subset* of  $\chi_{G_2} = (\psi_2, G_2)$ , if

- i.  $G_1 \subseteq G_2$
- ii.  $\forall \underline{x} \in G_1, \psi_1(\underline{x}) \subseteq \psi_2(\underline{x})$  i.e.  $r_{G_1}(\underline{x}) \leq r_{G_2}(\underline{x})$  and  $\omega_{G_1}(\underline{x}) \leq \omega_{G_2}(\underline{x})$ , where  $r_{G_1}(\underline{x})$  and  $\omega_{G_1}(\underline{x})$  are amplitude and phase terms of  $\psi_1(\underline{x})$ , whereas  $r_{G_2}(\underline{x})$  and  $\omega_{G_2}(\underline{x})$  are amplitude and phase terms of  $\psi_2(\underline{x})$ .

**Definition 2.11.** Two CFH-sets  $\chi_{G_1} = (\psi_1, G_1)$  and  $\chi_{G_2} = (\psi_2, G_2)$  over the same  $\mathbb{U}$ , are said to be *equal* if

- i.  $(\psi_1, G_1) \subseteq (\psi_2, G_2)$
- ii.  $(\psi_2, G_2) \subseteq (\psi_1, G_1)$ .

**Definition 2.12.** Let  $(\psi, G)$  be a CFH-set over  $\mathbb{U}$ . Then

- i.  $(\psi, G)$  is called a *null CFH-set*, denoted by  $(\psi, G)_\Phi$  if for all  $\underline{x} \in G$ , the amplitude and phase terms of the membership function are given by  $r_G(\underline{x}) = 0$  and  $\omega_G(\underline{x}) = 0\pi$  respectively.
- ii.  $(\psi, G)$  is called a *absolute CFH-set*, denoted by  $(\psi, G)_\Delta$  if for all  $\underline{x} \in G$ , the amplitude and phase terms of the membership function are given by  $r_G(\underline{x}) = 1$  and  $\omega_G(\underline{x}) = 2\pi$  respectively.

**Definition 2.13.** Let  $(\psi_1, G_1)$  and  $(\psi_2, G_2)$  are two CFH-sets over the same universe  $\mathbb{U}$ . Then

- i. A CFH-set  $(\psi_1, G_1)$  is called a *homogeneous CFH-set*, denoted by  $(\psi_1, G_1)_{Hom}$  if and only if  $\psi_1(\underline{x})$  is a homogeneous CF-set for all  $\underline{x} \in G_1$ .
- ii. A CFH-set  $(\psi_1, G_1)$  is called a *completely homogeneous CFH-set*, denoted by  $(\psi_1, G_1)_{CHom}$  if and only if  $\psi_1(\underline{x})$  is a homogeneous with  $\psi_1(\underline{y})$  for all  $\underline{x}, \underline{y} \in G_1$ .
- iii. A CFH-set  $(\psi_1, G_1)$  is said to be a completely homogeneous CFH-set with  $(\psi_2, G_2)$  if and only if  $\psi_1(\underline{x})$  is a homogeneous with  $\psi_2(\underline{x})$  for all  $\underline{x} \in G_1 \cap G_2$ .

## 2.2. Set Theoretic Operations and Laws on CFH-Sets

Here some basic set theoretic operations (i.e. complement, union and intersection) and laws (commutative laws, associative laws etc.) are discussed on CFH-sets.

**Definition 2.14.** The *complement* of CFH-set  $(\psi, G)$ , denoted by  $(\psi, G)^c$  is defined as

$$(\psi, G)^c = \{(\underline{x}, \psi^c(\underline{x})) : \underline{x} \in G, \psi^c(\underline{x}) \in C(\mathbb{U})\}$$

such that the amplitude and phase terms of the membership function  $\psi^c(\underline{x})$  are given by  $r_G^c(\underline{x}) = 1 - r_G(\underline{x})$  and  $\omega_G^c(\underline{x}) = 2\pi - \omega_G(\underline{x})$  respectively.

**Proposition 2.15.** Let  $(\psi, G)$  be a CFH-set over  $\mathbb{U}$ . Then  $((\psi, G)^c)^c = (\psi, G)$ .

*Proof.* Since  $\psi(\underline{x}) \in C(\mathbb{U})$ , therefore  $(\psi, G)$  can be written in terms of its amplitude and phase terms as

$$(\psi, G) = \left\{ \left( \underline{x}, r_G(\underline{x})e^{i\omega_G(\underline{x})} \right) : \underline{x} \in G \right\} \quad (1)$$

Now

$$\begin{aligned} \psi^c(\underline{x}) &= \left\{ \left( \underline{x}, r_G^c(\underline{x})e^{i\omega_G^c(\underline{x})} \right) : \underline{x} \in G \right\} \\ \psi^c(\underline{x}) &= \left\{ \left( \underline{x}, (1 - r_G(\underline{x}))e^{i(2\pi - \omega_G(\underline{x}))} \right) : \underline{x} \in G \right\} \\ ((\psi, G)^c)^c &= \left\{ \left( \underline{x}, (1 - r_G(\underline{x}))^c e^{i(2\pi - \omega_G(\underline{x})^c)} \right) : \underline{x} \in G \right\} \\ ((\psi, G)^c)^c &= \left\{ \left( \underline{x}, (1 - (1 - r_G(\underline{x})))e^{i(2\pi - (2\pi - \omega_G(\underline{x})))} \right) : \underline{x} \in G \right\} \\ ((\psi, G)^c)^c &= \left\{ \left( \underline{x}, r_G(\underline{x})e^{i\omega_G(\underline{x})} \right) : \underline{x} \in G \right\} \end{aligned} \quad (2)$$

from equations (1) and (2), we have  $((\psi, G)^c)^c = (\psi, G)$ .  $\square$

**Proposition 2.16.** Let  $(\psi, G)$  be a CFH-set over  $\mathbb{U}$ . Then

- i.  $((\psi, G)_\Phi)^c = (\psi, G)_\Delta$
- ii.  $((\psi, G)_\Delta)^c = (\psi, G)_\Phi$

**Definition 2.17.** The *intersection* of two CFH-sets  $(\psi_1, G_1)$  and  $(\psi_2, G_2)$  over the same universe  $\mathbb{U}$ , denoted by  $(\psi_1, G_1) \cap (\psi_2, G_2)$ , is the CFH-set  $(\psi_3, G_3)$ , where  $G_3 = G_1 \cap G_2$ , and  $\psi_3(\underline{x}) = \psi_1(\underline{x}) \cap \psi_2(\underline{x})$  for all  $\underline{x} \in G_3$ .

**Definition 2.18.** The *difference* between two CFH-sets  $(\psi_1, G_1)$  and  $(\psi_2, G_2)$  is defined as

$$(\psi_1, G_1) \setminus (\psi_2, G_2) = (\psi_1, G_1) \cap (\psi_2, G_2)^c$$

**Definition 2.19.** The *union* of two CFH-sets  $(\psi_1, G_1)$  and  $(\psi_2, G_2)$  over the same universe  $\mathbb{U}$ , denoted by  $(\psi_1, G_1) \cup (\psi_2, G_2)$ , is the CFH-set  $(\psi_3, G_3)$ , where  $G_3 = G_1 \cup G_2$ , and for all  $\underline{x} \in G_3$ ,

$$\psi_3(\underline{x}) = \begin{cases} \psi_1(\underline{x}) & , \text{if } \underline{x} \in G_1 \setminus G_2 \\ \psi_2(\underline{x}) & , \text{if } \underline{x} \in G_2 \setminus G_1 \\ \psi_1(\underline{x}) \cup \psi_2(\underline{x}) & , \text{if } \underline{x} \in G_1 \cap G_2 \end{cases}$$

**Proposition 2.20.** Let  $(\psi, G)$  be a CFH-set over  $\mathbb{U}$ . Then the following results hold true:

- i.  $(\psi, G) \cup (\psi, G)_{\Phi} = (\psi, G)$
- ii.  $(\psi, G) \cup (\psi, G)_{\Delta} = (\psi, G)_{\Delta}$
- iii.  $(\psi, G) \cap (\psi, G)_{\Phi} = (\psi, G)_{\Phi}$
- iv.  $(\psi, G) \cap (\psi, G)_{\Delta} = (\psi, G)$
- v.  $(\psi, G)_{\Phi} \cup (\psi, G)_{\Delta} = (\psi, G)_{\Delta}$
- vi.  $(\psi, G)_{\Phi} \cap (\psi, G)_{\Delta} = (\psi, G)_{\Phi}$

**Proposition 2.21.** Let  $(\psi_1, G_1)$ ,  $(\psi_2, G_2)$  and  $(\psi_3, G_3)$  are three CFH-sets over the same universe  $\mathbb{U}$ . Then the following commutative and associative laws hold true:

- i.  $(\psi_1, G_1) \cap (\psi_2, G_2) = (\psi_2, G_2) \cap (\psi_1, G_1)$
- ii.  $(\psi_1, G_1) \cup (\psi_2, G_2) = (\psi_2, G_2) \cup (\psi_1, G_1)$
- iii.  $(\psi_1, G_1) \cap ((\psi_2, G_2) \cap (\psi_3, G_3)) = ((\psi_1, G_1) \cap (\psi_2, G_2)) \cap (\psi_3, G_3)$
- iv.  $(\psi_1, G_1) \cup ((\psi_2, G_2) \cup (\psi_3, G_3)) = ((\psi_1, G_1) \cup (\psi_2, G_2)) \cup (\psi_3, G_3)$

**Proposition 2.22.** Let  $(\psi_1, G_1)$  and  $(\psi_2, G_2)$  are two CFH-sets over the same universe  $\mathbb{U}$ . Then the following De Morgan's laws hold true:

- i.  $((\psi_1, G_1) \cap (\psi_2, G_2))^c = (\psi_1, G_1)^c \cup (\psi_2, G_2)^c$
- ii.  $(\psi_1, G_1) \cup (\psi_2, G_2))^c = (\psi_1, G_1)^c \cap (\psi_2, G_2)^c$

### 3. Aggregation of Complex Fuzzy Hypersoft Set

In this section, we define an aggregation operator on complex fuzzy hypersoft set that produces an aggregate fuzzy set from a complex fuzzy hypersoft set and its cardinal set. The approximate functions of a complex fuzzy hypersoft set are fuzzy. Here  $G, E, \chi_G$  and  $C_H(\mathbb{U})$  will be in accordance with definition (2.8).

**Definition 3.1.** Let  $\chi_G \in C_H(\mathbb{U})$ . Assume that  $\mathbb{U} = \{u_1, u_2, \dots, u_m\}$  and  $E = \{A_1, A_2, \dots, A_n\}$  with

$$A_1 = \{e_{11}, e_{12}, \dots, e_{1n}\}, A_2 = \{e_{21}, e_{22}, \dots, e_{2n}\}, \dots, A_n = \{e_{n1}, e_{n2}, \dots, e_{nn}\}$$

and  $G = A_1 \times A_2 \times \dots \times A_n = \{x_1, x_2, \dots, x_n, \dots, x_{n^n} = x_r\}$ , each  $x_i$  is n-tuple element of  $G$  and  $|G| = r = n^n$  then the  $\chi_G$  can be presented as

$\chi_G$	$x_1$	$x_2$	$\dots$	$x_r$
$u_1$	$\mu_{\psi_G(x_1)}(u_1)$	$\mu_{\psi_G(x_2)}(u_1)$	$\dots$	$\mu_{\psi_G(x_r)}(u_1)$
$u_2$	$\mu_{\psi_G(x_1)}(u_2)$	$\mu_{\psi_G(x_2)}(u_2)$	$\dots$	$\mu_{\psi_G(x_r)}(u_2)$
$\vdots$	$\vdots$	$\vdots$	$\ddots$	$\vdots$
$u_m$	$\mu_{\psi_G(x_1)}(u_m)$	$\mu_{\psi_G(x_2)}(u_m)$	$\dots$	$\mu_{\psi_G(x_r)}(u_m)$

Where  $\mu_{\psi_G(x)}$  is the membership function of  $\psi_G$ . If  $a_{ij} = \mu_{\psi_G(x_j)}(u_i)$ , for  $i = \mathbb{N}_1^m$  and  $j = \mathbb{N}_1^r$  then CFH-set  $\chi_G$  is uniquely characterized by a matrix,

$$[a_{ij}] = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$

is called an  $m \times r$  CFH-set matrix..

**Definition 3.2.** Let  $\chi_G \in C_H(\mathbb{U})$ . Then, the *cardinal set* of  $\chi_G$  is defined as

$$\|\chi_G\| = \{\mu_{\|\chi_G\|}(\underline{x})/\underline{x} : \underline{x} \in G\},$$

where  $\mu_{\|\chi_G\|} : G \rightarrow [0, 1]$  is a membership function of  $\|\chi_G\|$  with  $\mu_{Card(\chi_G)(\underline{x})} = \frac{|\psi_G(\underline{x})|}{|U|}$ .

Note that  $\|C_H(\mathbb{U})\|$  is the collection of all cardinal sets of CFH-sets and  $\|C_H(\mathbb{U})\| \subseteq F(G)$ .

**Definition 3.3.** Let  $\chi_G \in C_H(\mathbb{U})$  and  $\|\chi_G\| \in \|C_H(\mathbb{U})\|$ . Consider  $E$  as in definition (4.1) then  $\|\chi_G\|$  can be presented as

G	$x_1$	$x_2$	$\dots$	$x_r$
$\mu_{\ \chi_G\ }$	$\mu_{\ \chi_G\ }(x_1)$	$\mu_{\ \chi_G\ }(x_2)$	$\dots$	$\mu_{\ \chi_G\ }(x_r)$

If  $a_{1j} = \mu_{\|\chi_G\|}(x_j)$ , for  $j = \mathbb{N}_1^r$  then the cardinal set  $\|\chi_G\|$  is represented by a matrix,

$$[a_{ij}]_{1 \times n} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \end{bmatrix}$$

and is called *cardinal matrix* of  $\|\chi_G\|$ .

**Definition 3.4.** Let  $\chi_G \in C_H(\mathbb{U})$  and  $\|\chi_G\| \in \|C_H(\mathbb{U})\|$ . Then *CFH-aggregation operator* is defined as

$$\widehat{\chi_G} = A_{CFH}(\|\chi_G\|, \chi_G)$$

where

$$A_{CFH} : \|C_H(\mathbb{U})\| \times C_H(\mathbb{U}) \rightarrow F(U).$$

$\widehat{\chi_G}$  is called the aggregate fuzzy set of CFH-set  $\chi_G$ .

Its membership function is given as

$$\mu_{\widehat{\chi_G}} : U \rightarrow [0, 1]$$

with

$$\mu_{\widehat{\chi}_G}(u) = \frac{1}{|G|} \sum_{\underline{x} \in G} \mu_{Card(\chi_G)}(\underline{x}) \mu_{Card(\psi_G)}(u).$$

**Definition 3.5.** Let  $\chi_G \in C_H(\mathbb{U})$  and  $\widehat{\chi}_G$  be its aggregate fuzzy set. Assume that  $\mathbb{U} = \{u_1, u_2, \dots, u_m\}$ , then  $\widehat{\chi}_G$  can be presented as

$\chi_G$	$\mu_{\widehat{\chi}_G}$
$u_1$	$\mu_{\widehat{\chi}_G}(u_1)$
$u_2$	$\mu_{\widehat{\chi}_G}(u_2)$
$\vdots$	$\vdots$
$u_m$	$\mu_{\widehat{\chi}_G}(u_m)$

If  $a_{i1} = \mu_{\widehat{\chi}_G}(u_i)$  for  $i = \mathbb{N}_1^m$  then  $\widehat{\chi}_G$  is represented by the matrix,

$$[a_{i1}]_{m \times 1} = \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix}$$

which is called *aggregate matrix* of  $\widehat{\chi}_G$  over  $\mathbb{U}$ .

### 3.1. Applications of Complex Fuzzy Hypersoft Set

In this section, an algorithm is presented to solve the problems in decision making by having under consideration the concept of aggregations defined in previous section. An example is demonstrated to explain the proposed algorithm.

It is necessary to determine an aggregate fuzzy set of CFH-set for choosing the best option (parameter) from the given set (set of choices/alternatives). Following algorithm may help in making appropriate decision.

**Step 1:** Determine a CFH-set  $\chi_G$  over  $\mathbb{U}$ ,

**Step 2:** Determine  $\|\chi_G\|$  for amplitude term and phase term separately,

**Step 3:** Find  $\widehat{\chi}_G$  for amplitude term and phase term separately,

**Step 4:** Find the best option by max modulus of  $\mu_{\widehat{\chi}_G}(u)$

**Example 3.6.** Suppose a business man wants to buy a share from share market. There are four same kind of share which form the set,  $\mathbb{U} = \{s_1, s_2, s_3, s_4\}$ . The expert committee consider a set of attributes,  $E = \{e_1, e_2, e_3\}$ . For  $i = 1, 2, 3, 4$ , the attributes  $e_i$  stand for current trend of company performance, particular company's stock price for last one year, and Home

country inflation rate, respectively. Corresponding to each attribute, the sets of attribute values are:  $A_1 = \{e_{11}, e_{12}\}$ ;  $A_2 = \{e_{21}\}$  and  $A_3 = \{e_{31}, e_{32}\}$ . Then the set  $G = A_1 \times A_2 \times A_3 = \{\epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4\}$  where each  $\epsilon_i$  is a 3-tuple. Complex fuzzy sets  $\psi_G(\epsilon_1), \psi_G(\epsilon_2), \psi_G(\epsilon_3), \psi_G(\epsilon_4)$  are defined as,

$$\psi_G(\epsilon_1) = \left\{ \frac{0.4e^{i0.5\pi}}{s_1}, \frac{0.8e^{i0.6\pi}}{s_2}, \frac{0.8e^{i0.8\pi}}{s_3}, \frac{1.0e^{i0.75\pi}}{s_4} \right\},$$

$$\psi_G(\epsilon_2) = \left\{ \frac{0.3e^{i0.7\pi}}{s_1}, \frac{0.6e^{i0.8\pi}}{s_2}, \frac{0.5e^{i0.2\pi}}{s_3}, \frac{1.0e^{i0.85\pi}}{s_4} \right\},$$

$$\psi_G(\epsilon_3) = \left\{ \frac{0.6e^{i0.7\pi}}{s_1}, \frac{0.9e^{i0.9\pi}}{s_2}, \frac{0.7e^{i0.95\pi}}{s_3}, \frac{0.75e^{i0.95\pi}}{s_4} \right\},$$

and

$$\psi_G(\epsilon_4) = \left\{ \frac{0.5e^{i0.6\pi}}{s_1}, \frac{0.7e^{i0.8\pi}}{s_2}, \frac{0.6e^{i0.85\pi}}{s_3}, \frac{0.75e^{i0.85\pi}}{s_4} \right\},$$

Step 1: CFH-set  $\chi_G$  is written as,

$$\chi_G = \left\{ \left( \epsilon_1, \frac{0.4e^{i0.5\pi}}{s_1}, \frac{0.8e^{i0.6\pi}}{s_2}, \frac{0.8e^{i0.8\pi}}{s_3}, \frac{1.0e^{i0.75\pi}}{s_4} \right), \left( \epsilon_2, \frac{0.3e^{i0.7\pi}}{s_1}, \frac{0.6e^{i0.8\pi}}{s_2}, \frac{0.5e^{i0.2\pi}}{s_3}, \frac{1.0e^{i0.85\pi}}{s_4} \right), \right. \\ \left. \left( \epsilon_3, \frac{0.6e^{i0.7\pi}}{s_1}, \frac{0.9e^{i0.9\pi}}{s_2}, \frac{0.7e^{i0.95\pi}}{s_3}, \frac{0.75e^{i0.95\pi}}{s_4} \right), \left( \epsilon_4, \frac{0.5e^{i0.6\pi}}{s_1}, \frac{0.7e^{i0.8\pi}}{s_2}, \frac{0.6e^{i0.85\pi}}{s_3}, \frac{0.75e^{i0.85\pi}}{s_4} \right) \right\}$$

Step 2: The cardinal is computed as,

$$\|\chi_G\| (Amplitude Term) = \{0.75/\epsilon_1, 0.6/\epsilon_2, 0.74/\epsilon_3, 0.64/\epsilon_4\}$$

$$\|\chi_G\| (Phase Term) = \{0.66/\epsilon_1, 0.64/\epsilon_2, 0.87/\epsilon_3, 0.78/\epsilon_4\}$$

Step 3: The set  $\widehat{\chi}_G$  can be determined as,

$$\widehat{\chi}_G (Amplitude Term) = \frac{1}{4} \begin{bmatrix} 0.4 & 0.3 & 0.6 & 0.5 \\ 0.8 & 0.6 & 0.9 & 0.7 \\ 0.8 & 0.5 & 0.7 & 0.6 \\ 1.0 & 1.0 & 0.75 & 0.75 \end{bmatrix} \begin{bmatrix} 0.75 \\ 0.6 \\ 0.74 \\ 0.64 \end{bmatrix} = \begin{bmatrix} 0.3110 \\ 0.5185 \\ 0.4505 \\ 0.5963 \end{bmatrix}$$

$$\widehat{\chi}_G (Phase Term) = \frac{1}{4} \begin{bmatrix} 0.5 & 0.7 & 0.7 & 0.6 \\ 0.6 & 0.8 & 0.9 & 0.8 \\ 0.8 & 0.2 & 0.95 & 0.85 \\ 0.75 & 0.85 & 0.95 & 0.85 \end{bmatrix} \begin{bmatrix} 0.66 \\ 0.64 \\ 0.87 \\ 0.78 \end{bmatrix} = \begin{bmatrix} 0.4638 \\ 0.5788 \\ 0.5364 \\ 0.6321 \end{bmatrix}$$

$$\widehat{\chi}_G = \{0.3110e^{i0.4638\pi}/s_1, 0.5185e^{i0.5788\pi}/s_2, 0.4505e^{i0.5364\pi}/s_3, 0.5963e^{i0.6321\pi}/s_4\}$$

Consider the modulus value of  $Max(\mu_{\widehat{\chi}_G}) = \{0.3109/s_1, 0.5185/s_2, 0.4504/s_3, 0.5963/s_4\} = 0.5963/s_4$  This means that the 4th share  $s_4$  may be recommended for suitable investment.

#### 4. Interval-Valued Complex Fuzzy Hypersoft Set(IV-CFHS)

In this section, the basic theory of interval-valued complex fuzzy hypersoft set is developed.

**Definition 4.1.** Let  $W_1, W_2, W_3, \dots, W_n$  are disjoint sets having attribute values of  $n$  distinct attributes  $w_1, w_2, w_3, \dots, w_n$  respectively for  $n \geq 1, W = W_1 \times W_2 \times W_3 \times \dots \times W_n$  and  $\Psi(\underline{\omega})$  be a IV-CFS over  $\mathbb{U}$  for all  $\underline{\omega} = (b_1, b_2, b_3, \dots, b_n) \in W$ . Then, *interval-valued complex fuzzy hypersoft set* (IV-CFHS)  $\Omega_W = (\Psi, W)$  over  $\mathbb{U}$  is defined as

$$\Omega_W = \{(\underline{\omega}, \Psi(\underline{\omega})) : \underline{\omega} \in W, \Psi(\underline{\omega}) \in C_{IV}(\mathbb{U})\}$$

where

$$\Psi : W \rightarrow C_{IV}(\mathbb{U}), \quad \Psi(\underline{\omega}) = \emptyset \text{ if } \underline{\omega} \notin W.$$

is a IV-CF approximate function of  $\Omega_W$  and  $\Psi(\underline{\omega}) = (\overleftarrow{\Psi}(\underline{\omega}), \overrightarrow{\Psi}(\underline{\omega}))$ .  $\overleftarrow{\Psi}(\underline{\omega}) = \overleftarrow{r} e^{i\overleftarrow{\theta}}$  and  $\overrightarrow{\Psi}(\underline{\omega}) = \overrightarrow{r} e^{i\overrightarrow{\theta}}$  are lower and upper bounds of the membership function of  $\Omega_W$  respectively and its value  $\Psi(\underline{\omega})$  is called  $\underline{\omega}$ -member of IV-CFHS  $\forall \underline{\omega} \in W$ .

**Example 4.2.** Considering example 2.9 with  $W = \{e_1, e_2, e_3, \dots, e_8\}$ , IV-Complex fuzzy sets  $\Psi_W(e_1), \Psi_W(e_2), \dots, \Psi_W(e_8)$  are defined as,

$$\begin{aligned} \Psi_W(e_1) &= \left\{ \frac{[0.4, 0.5]e^{i[0.5, 0.6]\pi}}{t_1}, \frac{[0.7, 0.8]e^{i[0.5, 0.6]\pi}}{t_2}, \frac{[0.6, 0.7]e^{i[0.7, 0.8]\pi}}{t_3}, \frac{[0.3, 0.4]e^{i[0.65, 0.75]\pi}}{t_4} \right\}, \\ \Psi_W(e_2) &= \left\{ \frac{[0.5, 0.6]e^{i[0.6, 0.7]\pi}}{t_1}, \frac{[0.8, 0.9]e^{i[0.8, 0.9]\pi}}{t_2}, \frac{[0.6, 0.7]e^{i[0.8, 0.9]\pi}}{t_3}, \frac{[0.65, 0.75]e^{i[0.85, 0.95]\pi}}{t_4} \right\}, \\ \Psi_W(e_3) &= \left\{ \frac{[0.4, 0.5]e^{i[0.5, 0.6]\pi}}{t_1}, \frac{[0.7, 0.8]e^{i[0.8, 0.9]\pi}}{t_2}, \frac{[0.5, 0.6]e^{i[0.8, 0.9]\pi}}{t_3}, \frac{[0.55, 0.65]e^{i[0.85, 0.95]\pi}}{t_4} \right\}, \\ \Psi_W(e_4) &= \left\{ \frac{[0.2, 0.3]e^{i[0.6, 0.7]\pi}}{t_1}, \frac{[0.6, 0.7]e^{i[0.8, 0.9]\pi}}{t_2}, \frac{[0.4, 0.5]e^{i[0.8, 0.9]\pi}}{t_3}, \frac{[0.65, 0.75]e^{i[0.55, 0.65]\pi}}{t_4} \right\}, \\ \Psi_W(e_5) &= \left\{ \frac{[0.1, 0.2]e^{i[0.4, 0.5]\pi}}{t_1}, \frac{[0.2, 0.3]e^{i[0.7, 0.8]\pi}}{t_2}, \frac{[0.7, 0.8]e^{i[0.6, 0.7]\pi}}{t_3}, \frac{[0.35, 0.45]e^{i[0.55, 0.65]\pi}}{t_4} \right\}, \\ \Psi_W(e_6) &= \left\{ \frac{[0.4, 0.5]e^{i[0.8, 0.9]\pi}}{t_1}, \frac{[0.2, 0.3]e^{i[0.8, 0.9]\pi}}{t_2}, \frac{[0.6, 0.7]e^{i[0.7, 0.8]\pi}}{t_3}, \frac{[0.75, 0.85]e^{i[0.85, 0.95]\pi}}{t_4} \right\}, \\ \Psi_W(e_7) &= \left\{ \frac{[0.5, 0.6]e^{i[0.8, 0.9]\pi}}{t_1}, \frac{[0.8, 0.9]e^{i[0.5, 0.6]\pi}}{t_2}, \frac{[0.4, 0.5]e^{i[0.5, 0.6]\pi}}{t_3}, \frac{[0.75, 0.85]e^{i[0.65, 0.75]\pi}}{t_4} \right\}, \end{aligned}$$

and

$$\Psi_W(e_8) = \left\{ \frac{[0.7, 0.8]e^{i[0.8, 0.9]\pi}}{t_1}, \frac{[0.7, 0.8]e^{i[0.7, 0.8]\pi}}{t_2}, \frac{[0.5, 0.6]e^{i[0.7, 0.8]\pi}}{t_3}, \frac{[0.55, 0.65]e^{i[0.75, 0.85]\pi}}{t_4} \right\}$$

then IV-CFHS  $\Omega_W$  is written by,

$$\Omega_W = \left\{ \begin{array}{l} (e_1, \frac{[0.4,0.5]e^{i[0.5,0.6]\pi}}{t_1}, \frac{[0.7,0.8]e^{i[0.5,0.6]\pi}}{t_2}, \frac{[0.6,0.7]e^{i[0.7,0.8]\pi}}{t_3}, \frac{[0.3,0.4]e^{i[0.65,0.75]\pi}}{t_4}), \\ (e_2, \frac{[0.5,0.6]e^{i[0.6,0.7]\pi}}{t_1}, \frac{[0.8,0.9]e^{i[0.8,0.9]\pi}}{t_2}, \frac{[0.6,0.7]e^{i[0.8,0.9]\pi}}{t_3}, \frac{[0.65,0.75]e^{i[0.85,0.95]\pi}}{t_4}), \\ (e_3, \frac{[0.4,0.5]e^{i[0.5,0.6]\pi}}{t_1}, \frac{[0.7,0.8]e^{i[0.8,0.9]\pi}}{t_2}, \frac{[0.5,0.6]e^{i[0.8,0.9]\pi}}{t_3}, \frac{[0.55,0.65]e^{i[0.85,0.95]\pi}}{t_4}), \\ (e_4, \frac{[0.2,0.3]e^{i[0.6,0.7]\pi}}{t_1}, \frac{[0.6,0.7]e^{i[0.8,0.9]\pi}}{t_2}, \frac{[0.4,0.5]e^{i[0.8,0.9]\pi}}{t_3}, \frac{[0.65,0.75]e^{i[0.55,0.65]\pi}}{t_4}), \\ (e_5, \frac{[0.1,0.2]e^{i[0.4,0.5]\pi}}{t_1}, \frac{[0.2,0.3]e^{i[0.7,0.8]\pi}}{t_2}, \frac{[0.7,0.8]e^{i[0.6,0.7]\pi}}{t_3}, \frac{[0.35,0.45]e^{i[0.55,0.65]\pi}}{t_4}), \\ (e_6, \frac{[0.4,0.5]e^{i[0.8,0.9]\pi}}{t_1}, \frac{[0.2,0.3]e^{i[0.8,0.9]\pi}}{t_2}, \frac{[0.6,0.7]e^{i[0.7,0.8]\pi}}{t_3}, \frac{[0.75,0.85]e^{i[0.85,0.95]\pi}}{t_4}), \\ (e_7, \frac{[0.5,0.6]e^{i[0.8,0.9]\pi}}{t_1}, \frac{[0.8,0.9]e^{i[0.5,0.6]\pi}}{t_2}, \frac{[0.4,0.5]e^{i[0.5,0.6]\pi}}{t_3}, \frac{[0.75,0.85]e^{i[0.65,0.75]\pi}}{t_4}), \\ (e_8, \frac{[0.7,0.8]e^{i[0.8,0.9]\pi}}{t_1}, \frac{[0.7,0.8]e^{i[0.7,0.8]\pi}}{t_2}, \frac{[0.5,0.6]e^{i[0.7,0.8]\pi}}{t_3}, \frac{[0.55,0.65]e^{i[0.75,0.85]\pi}}{t_4}) \end{array} \right\}$$

**Definition 4.3.** Let  $\Omega_{W_1} = (\Psi_1, W_1)$  and  $\Omega_{W_2} = (\Psi_2, W_2)$  be two IV-CFHS over the same  $\mathbb{U}$ .

The set  $\Omega_{W_1} = (\Psi_1, W_1)$  is said to be the *subset* of  $\Omega_{W_2} = (\Psi_2, W_2)$ , if

- i.  $W_1 \subseteq W_2$
- ii.  $\forall \underline{x} \in W_1, \Psi_1(\underline{x}) \subseteq \Psi_2(\underline{x})$  implies  $\overleftarrow{\Psi}_1(\underline{x}) \subseteq \overleftarrow{\Psi}_2(\underline{x}), \overrightarrow{\Psi}_1(\underline{x}) \subseteq \overrightarrow{\Psi}_2(\underline{x})$  i.e.  
 $\overleftarrow{r}_{W_1}(\underline{x}) \leq \overleftarrow{r}_{W_2}(\underline{x}), \overrightarrow{r}_{W_1}(\underline{x}) \leq \overrightarrow{r}_{W_2}(\underline{x}), \overleftarrow{\theta}_{W_1}(\underline{x}) \leq \overleftarrow{\theta}_{W_2}(\underline{x})$  and  $\overrightarrow{\theta}_{W_1}(\underline{x}) \leq \overrightarrow{\theta}_{W_2}(\underline{x})$ ,

where

- $\overleftarrow{r}_{W_1}(\underline{x})$  and  $\overleftarrow{\theta}_{W_1}(\underline{x})$  are amplitude and phase terms of  $\overleftarrow{\Psi}_1(\underline{x})$ ,
- $\overrightarrow{r}_{W_1}(\underline{x})$  and  $\overrightarrow{\theta}_{W_1}(\underline{x})$  are amplitude and phase terms of  $\overrightarrow{\Psi}_1(\underline{x})$ ,
- $\overleftarrow{r}_{W_2}(\underline{x})$  and  $\overleftarrow{\theta}_{W_2}(\underline{x})$  are amplitude and phase terms of  $\overleftarrow{\Psi}_2(\underline{x})$ , and
- $\overrightarrow{r}_{W_2}(\underline{x})$  and  $\overrightarrow{\theta}_{W_2}(\underline{x})$  are amplitude and phase terms of  $\overrightarrow{\Psi}_2(\underline{x})$ .

**Definition 4.4.** Two IV-CFHS  $\Omega_{W_1} = (\Psi_1, W_1)$  and  $\Omega_{W_2} = (\Psi_2, W_2)$  over the same  $\mathbb{U}$ , are said to be *equal* if

- i.  $(\Psi_1, W_1) \subseteq (\Psi_2, W_2)$
- ii.  $(\Psi_2, W_2) \subseteq (\Psi_1, W_1)$ .

**Definition 4.5.** Let  $(\Psi, W)$  be a IV-CFHS over  $\mathbb{U}$ . Then

- i.  $(\Psi, W)$  is called a *null IV-CFHS*, denoted by  $(\Psi, W)_\Phi$  if for all  $\underline{x} \in W$ , the amplitude and phase terms of the membership function are given by  $\overleftarrow{r}_W(\underline{x}) = \overrightarrow{r}_W(\underline{x}) = 0$  and  $\overleftarrow{\theta}_W(\underline{x}) = \overrightarrow{\theta}_W(\underline{x}) = 0\pi$  respectively.
- ii.  $(\Psi, W)$  is called a *absolute IV-CFHS*, denoted by  $(\Psi, W)_\Delta$  if for all  $\underline{x} \in W$ , the amplitude and phase terms of the membership function are given by  $\overleftarrow{r}_W(\underline{x}) = \overrightarrow{r}_W(\underline{x}) = 1$  and  $\overleftarrow{\theta}_W(\underline{x}) = \overrightarrow{\theta}_W(\underline{x}) = 2\pi$  respectively.

**Definition 4.6.** Let  $(\Psi_1, W_1)$  and  $(\Psi_2, W_2)$  are two CFH-sets over the same universe  $\mathbb{U}$ . Then

- i. A IV-CFHS  $(\Psi_1, W_1)$  is called a *homogeneous IV-CFHS*, denoted by  $(\Psi_1, W_1)_{Hom}$  if and only if  $\Psi_1(\underline{x})$  is a homogeneous CF-set for all  $\underline{x} \in W_1$ .

ii. A IV-CFHS  $(\Psi_1, W_1)$  is called a *completely homogeneous IV-CFHS*, denoted by  $(\Psi_1, W_1)_{CHom}$  if and only if  $\Psi_1(\underline{x})$  is a homogeneous with  $\Psi_1(\underline{y})$  for all  $\underline{x}, \underline{y} \in W_1$ .

iii. A IV-CFHS  $(\Psi_1, W_1)$  is said to be a completely homogeneous IV-CFHS with  $(\Psi_2, W_2)$  if and only if  $\Psi_1(\underline{x})$  is a homogeneous with  $\Psi_2(\underline{x})$  for all  $\underline{x} \in W_1 \coprod W_2$ .

#### 4.1. Set Theoretic Operations and Laws on IV-CFHS

Here some basic set theoretic operations (i.e. complement, union and intersection) and laws (commutative laws, associative laws etc.) are discussed on IV-CFHS.

**Definition 4.7.** The *complement* of IV-CFHS  $(\Psi, W)$ , denoted by  $(\Psi, W)^c$  is defined as

$$(\Psi, W)^c = \{(\underline{x}, (\Psi(\underline{x}))^c) : \underline{x} \in W, (\Psi(\underline{x}))^c \in C_{IV}(\mathbb{U})\}$$

such that the amplitude and phase terms of the membership function  $(\Psi(\underline{x}))^c$  are given by

$$(\overleftarrow{r}_W(\underline{x}))^c = 1 - \overleftarrow{r}_W(\underline{x})$$

$$(\overrightarrow{r}_W(\underline{x}))^c = 1 - \overrightarrow{r}_W(\underline{x})$$

and

$$(\overleftarrow{\theta}_W(\underline{x}))^c = 2\pi - \overleftarrow{\theta}_W(\underline{x}),$$

$$(\overrightarrow{\theta}_W(\underline{x}))^c = 2\pi - \overrightarrow{\theta}_W(\underline{x}) \text{ respectively.}$$

**Proposition 4.8.** Let  $(\Psi, W)$  be a IV-CFHS over  $\mathbb{U}$ . Then  $((\Psi, W)^c)^c = (\Psi, W)$ .

*Proof.* Since  $\Psi(\underline{x}) \in C_{IV}(\mathbb{U})$ , therefore  $(\Psi, W)$  can be written in terms of its amplitude and phase terms as

$$(\Psi, W) = \left\{ \left( \underline{x}, \left( \overleftarrow{r}_W(\underline{x})e^{i\overleftarrow{\theta}_W(\underline{x})}, \overrightarrow{r}_W(\underline{x})e^{i\overrightarrow{\theta}_G(\underline{x})} \right) \right) : \underline{x} \in W \right\} \quad (3)$$

Now

$$\begin{aligned} (\Psi, W)^c(\underline{x}) &= \left\{ \left( \underline{x}, \left( (\overleftarrow{r}_W(\underline{x}))^c e^{i(\overleftarrow{\theta}_W(\underline{x}))^c}, (\overrightarrow{r}_W(\underline{x}))^c e^{i(\overrightarrow{\theta}_G(\underline{x}))^c} \right) \right) : \underline{x} \in W \right\} \\ (\Psi, W)^c(\underline{x}) &= \left\{ \left( \underline{x}, \left( (1 - \overleftarrow{r}_W(\underline{x}))e^{i(2\pi - \overleftarrow{\theta}_W(\underline{x}))}, (1 - \overrightarrow{r}_W(\underline{x}))e^{i(2\pi - \overrightarrow{\theta}_G(\underline{x}))} \right) \right) : \underline{x} \in W \right\} \\ ((\Psi, G)^c)^c &= \left\{ \left( \underline{x}, \left( (1 - \overleftarrow{r}_W(\underline{x}))^c e^{i(2\pi - \overleftarrow{\theta}_W(\underline{x}))^c}, (1 - \overrightarrow{r}_W(\underline{x}))^c e^{i(2\pi - \overrightarrow{\theta}_G(\underline{x}))^c} \right) \right) : \underline{x} \in W \right\} \\ ((\Psi, W)^c)^c &= \left\{ \left( \underline{x}, \left( (1 - (1 - \overleftarrow{r}_W(\underline{x})))e^{i(2\pi - (2\pi - \overleftarrow{\theta}_W(\underline{x})))}, (1 - (1 - \overrightarrow{r}_W(\underline{x})))e^{i(2\pi - (2\pi - \overrightarrow{\theta}_G(\underline{x})))} \right) \right) : \underline{x} \in W \right\} \\ ((\Psi, W)^c)^c &= \left\{ \left( \underline{x}, \left( \overleftarrow{r}_W(\underline{x})e^{i\overleftarrow{\theta}_W(\underline{x})}, \overrightarrow{r}_W(\underline{x})e^{i\overrightarrow{\theta}_G(\underline{x})} \right) \right) : \underline{x} \in W \right\} \end{aligned} \quad (4)$$

from equations (3) and (4), we have  $((\Psi, W)^c)^c = (\Psi, W)$ .  $\square$

**Proposition 4.9.** Let  $(\Psi, W)$  be a IV-CFHS over  $\mathbb{U}$ . Then

$$\text{i. } ((\Psi, W)_\Phi)^c = (\Psi, W)_\Delta$$

$$\text{ii. } ((\Psi, W)_\Delta)^c = (\Psi, W)_\Phi$$

**Definition 4.10.** The *intersection* of two IV-CFHS  $(\Psi_1, W_1)$  and  $(\Psi_2, W_2)$  over the same universe  $\mathbb{U}$ , denoted by  $(\Psi_1, W_1) \prod (\Psi_2, W_2)$ , is the IV-CFHS  $(\Psi_3, W_3)$ , where  $W_3 = W_1 \prod W_2$ , and for all  $\underline{x} \in W_3$ ,

$$\overleftarrow{\Psi}_3(\underline{x}) = \begin{cases} \overleftarrow{r}_{W_1}(\underline{x})e^{i\overleftarrow{\theta}_{W_1}(\underline{x})} & , if \underline{x} \in W_1 \setminus W_2 \\ \overleftarrow{r}_{W_2}(\underline{x})e^{i\overleftarrow{\theta}_{W_2}(\underline{x})} & , if \underline{x} \in W_2 \setminus W_1 \\ \min[\overleftarrow{r}_{W_1}(\underline{x}), \overleftarrow{r}_{W_2}(\underline{x})]e^{i\min[\overleftarrow{\theta}_{W_1}(\underline{x}), \overleftarrow{\theta}_{W_2}(\underline{x})]} & , if \underline{x} \in W_1 \prod W_2 \end{cases}$$

and

$$\overrightarrow{\Psi}_3(\underline{x}) = \begin{cases} \overrightarrow{r}_{W_1}(\underline{x})e^{i\overrightarrow{\theta}_{W_1}(\underline{x})} & , if \underline{x} \in W_1 \setminus W_2 \\ \overrightarrow{r}_{W_2}(\underline{x})e^{i\overrightarrow{\theta}_{W_2}(\underline{x})} & , if \underline{x} \in W_2 \setminus W_1 \\ \min[\overrightarrow{r}_{W_1}(\underline{x}), \overrightarrow{r}_{W_2}(\underline{x})]e^{i\min[\overrightarrow{\theta}_{W_1}(\underline{x}), \overrightarrow{\theta}_{W_2}(\underline{x})]} & , if \underline{x} \in W_1 \prod W_2 \end{cases}$$

**Definition 4.11.** The *difference* between two IV-CFHS  $(\Psi_1, W_1)$  and  $(\Psi_2, W_2)$  is defined as

$$(\Psi_1, W_1) \setminus (\Psi_2, W_2) = (\Psi_1, W_1) \prod (\Psi_2, W_2)^c$$

**Definition 4.12.** The *union* of two IV-CFHS  $(\Psi_1, W_1)$  and  $(\Psi_2, W_2)$  over the same universe  $\mathbb{U}$ , denoted by  $(\Psi_1, W_1) \coprod (\Psi_2, W_2)$ , is the IV-CFHS  $(\Psi_3, W_3)$ , where  $W_3 = W_1 \coprod W_2$ , and for all  $\underline{x} \in W_3$ ,

$$\overleftarrow{\Psi}_3(\underline{x}) = \begin{cases} \overleftarrow{r}_{W_1}(\underline{x})e^{i\overleftarrow{\theta}_{W_1}(\underline{x})} & , if \underline{x} \in W_1 \setminus W_2 \\ \overleftarrow{r}_{W_2}(\underline{x})e^{i\overleftarrow{\theta}_{W_2}(\underline{x})} & , if \underline{x} \in W_2 \setminus W_1 \\ \max[\overleftarrow{r}_{W_1}(\underline{x}), \overleftarrow{r}_{W_2}(\underline{x})]e^{i\max[\overleftarrow{\theta}_{W_1}(\underline{x}), \overleftarrow{\theta}_{W_2}(\underline{x})]} & , if \underline{x} \in W_1 \prod W_2 \end{cases}$$

and

$$\overrightarrow{\Psi}_3(\underline{x}) = \begin{cases} \overrightarrow{r}_{W_1}(\underline{x})e^{i\overrightarrow{\theta}_{W_1}(\underline{x})} & , if \underline{x} \in W_1 \setminus W_2 \\ \overrightarrow{r}_{W_2}(\underline{x})e^{i\overrightarrow{\theta}_{W_2}(\underline{x})} & , if \underline{x} \in W_2 \setminus W_1 \\ \max[\overrightarrow{r}_{W_1}(\underline{x}), \overrightarrow{r}_{W_2}(\underline{x})]e^{i\max[\overrightarrow{\theta}_{W_1}(\underline{x}), \overrightarrow{\theta}_{W_2}(\underline{x})]} & , if \underline{x} \in W_1 \prod W_2 \end{cases}$$

**Proposition 4.13.** Let  $(\Psi, W)$  be a IV-CFHS over  $\mathbb{U}$ . Then the following results hold true:

- i.  $(\Psi, W) \prod (\Psi, W)_\Phi = (\Psi, W)$
- ii.  $(\Psi, W) \prod (\Psi, W)_\Delta = (\Psi, W)_\Delta$
- iii.  $(\Psi, W) \prod (\Psi, W)_\Phi = (\Psi, W)_\Phi$
- iv.  $(\Psi, W) \prod (\Psi, W)_\Delta = (\Psi, W)$
- v.  $(\Psi, W)_\Phi \prod (\Psi, W)_\Delta = (\Psi, W)_\Delta$
- vi.  $(\Psi, W)_\Phi \prod (\Psi, W)_\Delta = (\Psi, W)_\Phi$

**Proposition 4.14.** Let  $(\Psi_1, W_1)$ ,  $(\Psi_2, W_2)$  and  $(\Psi_3, W_3)$  are three CFH-sets over the same universe  $\mathbb{U}$ . Then the following commutative and associative laws hold true:

- i.  $(\Psi_1, W_1) \prod (\Psi_2, W_2) = (\Psi_2, W_2) \prod (\Psi_1, W_1)$
- ii.  $(\Psi_1, W_1) \coprod (\Psi_2, W_2) = (\Psi_2, W_2) \coprod (\Psi_1, W_1)$
- iii.  $(\Psi_1, W_1) \prod ((\Psi_2, W_2) \prod (\Psi_3, W_3)) = ((\Psi_1, W_1) \prod (\Psi_2, W_2)) \prod (\Psi_3, W_3)$

$$\text{iv. } (\Psi_1, W_1) \coprod ((\Psi_2, W_2) \coprod (\Psi_3, W_3)) = ((\Psi_1, W_1) \coprod (\Psi_2, W_2)) \coprod (\Psi_3, W_3)$$

**Proposition 4.15.** *Let  $(\Psi_1, W_1)$  and  $(\Psi_2, W_2)$  are two CFH-sets over the same universe  $\mathbb{U}$ . Then the following De Morgans laws hold true:*

- i.  $((\Psi_1, W_1) \coprod (\Psi_2, W_2))^c = (\Psi_1, W_1)^c \coprod (\Psi_2, W_2)^c$
- ii.  $(\Psi_1, W_1) \coprod ((\Psi_2, W_2))^c = (\Psi_1, W_1)^c \coprod (\Psi_2, W_2)^c$

## Conclusion

In this work, the complex fuzzy hypersoft sets (CFH-sets) are developed along with some fundamentals, theoretic set operations and aggregations. Also a method is proposed to solve decision making problems and demonstrated with a commerce-based application. Moreover, the rudiments of interval-valued fuzzy hypersoft set (IV-CFHS) are characterized with suitable examples. CFH-sets and IV-CFHS generalize the existing structures of complex fuzzy soft set, permit a broad range of values for membership function by expanding them to the unit circle in a complex plane, consider the periodic nature of the information through the phase-terms and classify distinct attributes into corresponding attribute-values sets for vivid understanding. Further work may include:

- (i) the extension of proposed work to the development of:
  - complex intuitionistic fuzzy hypersoft set,
  - complex neutrosophic hypersoft set,
  - interval-valued complex intuitionistic fuzzy hypersoft set,
  - interval-valued complex neutrosophic hypersoft set,
- (ii) the application of proposed work in multi-criteria decision-making,
- (iii) the determination of similarity measures and entropies for proposed structures,
- (iv) the parameterization of proposed structures with fuzzy, intuitionistic fuzzy and neutrosophic settings,
- (v) the characterization of proposed structures under multi-decisive environment,
- (vi) the introduction of refinement in the proposed structures for sub-membership grades.

## References

- [1] Ramot, D.; Milo, R.; Friedman, M.; Kandel, A. Complex fuzzy sets. *IEEE Transactions on Fuzzy Systems* **2002**, *10*(2), 171-186. <http://doi.org/10.1109/91.995119>
- [2] Zadeh, L. Fuzzy sets. *Information and control* **1965**, *8*(3), 338-353. [http://doi.org/10.1016/S0019-9958\(65\)90241-X](http://doi.org/10.1016/S0019-9958(65)90241-X)
- [3] Buckley. J.J., Fuzzy complex numbers. *Fuzzy Sets and Systems* **1989**, *33*(3), 333-345. [https://doi.org/10.1016/0165-0114\(89\)90122-X](https://doi.org/10.1016/0165-0114(89)90122-X)
- [4] Zhang, G.; Dillon, T. S.; Cai, K. Y.; Ma, J.; Lu, J. Operation properties and  $\delta$ -equalities of complex fuzzy sets. *International journal of approximate reasoning* **2009**, *50*(8), 1227-1249. <https://doi.org/10.1016/j.ijar.2009.05.010>

---

Atiqe Ur Rahman, Muhammad Saeed, Arooj Khalid, Muhammad Rayees Ahmad and Sehrish Ayaz, Decision-Making Application Based on Aggregations of Complex Fuzzy Hypersoft Set and Development of Interval-Valued Complex Fuzzy Hypersoft Set

- [5] Ramot, D.; Milo, R.; Friedman, M.; Kandel, A. Complex fuzzy sets. *IEEE Transactions on Fuzzy Systems* **2002**, 10(2), 171-186. <https://doi.org/10.1109/91.995119>
- [6] Ramot, D.; Friedman, M.; Langholz, G.; Kandel, A. Complex fuzzy logic. *IEEE Transactions on Fuzzy Systems* **2003**, 11(4), 450-461. <https://doi.org/10.1109/TFUZZ.2003.814832>
- [7] Molodtsov, D. Soft Set Theory - First Results. *Computers and Mathematics with Applications* **1999**, 37, 19-31. [http://doi.org/10.1016/S0898-1221\(99\)00056-5](http://doi.org/10.1016/S0898-1221(99)00056-5)
- [8] Maji, P.K.; Biswas R.; Roy, A. R. Soft Set Theory. *Computers and Mathematics with Applications* **2003**, 45, 555-562. [http://doi.org/10.1016/S0898-1221\(03\)00016-6](http://doi.org/10.1016/S0898-1221(03)00016-6)
- [9] Maji, P. K.; Biswas, R.; Roy, A. R. Fuzzy soft sets. *Journal of Fuzzy Mathematics* **2001**, 9(3), 589-602.
- [10] Pei, D.; Miao, D. From soft set to information system. *In international conference of granular computing IEEE* **2005**, 2 , 617-621. <http://doi.org/10.1109/GRC.2005.1547365>
- [11] Ali, M. I.; Feng, F.; Liu, X.; Min, W. K.; Sabir, M. On some new operations in soft set theory. *Computers and Mathematics with Applications* **2009**, 57 , 1547-1553. <http://doi.org/10.1016/j.camwa.2008.11.009>
- [12] Babitha, K. V.; Sunil, J. J. Soft set relations and functions. *Computers and Mathematics with Applications* **2010**, 60, 1840-1849. <http://doi.org/10.1016/j.camwa.2010.07.014>
- [13] Babitha, K. V.; Sunil, J. J. Transitive closure and ordering in soft set. *Computers and Mathematics with Applications* **2010**, 61 2235-2239. <http://doi.org/10.1016/j.camwa.2011.07.010>
- [14] Sezgin, A.; Atagün A. O. On operations of soft sets. *Computers and Mathematics with Applications* **2011**, 61(5), 1457-1467. <http://doi.org/10.1016/j.camwa.2011.01.018>
- [15] Ge, X.; Yang, S. Investigations on some operations of soft sets. *World Academy of Science Engineering and Technology* **2011**, 75, 1113-1116.
- [16] Li, F. Notes on soft set operations. *ARPJ Journal of systems and softwares* **2011**, 205-208.
- [17] Maji, P. K.; Biswas, R.; Roy, A. R. Intuitionistic fuzzy soft sets. *The Journal of Fuzzy Mathematics* **2001**, 9(3), 677-692.
- [18] Maji, P. K. Neutrosophic Soft Set. *Annals of Fuzzy Mathematics and Informatics* **2013**, 5(1), 2287-623.
- [19] Smarandache, F. Extension of Soft Set of Hypersoft Set, and then to Plithogenic Hypersoft Set. *Neutrosophic Sets and Systems* **2018**, 22 , 168-170. <http://doi.org/10.5281/zenodo.2838716>
- [20] Saeed, M.; Ahsan, M.; Siddique, M.k.; Ahmad, M.R. A Study of the Fundamentals of Hypersoft Set Theory. *International Journal of Scientific and Engineering Research* **2020**, 11(1), 320-329.
- [21] Saeed, M.; Rahman, A. U.; Ahsan, M.; Smarandache, F. An Inclusive Study on Fundamentals of Hypersoft Set, In Theory and Application of Hypersoft Set, Pons Publishing House, Brussel, 2021, 1-23.
- [22] Abbas, F.; Murtaza, G.; Smarandache, F. (2020). Basic operations on hypersoft sets and hypersoft points. *Neutrosophic Sets and Systems* **2020**, 35, 407-421. <http://doi.org/10.5281/zenodo.3951694>
- [23] Saqlain, M.; Jafar, N.; Moin, S.; Saeed, M.; Broumi, S. Single and Multi-valued Neutrosophic Hypersoft Set and Tangent Similarity Measure of Single valued Neutrosophic Hypersoft Sets. *Neutrosophic Sets and Systems* **2020**, 32, 317-329. <http://doi.org/10.5281/zenodo.3723165>
- [24] Saqlain, M.; Moin, S.; Jafar, N.; Saeed, M.; Smarandache, F. Aggregate Operators of Neutrosophic Hypersoft Sets. *Neutrosophic Sets and Systems* **2020**, 32, 294-306. <http://doi.org/10.5281/zenodo.3723155>
- [25] Saqlain, M.; Saeed, M.; Ahmad, M.R.; Smarandache, F. Generalization of TOPSIS for Neutrosophic Hypersoft Sets using Accuracy Function and its Application. *Neutrosophic Sets and Systems* **2020**, 27, 131-137. <http://doi.org/10.5281/zenodo.3275533>
- [26] Martin, N.; Smarandache, F. Concentric Plithogenic Hypergraph based on Plithogenic Hypersoft Sets A Novel Outlook. *Neutrosophic Sets and Systems* **2020**, 33, 78-91. <http://doi.org/10.5281/zenodo.3782824>
- [27] Rahman, A. U.; Saeed, M.; Alodhaibi, S.S.; Abd, H. Decision Making Algorithmic Approaches Based on Parameterization of Neutrosophic Set under Hypersoft Set Environment with Fuzzy, Intuitionistic Fuzzy

- and Neutrosophic Settings. *CMES-Computer Modeling in Engineering & Sciences* **2021**, 128(2), 743-777. <http://doi.org/10.32604/cmes.2021.016736>
- [28] Rahman, A. U.; Saeed, M.; Smarandache, F.; Ahmad, M. R. Development of Hybrids of Hypersoft Set with Complex Fuzzy Set, Complex Intuitionistic Fuzzy set and Complex Neutrosophic Set. *Neutrosophic Sets and Systems* **2020**, 38, 335-354. <http://doi.org/10.5281/zenodo.4300520>
- [29] Rahman, A. U.; Saeed, M.; Smarandache, F. Convex and Concave Hypersoft Sets with Some Properties. *Neutrosophic Sets and Systems* **2020**, 38, 497-508. <http://doi.org/10.5281/zenodo.4300580>
- [30] Deli, I. Hybrid set structures under uncertainly parameterized hypersoft sets, Theory and applications. In Theory and Application of Hypersoft Set, Pons Publishing House, Brussel, 2021, 24-49.
- [31] Gayen, S.; Smarandache, F.; Jha, S.; Singh, M. K.; Broumi, S.; Kumar, R. Introduction to Plithogenic Hypersoft Subgroup. *Neutrosophic Sets and Systems* **2020**, 33, 14. <http://doi.org/10.5281/zenodo.3782897>
- [32] Saeed, M.; Ahsan, M.; Saeed, M. H.; Mehmood, A.; Abdeljawad, T. An Application of Neutrosophic Hypersoft Mapping to Diagnose Hepatitis and Propose Appropriate Treatment. *IEEE Access* **2021**, 9, 70455-70471. <http://doi.org/10.1109/ACCESS.2021.3077867>
- [33] Saeed, M.; Ahsan, M.; Abdeljawad, T. A Development of Complex Multi-Fuzzy Hypersoft Set With Application in MCDM Based on Entropy and Similarity Measure. *IEEE Access* **2021**, 9, 60026-60042. [10.1109/ACCESS.2021.3073206](http://doi.org/10.1109/ACCESS.2021.3073206)
- [34] Rahman, A.U.; Saeed, M.; Dhital, A. Decision Making Application Based on Neutrosophic Parameterized Hypersoft Set Theory. *Neutrosophic Sets and Systems* **2021**, 41, 1-14. <http://doi.org/10.5281/zenodo.4625665>
- [35] Rahman, A.U.; Saeed, M.; Zahid, S. Application in Decision Making Based on Fuzzy Parameterized Hypersoft Set Theory. *Asia Mathematica* **2021**, 5(1), 19-27. <http://doi.org/10.5281/zenodo.4721481>
- [36] Rahman, A.U.; Hafeez, A.; Saeed, M.; Ahmad, M.R.; Farwa, U. Development of Rough Hypersoft Set with Application in Decision Making for the Best Choice of Chemical Material, In Theory and Application of Hypersoft Set, Pons Publication House, Brussel, 2021, 192-202. <http://doi.org/10.5281/zenodo.4743367>
- [37] Ihsan, M.; Rahman, A.U.; Saeed, M. Hypersoft Expert Set With Application in Decision Making for Recruitment Process. *Neutrosophic Sets and Systems* **2021**, 42, 191-207. <http://doi.org/10.5281/zenodo.4711524>
- [38] Thirunavukarasu, P.; Suresh, R.; Ashokkumar, V. Theory of Complex Fuzzy Soft Set and its Applications. *International Journal for Innovative Research in Science and Technology* **2013**, 3, 13-18.
- [39] Kumar, T.; Bajaj, R. K. On Complex Intuitionistic Fuzzy Soft Sets with Distance Measures and Entropies. *Journal of Mathematics* **2014**, 2014(3), 1-12. <http://doi.org/10.1155/2014/972198>
- [40] Selvachandran, G.; Singh, P. K. Interval-valued complex fuzzy soft set and its application. *International Journal for Uncertainty Quantification* **2018**, 8(2), 101-117. <http://doi.org/10.1615/Int.J.UncertaintyQuantification.2018020362>
- [41] Abd Ulzeez M. J. S. Alkouri. Complex Generalised Fuzzy Soft Sets and its Application. *WSEAS Transactions on Mathematics* **2020**, 19, 323-333. <http://doi.org/10.37394/23206.2020.19.32>

Received: May 5, 2021. Accepted: October 5, 2021