

Decision Making in Neutrosophic Environment

MICHAEL GR. VOSKOGLLOU

Mathematical Sciences, School of Technological Applications

University of Peloponnese (ex T.E.I. of Western Greece)

Meg. Alexandrou 1, 26334 Patras

GREECE

Abstract: - Frequently in real life situations decision making takes place under fuzzy conditions, because the corresponding goals and/or the existing constraints are not clearly defined. Maji et al. introduced a method of parametric decision making using soft sets as tools. In reality, however, cases appear in which the decision maker has doubts about the correctness of the parameters assigned to some or all of the elements of the set of the discourse. In order to study more effectively the decision making process in such cases, we introduce neutrosophic sets here and we replace the binary elements in the tabular matrix of the method of Maji et al. by neutrosophic triplets. Our method is illustrated by suitable examples.

Key-Words: - Decision making (DM), fuzzy set (FS), fuzzy logic (FL), grey number (GN), soft set (SS), neutrosophic set (NS), neutrosophic triplets.

Received: August 28, 2022. Revised: November 22, 2022. Accepted: January 7, 2023. Published: February 20, 2023.

1 Introduction

Decision Making (DM) is the cognitive process of selecting, with the help of the suitable criteria, the best option among the set of all possible alternatives. DM plays a fundamental role in a great variety of human activities and many books have been written advising people how to make smarter choices easier and quicker; e.g. [1, 2], etc. The decision-maker's environment seriously affects the DM process. Frequently, for example, in real life situations DM takes place under fuzzy conditions, because the corresponding goals and/or the existing constraints are not clearly defined. Several methods have been proposed for successful DM in such cases, e.g. [3-5], etc.

Maji et al. [6] introduced in 2002 a method of parametric DM using *soft sets (SSs)* as tools and representing their tabular form as a binary matrix. When some or all of the parameters used for the characterization of the elements of the set of the discourse (houses in their example) are of fuzzy texture (beautiful and cheap in their example), however, their method does not always give the best solution to the corresponding DM problem. This happens, because they replace the parameters in the tabular form of the corresponding SS with the binary elements 0, 1. In other words, their method, although it starts with a fuzzy framework (SSs), it continues by using bivalent logic for obtaining the

required decision (beautiful or not beautiful and cheap or not cheap in their example)! In order to tackle this problem, we modified in earlier works the method of Maji et al. by using either *triangular fuzzy numbers (TFNs)* [7] or *grey numbers (GNs)* – see [8] and [9] (section 5.3) – instead of the binary elements in the tabular form of the corresponding SS.

In reality, however, cases appear in which the decision maker has doubts about the correctness of the parameters assigned to some or all of the elements of the set of the discourse (e.g. very beautiful, rather beautiful, etc.). In order to study more effectively the DM process in such cases, we introduce here *neutrosophic sets (NSs)* and we replace the binary elements of the method of Maji et al. by *neutrosophic triplets*.

The rest of the paper is organized as follows: Section 2 contains the necessary background about NSs, SSs and GNs needed for the understanding of the paper. The DM method of Maji et al., our modification using GNs, as well as the new neutrosophic DM method, are developed in section 3, illustrated by suitable examples. The article closes with a discussion of the results obtained, including some hints for further research, and the final conclusions, which are presented in section 4.

2 Mathematical Background

2.1 Fuzzy Sets and Fuzzy Logic

Zadeh extended in 1965 the concept of crisp set to the concept of *fuzzy set (FS)* [10] on the purpose of dealing with the existing in real life partial truths (e.g. rather good, almost true, etc.) and of expressing mathematically definitions having no clear boundaries (e.g. high mountains, clever people, etc.) This was succeeded by replacing the characteristic function of crisp sets by the *membership function*, which maps each element of the universal set U to the unit interval $[0, 1]$. If A is the corresponding FS in U and $m: U \rightarrow [0, 1]$ is its membership function, $m(x)$ is called the *membership degree* of x in A , for all elements x in U . The closer $m(x)$ to 1, the better x satisfies the characteristic property of A . Although the FS A is typically defined as the set of all ordered pairs of the form $(x, m(x))$, $\forall x \in U$, for reasons of simplicity many authors identify A with its membership function.

Most notions and operations on crisp sets are extended in a natural way to FSs (e.g. see [11]). Based on the concept of FS, Zadeh introduced the infinite-valued *fuzzy logic (FL)*, in which the truth values are modelled by numbers in the unit interval [12]. FL does not oppose the traditional bivalent logic of Aristotle (384-322 BC), which used to be for many centuries the basic tool of human reasoning being “responsible” for the growth of science and human civilization; on the contrary it completes and extends it [13]

When membership functions were reinterpreted as *possibility distributions*, FSs and FL were used to embrace *uncertainty* modelling [14, 15]. The frequently appearing in real life uncertainty is due to several reasons, like randomness, imprecise or incomplete data, vague information, etc. *Probability* theory has been proved sufficient to tackle only the cases of uncertainty which are due to randomness [16]. Starting from Zadeh’s FSs, however, several theories have been developed during the last years on the purpose of tackling more effectively all the forms of the existing uncertainty. The main among those theories are briefly reviewed in [17]. In the present paper we are going to use elements from the theories of NSs, SSs and GNs needed for its understanding, which are exposed below.

2.2 Neutrosophic Sets

Atanassov in 1986, considered, in addition to Zadeh’s membership degree, the degree of *non-membership* and extended FS to the notion of

intuitionistic FS (IFS) [18]. Smarandache in 1995, inspired by the frequently appearing in real life neutralities - like <friend, neutral, enemy>, <win, draw, defeat>, <high, medium, short>, etc. - generalized IFS to the concept of NS by adding the degree of *indeterminacy* or *neutrality* [19]. The word “neutrosophy” is a synthesis of the word “neutral” and the Greek word “sophia” (wisdom) and means “the knowledge of neutral thought”. The simplest form of a NS is defined as follows:

Definition 1: A *single valued NS (SVNS)* A in the universe U is of the form

$$A = \{(x, T(x), I(x), F(x)): x \in U, T(x), I(x), F(x) \in [0, 1], 0 \leq T(x) + I(x) + F(x) \leq 3\} \quad (1)$$

In equation (1) $T(x)$, $I(x)$, $F(x)$ are the degrees of *truth* (or membership), *indeterminacy* (or neutrality) and *falsity* (or non-membership) of x in A respectively, called the *neutrosophic components* of x . For simplicity, we write $A \langle T, I, F \rangle$.

Indeterminacy is defined to be in general everything that exists between the opposites of truth and falsity [20].

Example 1: Let U be the set of the players of a soccer club and let A be the SVNS of the good players of the club. Then each player x is characterized by a *neutrosophic triplet* (t, i, f) with respect to A , with t, i, f in $[0, 1]$. For example, $x(0.7, 0.1, 0.4) \in A$ means that the coach of the club is 70% sure that x is a good player, but at the same time has a 10% doubt about it and a 40% belief that x is not a good player. In particular, $x(0, 1, 0) \in A$ means that the coach does not know absolutely nothing about player x (new player).

If the sum $T(x) + I(x) + F(x) < 1$, then it leaves room for incomplete information about x , if it is equal to 1 for complete information and if it is >1 for *inconsistent* (i.e. contradiction tolerant) information about x . A SVNS may contain simultaneously elements leaving room to all the previous types of information. All notions and operations defined on FSs are naturally extended to SVNSs [21].

Summation of neutrosophic triplets is equivalent to the union of NSs. That is why the neutrosophic summation and implicitly its extension to neutrosophic scalar multiplication can be defined in many ways, equivalently to the known in the literature neutrosophic union operators [22]. For the needs of the present work, writing the elements of a SVNS A in the form of neutrosophic triplets and considering them simply as ordered triplets we define addition and scalar product as follows:

Definition 2: Let $(t_1, i_1, f_1), (t_2, i_2, f_2)$ be in A and let k be a positive number. Then:

- The *sum* $(t_1, i_1, f_1) + (t_2, i_2, f_2) = (t_1 + t_2, i_1 + i_2, f_1 + f_2)$ (2)
- The *scalar product* $k(t_1, i_1, f_1) = (kt_1, ki_1, kf_1)$ (3)

Remark 1: The summation and scalar product of the elements of a SVNS A with respect to Definition 2 need not be a closed operation in A, since it may happen that $(t_1 + t_2) + (i_1 + i_2) + (f_1 + f_2) > 3$ or $kt_1 + ki_1 + kf_1 > 3$. With the help of Definition 2, however, one can define in A the *mean value* of a finite number of elements of A as follows:

Definition 3: Let A be a SVNS and let $(t_1, i_1, f_1), (t_2, i_2, f_2), \dots, (t_k, i_k, f_k)$ be a finite number of elements of A. Assume that (t_i, i_i, f_i) appears n_i times in an application, $i = 1, 2, \dots, k$. Set $n = n_1 + n_2 + \dots + n_k$. Then the *mean value* of all these elements of A is defined to be the element of A

$$(t_m, i_m, f_m) = \frac{1}{n} [n_1(t_1, i_1, f_1) + n_2(t_2, i_2, f_2) + \dots + n_k(t_k, i_k, f_k)] \quad (4)$$

2.3 Soft Sets

A disadvantage of FSs and of all their extensions involving membership degrees (like IFSs, NSs, etc.), is that there is not any exact rule for defining properly the corresponding membership functions. The methods used for this are usually statistical or intuitive/empirical. Moreover, the definition of the membership function is not unique depending on the “signals” that each observer receives from the environment. For example, defining the FS of “young people” one could consider all persons aged less than 30 years as young and another one all persons aged less than 40 years. As a result the second observer will assign membership degree 1 to all people aged between 30 and 40 years, whereas the first one will assign to them membership degrees less than 1. Analogous differences are logically expected to appear with the membership degrees of all the other ages. In other words, the only restriction for the definition of the membership function is that it must be compatible to the common sense; otherwise the resulting FS does not give a creditable description of the corresponding real situation. This could happen, for instance, if in the previous example people aged more than 70 years possessed membership degrees ≥ 0.5 .

To overcome this problem, the concept of *interval-valued FS (IVFS)* was introduced in 1975. An IVFS is defined by mapping the universe U to the set of the closed subintervals of $[0, 1]$ [23]. Other related to FS theories were also developed in which the definition of a membership function is not necessary

(*grey systems and numbers* [24]), or it is overpassed by using a pair of crisp sets giving the lower and upper bound of the original set (*rough sets*) [25]. Molodstov, introduced in 1999 the concept of SS for tackling the uncertainty in a parametric manner, not needing, therefore, the definition of a membership function [26]. Namely, a SS is defined as follows:

Definition 4: Let E be a set of parameters, let A be a subset of E, and let f be a map from A into the power set $P(U)$ of the universe U. Then the SS (f, A) in U is defined as the set of the ordered pairs

$$(f, A) = \{(e, f(e)): e \in A\} \quad (5)$$

In other words, a SS is a parametrized family of subsets of U. The name “soft” was given because the form of (f, A) depends on the parameters of A.

Example 2: Let $U = \{H_1, H_2, H_3\}$ be a set of houses and let $E = \{e_1, e_2, e_3\}$ be the set of parameters $e_1 = \text{cheap}$, $e_2 = \text{beautiful}$ and $e_3 = \text{expensive}$. Let us further assume that H_1, H_2 are cheap, H_3 is expensive and H_2, H_3 are beautiful houses. Then, a map $f: E \rightarrow P(U)$ is defined by $f(e_1) = \{H_1, H_2\}$, $f(e_2) = \{H_2, H_3\}$ and $f(e_3) = \{H_3\}$. Therefore, the SS (f, E) in U is the set of the ordered pairs

$$(f, E) = \{(e_1, \{H_1, H_2\}), (e_2, \{H_2, H_3\}), (e_3, \{H_3\})\} \quad (6)$$

Maji et al. [6] introduced a *tabular representation* of SSs in the form of a binary matrix in order to be stored easily in a computer’s memory. For instance, the tabular representation of the soft set (f, E) of Example 2 is given by Table 1.

Table 1. Tabular representation of the SS of Example 2

| | e_1 | e_2 | e_3 |
|-------|-------|-------|-------|
| H_1 | 1 | 0 | 0 |
| H_2 | 1 | 1 | 0 |
| H_3 | 0 | 1 | 1 |

A FS in U with membership function $y = m(x)$ is a SS in U of the form $(f, [0, 1])$, where $f(\alpha) = \{x \in U: m(x) \geq \alpha\}$ is the corresponding *α -cut* of the FS, for each α in $[0, 1]$. Consequently the concept of SS is a generalization of the concept of FS. All notions and operations defined on FSs are extended in a natural way to SSs [27].

2.4 Grey Numbers

The theory of *grey systems* [24] introduces an alternative way for managing the uncertainty in case of approximate data. A grey system is understood to be any system which lacks information, such as structure message, operation mechanism or/and behavior document.

Closed real intervals are used for performing the necessary calculations in grey systems. In fact, a closed real interval $[x, y]$ could be considered as representing a real number T , termed as a GN, whose exact value in $[x, y]$ is unknown. We write then $T \in [x, y]$. A GN T , however, is frequently accompanied by a *whitenization function* $f: [x, y] \rightarrow [0, 1]$, such that, if $f(a)$ approaches 1, then a in $[x, y]$ approaches the unknown value of T . If no whitenization function is defined, it is logical to consider as a representative crisp approximation of the GN A the real number

$$V(A) = \frac{x+y}{2} \quad (7)$$

The arithmetic operations on GNs are introduced with the help of the known arithmetic of the real intervals [28]. In this work we are going to make use only of the addition of GNs and of the scalar multiplication of a GN with a positive number, which are defined as follows:

Definition 5: Let $A \in [x_1, y_1]$, $B \in [x_2, y_2]$ be two GNs and let k be a positive number. Then:

- The sum: $A+B$ is the GN $A+B \in [x_1+y_1, x_2+y_2]$ (8)
- The scalar product kA is the GN $kA \in [kx_1, ky_1]$ (9)

3. The Parametric Assessment Method

The parametric assessment method of Maji et al. [6], our modification using GNs [8, 9] and our new method using NSs are developed in this section through suitable examples concerning the choice of a new player by a soccer club.

3.1 The Method of Maji et al.

Let us consider the following example:

Example 3: A soccer club wants to choose a new player among 6 candidates, say P_1, P_2, P_3, P_4, P_5 and P_6 . The desired qualifications of the new player are to be fast, younger than 30 years, higher than 1.70 m and experienced. Assume that P_1, P_2, P_6 are the fast players, P_2, P_3, P_5, P_6 are the players being younger than 30 years, P_3, P_5 are the players with heights

greater than 1.70 m and P_4 is the unique experienced player. Which is the best choice for the club?

Solution: Let U be the set of the 6 candidate players. Consider the parameters e_1 =fast, e_2 =younger than 30 years, e_3 =higher than 1.70 m, e_4 =experienced and set $E = \{e_1, e_2, e_3, e_4\}$. Define the map $f: E \rightarrow P(U)$ by $f(e_1) = \{P_1, P_2, P_6\}$, $f(e_2) = \{P_2, P_3, P_5, P_6\}$, $f(e_3) = \{P_3, P_5\}$ and $f(e_4) = \{P_4\}$. Then the tabular form of the SS (f, E) is shown in Table 2.

Table 2. Tabular representation of the SS of Example 3

| | e_1 | e_2 | e_3 | e_4 |
|-------|-------|-------|-------|-------|
| P_1 | 1 | 0 | 0 | 0 |
| P_2 | 1 | 1 | 0 | 0 |
| P_3 | 0 | 1 | 1 | 0 |
| P_4 | 0 | 0 | 0 | 1 |
| P_5 | 0 | 1 | 1 | 0 |
| P_6 | 1 | 1 | 0 | 0 |

The *choice value* of each player is calculated by adding the binary elements of the row of Table 2 in which he belongs. The players P_1 and P_4 , therefore, have choice value 1 and all the others have choice value 2. Consequently, the right decision is to choose one of the players P_2, P_3, P_5 , and P_6 .

3.2 Our Method Using GNs

The previous decision, obtained by the method of Maji et al., is obviously not very helpful for the soccer club. Observe, however, that the parameters e_1 and e_4 , in contrast to e_2 and e_3 , have not a bivalent texture. A player, for example, could not be very fast, but quite fast, or rather experienced, etc. This inspired us to characterize the qualifications of the players with respect to the parameters e_1 and e_4 in the tabular matrix of the previous SS (f, E) by using the linguistic grades A =excellent, B =very good, C =good, D =mediocre and F = not satisfactory instead of the binary element 0. To show how one

works in this case for making the right decision, let us modify Example 3 as follows:

Example 4: Reconsider Example 3 and assume that the technical manager of the soccer club, after a more careful inspection of the qualifications of the 6 candidate players, decided to use the following Table 3 instead of Table 2 for the DM process. Which is the best choice for the club in this case?

Table 3. Tabular representation of the SS of Example 4

| | e ₁ | e ₂ | e ₃ | e ₄ |
|----------------|----------------|----------------|----------------|----------------|
| P ₁ | 1 | 0 | 0 | C |
| P ₂ | 1 | 1 | 0 | F |
| P ₃ | C | 1 | 1 | C |
| P ₄ | D | 0 | 0 | 1 |
| P ₅ | D | 1 | 1 | C |
| P ₆ | 1 | 1 | 0 | D |

Solution: Assign to each of the qualitative grades A, B, C, D, F a GN, denoted for simplicity by the same letter, as follows: A = [0.85, 1], B = [0.75, 0.84], C = [0.6, 0.74], D = [0.5, 0.59], F = [0, 0.49]. From Table 3 then, one calculates, with the help of formulas (7), (8) and (9), the choice value of each player in the following way. P₁: $1+V(C) = 1 + \frac{0.6+0.74}{2} = 1.67$, P₂: $2+V(F) = 2.245$, P₃: $2+V(2C) = 3.34$, P₄: $1+V(D) = 1.545$, P₅: $2+V(D+C) = 3.215$, P₆: $2+V(D) = 2.545$. The right decision, therefore, is to choose the player P₃.

Remark 2: The choices of the qualitative grades A, B, C, D, F, as well as of the intervals for translating them in the numerical scale 0-1, correspond to generally accepted standards. The decision maker, however, could use, with respect to his/her goals, more or less qualitative grades (e.g. by adding E between D and F, etc.) and could also change the corresponding intervals, e.g. by setting A = [0.9, 1], B = [0.8, 0.89], C = [0.7, 0.79], D = [0.6, 0.69], F = [0, 0.59], or otherwise. Such changes, however, does not affect the generality of our method

3.3. The Neutrosophic DM Method

As it was already mentioned in our Introduction, DM situations appear frequently in reality, in which

the decision maker has doubts about the correctness of the parameters assigned to some or all of the elements of the set of the discourse. In such cases, the best way to perform the DM process is to use NSs. In Example 4, for instance, considering the set U of the 6 candidate players as the universal set, the decision maker could define in U the NSs of the fast and of the experienced players by assigning the suitable neutrosophic triplets to each player of U. In order to obtain complete information, he/she must arrange to have $t+i+f = 1$, for each triplet (t, i, f). Then, he/she could continue the DM process by replacing in Table 3 the qualitative grades by the corresponding neutrosophic triplets and the binary elements 0, 1 by the neutrosophic triplets (0, 0, 1) and (1, 0, 0) respectively. This process will be illustrated by the following Example 5.

Example 5: Reconsider Example 4 and assume that the technical manager of the soccer club, being not sure about the qualitative grades assigned to each of the 6 candidate players, he decided to proceed by replacing them by neutrosophic triplets, in the way that we have previously described. As a result, the tabular matrix of the DM process took the form of the following Table 4. Which is the best decision for the club in this case?

Table 4. Tabular representation of the SS of Example 5

| | e ₁ | e ₂ | e ₃ | e ₄ |
|----------------|-----------------|----------------|----------------|-----------------|
| P ₁ | (1, 0, 0) | (0, 0, 1) | (0, 0, 1) | (0.6, 0.3, 0.1) |
| P ₂ | (1, 0, 0) | (1, 0, 0) | (0, 0, 1) | (0.2, 0.2, 0.6) |
| P ₃ | (0.5, 0.4, 0.1) | (1, 0, 0) | (1, 0, 0) | (0.6, 0.2, 0.2) |
| P ₄ | (0.5, 0.2, 0.3) | (0, 0, 1) | (0, 0, 1) | (1, 0, 0) |
| P ₅ | (0.5, 0.1, 0.4) | (1, 0, 0) | (1, 0, 0) | (0.6, 0.3, 0.1) |
| P ₆ | (1, 0, 0) | (1, 0, 0) | (0, 0, 1) | (0.4, 0.4, 0.2) |

Solution: The choice value of each player in this case is defined to be the mean value of the neutrosophic triplets of the line of Table 4 in which he belongs. Thus, by equation (4), the choice value of P₁ is equal to $\frac{1}{4}[(1, 0, 0)+2(0, 0, 1)+(0.6, 0.3, 0.1)] = \frac{1}{4}(1.6, 0.3, 2.1) = (0.4, 0.075, 0.525)$. In the same way one finds that the choice values of P₂, P₃,

P_4 , P_5 and P_6 are (0.55, 0.005, 0.4), (0.775, 0.15, 0.075), (0.375, 0.05, 0.575), (0.775, 0.1, 0.125) and (0.6, 0.1, 0.3) respectively.

In this case the club's technical manager could use either an optimistic criterion by choosing the player with the greatest truth degree, or a conservative criterion by choosing the player with the lower falsity degree. Consequently, using the optimistic criterion he must choose one of the players P_3 and P_5 , whereas using the conservative criterion he must choose the player P_3 . A combination of the two criteria leads to the final choice of player P_3 . Observe, however, that, since the indeterminacy degree of P_3 is 0.15 and of P_5 is 0.1, there is a slightly greater doubt about the qualifications of P_3 with respect to the qualifications of P_5 . In other words, the choice of P_3 is connected with a slightly greater risk. In final analysis, therefore, all the neutrosophic components assigned to each player give useful information about his qualifications.

4. Discussion and Conclusions

In this paper a parametric DM method of hybrid character was presented illustrated by suitable examples about the choice of a new player from a football club. The whole discussion performed leads to the following conclusions:

- The parametric DM method is based on the introduction of a set E of parameters characterizing the elements of the set U of all possible solutions of the corresponding DM problem, the definition with respect to E of a suitable SS in U and the use of its tabular representation T as a tool for the DM process.
- When all the parameters of E are of bivalent texture (yes or no), then T takes the form of a binary matrix, wherefrom the decision maker calculates the choice value of each element of U by adding the binary elements of the row of T in which this element appears.
- When some or all of the parameters of E are of fuzzy texture and the decision maker has no doubts about the qualitative characterizations assigned to the elements of U with respect to these parameters, then the binary elements of T corresponding to the fuzzy parameters are replaced by suitable GNs and the choice value of each element of U is calculated by adding the remaining in the corresponding row of T

binary elements and the representative real values of the GNs appearing in it.

- When some or all of the parameters of E are of fuzzy texture and the decision maker has doubts about the qualitative characterizations assigned to the elements of U with respect to these parameters, then each parameter of E is expressed in the form of a NS in U and the binary elements of T are replaced by the corresponding neutrosophic triplets. In this case, the choice value of each element of U is obtained by calculating the mean value of the neutrosophic triplets of the row of T in which this element appears.

As it has been already mentioned in section 2.1 of this paper, several theories extending / generalizing Zadeh's FSs have been developed during the last years on the purpose of tackling more effectively the existing in real life uncertainty. None of these theories, however, can tackle efficiently alone all the forms of the existing uncertainty, but the combination of all of them provides an adequate framework towards this target.

The results obtained in this and earlier works of the present author give strong indications that hybrid methods applied to several situations in fuzzy environments could give better results, not only for DM, as it happened here, but also for assessment - e.g. see [9] (section 5.2) – and probably for many other topics. This is, therefore, a promising area for further research.

References

- [1] Barker, R.C., *The Power of Decision*, Penguin Putman: LA, USA, 2011
- [2] Heath C., Heath, D., *Decisive: How to make better choices in life and work*, Crown Business, NY, USA, 2013
- [3] Bellman, R.A., Zadeh, L.A., Decision Making in Fuzzy Environment, *Management Science*, 17, 1970, pp. 141-164.
- [4] Alcantud, J.C.R. (Ed.), *Fuzzy Techniques for Decision Making* (reprint of the special issue published in *Symmetry*), MDPI, Basel, Switzerland, 2018.
- [5] Zhu, B. & Ren, P., Type-2 fuzzy numbers made simple in decision making, *Fuzzy Optimization and Decision Making*, 21, 2022, pp.175-195.
- [6] Maji, P.K., Roy, A.R., Biswas, R., An Application of Soft Sets in a Decision Making Problem, *Computers and Mathematics with*

- Applications*, 44, 2002, pp. 1077-1083..
- [7] Voskoglou, M.Gr., A Hybrid Method for Decision Making Utilizing TFNs and Soft Sets as Tools, *Equations*, 2, 2022, pp. 65-69.
- [8] Voskoglou, M.Gr., A Combined Use of Soft Sets and Grey Numbers in Decision Making, *Journal of Computational and Cognitive Engineering*, doi: 10.47852/bonviewjce 2202237, 2022.
- [9] Voskoglou, M.Gr., Fuzziness, Indeterminacy and Soft Sets: Frontiers and Perspectives, *Mathematics*, 10, 2022, paper no. 3909.
- [10] Zadeh, L.A. Fuzzy Sets, *Information and Control*, 8, 1965, pp. 338-353.
- [11] Klir, G. J. & Folger, T. A., *Fuzzy Sets, Uncertainty and Information*, Prentice-Hall, London, 1988
- [12] Zadeh, L.A., Outline of a new approach to the analysis of complex systems and decision processes. *IEEE Trans. Syst. Man Cybern.*, 3, 1973, pp.28-44.
- [13] Kosko, B. *Fuzzy Thinking: The New Science of Fuzzy Logic*. Hyperion New York 1993
- [14] Zadeh, L.A. Fuzzy Sets as a basis for a theory of possibility, *Fuzzy Sets Syst.*, 1, 1978, pp. 3-28.
- [15] Dubois, D.; Prade, H. Possibility theory, probability theory and multiple-valued logics: A clarification, *Ann. Math. Artif. Intell.*, 32, 2001,
- [16] Kosko, B., Fuzziness Vs. Probability, *Int. J. of General Systems*, 17(2-3), 1990, pp. 211-240.
- [17] Voskoglou, M.Gr., Fuzzy Systems, Extensions and Relative Theories, *WSEAS Transactions on Advances in Engineering Education*, 16, 2019, pp. 63-69
- [18] Atanassov, K.T., Intuitionistic Fuzzy Sets, *Fuzzy Sets and Systems*, 20(1), 1986, pp.87-96.
- [19] Smarandache, F., *Neutrosophy / Neutrosophic probability, set, and logic*, Proquest, Michigan, USA, 1998.
- [20].Smarandache, F., Indeterminacy in Neutrosophic Theories and their Applications, *International Journal of Neutrosophic Science*, , 15(2), 2021, pp. 89-97.
- [21] Wang, H., Smarandache, F., Zhang, Y. and Sunderraman, R., Single Valued Neutrosophic Sets, *Review of the Air Force Academy (Brasov)*, 1(16), 2010, pp. 10-14.
- [22] Smarandache, F., Subtraction and Division of Neutrosophic Numbers, *Critical Review*, Vol. XIII, 2016, pp. 103-110.
- [23] Dubois, D., Prade, H., Interval-Valued Fuzzy Sets, Possibility Theory and Imprecise Probability, *Proceedings EUSFLAT-LFA*, 2005, pp. 314-319.
- [24] Deng, J., Control Problems of Grey Systems, *Systems and Control Letters*, 1982, 288-294.
- [25] Pawlak, Z., *Rough Sets: Aspects of Reasoning about Data*, Kluwer Academic Publishers, Dordrecht, 1991.
- [26] Molodtsov, D. , Soft Set Theory-First Results, *Computers and Mathematics with Applications*, 37(4-5), 1999, pp. 19-31.
- [27] Maji, P.K., Biswas, R., & Ray, A.R., Soft Set Theory, *Computers and Mathematics with Applications*, 45, 2003, pp. 555-562
- [28] Moore, R.A., Kearfort, R.B., Cloud, M.G., *Introduction to Interval Analysis*, 2nd ed., SIAM, Philadelphia, PA, USA, 1995.

Creative Commons Attribution License 4.0 (Attribution 4.0 International , CC BY 4.0)

This article is published under the terms of the Creative Commons Attribution License 4.0
https://creativecommons.org/licenses/by/4.0/deed.en_US