

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/364921918>

On The Classification of Neutrosophic Complex Inner Product Spaces

Article · January 2022

DOI: 10.54216/GJMSA.020104

CITATIONS

0

READS

4

2 authors:



Mehmet Çelik

9 PUBLICATIONS 31 CITATIONS

[SEE PROFILE](#)



Necati Olgun

Gaziantep University

69 PUBLICATIONS 580 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Symmetric Derivations, Exterior Derivation [View project](#)



Z-numbers [View project](#)



On The Classification of Neutrosophic Complex Inner Product Spaces

Mehmet Celik, Necati Olgun

Department Of Mathematics, Gaziantep University, Gaziantep, Turkey

Emails: mathcelik@gmail.com; olgun@gantep.edu.tr

Abstract

The objective of this paper is to study the neutrosophic complex inner product spaces over neutrosophic complex field $C(I)$. Also, it determines the necessary and sufficient condition of a neutrosophic complex vector space to be a complex inner product space by using semi module isomorphisms.

Key words: Neutrosophic vector space; neutrosophic complex number; neutrosophic inner product.

1.Introduction

Neutrosophy is a generalization of fuzzy ideas presented by Smarandache [13], where it was built over the idea of extending the degree of truth (T) and falsity (F) to a third logical case (the indeterminacy I).

Neutrosophic algebraic structures have been studied by many authors such as neutrosophic groups, neutrosophic rings, and matrices [2-7].

The study of neutrosophic real inner product spaces began in [15], where we find the structure of canonical neutrosophic inner products and orthogonality. This lead to many interesting results about isometries and norms in neutrosophic Euclidean geometry [11].

In this work, we combine the classical complex inner products , with neutrosophic vector spaces defined over neutrosophic complex numbers to get the complex neutrosophic inner product spaces. Also, we use the concept of semi-module isomorphisms [16], to classify the complex neutrosophic inner product spaces.

Main Concepts and Discussion

Definition :

Let V be a vector space over C , the corresponding neutrosophic complex vector space is defined as follows:

$$V(I) = \{a + bI; a, b \in V\}.$$

The scalars are taken for the neutrosophic complex field $C(I)$:

$$C(I) = \{(m + in) + (t + il)I; m, n, t, l \in C\}.$$

The following example clarifies operations on $V(I)$.

Example :

Let $V = C^2 = C \times C$ be the Euclidean complex vector space over C .

The corresponding neutrosophic complex vector space is:

$$C^2(I) = \{(x, y) + (z, t)I = (x + zI, y + tI); x, y, z, t \in C\}.$$

Doi : <https://doi.org/10.54216/GJMSA.020104>

Received: March 03, 2022 Accepted: July 18, 2022

Multiplication by a neutrosophic complex scalar can be showed as follows:

Take $X = (1 + i + (2 - i)I, i + I(1 + i))$, $\lambda = 1 - i + I(3i)$.

$$\lambda X = ([1 + i + (2 - i)I][1 - i + 3iI], [i + I(1 + i) \cdot [1 - i + 3iI]])$$

$$\lambda X = (2 + 3iI - 3I + (1 - 3i)I + (3 + 6i)I, 1 + i - 2I + 2I + (-3 + 3i)I)$$

$$\lambda X = (2 + (1 + 6i)I, 1 + i + (-4 + 3i)I).$$

$V(I)$ is a module over $C(I)$, that is because the neutrosophic complex field $C(I)$ is a ring in the ordinary meaning not a field.

Definition :

Let V be a vector space over C , $f: V \times V \rightarrow C$ is called a complex inner product if:

- 1) $f(a, a) = 0 \Rightarrow a = 0$, $f(a, a) \geq 0$, $f(a, a)$ is a real number.
- 2) $f(a, b) = \overline{f(b, a)}$
- 3) $f(a + b, c) = f(a, c) + f(b, c)$
- 4) $f(\lambda a, \mu b) = \lambda \bar{\mu} f(a, b)$; $a, b, c \in V$, $\lambda, \mu \in C$

Example :

Consider the complex vector space C over itself.

Define: $f: C \times C \rightarrow C$; $f(a, b) = a\bar{b}$.

We remark that: $f(a, a) = |a|^2 \geq 0$.

If $f(a, a) = 0 \Rightarrow a = 0$, $f(a, b) = a\bar{b} = \overline{(b \cdot \bar{a})} = \overline{f(b, a)}$.

$f(a + b, c) = (a + b)\bar{c} = a\bar{c} + b\bar{c} = f(a, c) + f(b, c)$.

$f(\lambda a, \mu b) = \lambda a(\bar{\mu} \bar{b}) = \lambda \bar{\mu} f(a, b)$ for all $a, b, c, \lambda, \mu \in C$

Definition :

Let $V(I)$ be a neutrosophic complex vector space over $C(I)$, consider the mapping $f: V(I) \times V(I) \rightarrow C(I)$ We say that f is a neutrosophic complex inner product if:

- 1) $f(a + bI, c + dI) = f(c + dI, a + bI)$.
- 2) $f(a + bI, a + bI) = 0 \Rightarrow a + bI = 0$, $f(a + bI, a + bI) \geq 0$ and $f(a + bI, a + bI) \in R(I)$.
- 3) $f([a + bI] + [c + dI], m + nI) = f(a + bI, m + nI) + f(c + dI, m + nI)$.
- 4) $f(\lambda(a + bI), \mu(c + dI)) = \lambda \bar{\mu} f(a + bI, c + dI)$; $a, b, c, d, m, n \in V$ and $\lambda, \mu \in C(I)$.

Theorem 10:

Let V be a vector space over C , with $g: V \times V \rightarrow C$ as a complex inner product.

Let $V(I)$ be the corresponding neutrosophic vector space over $C(I)$, then $V(I)$ has a neutrosophic complex inner product generated by g .

Proof.

Define $f: V(I) \times V(I) \rightarrow C(I)$; $f(a + bI, c + dI) = g(a, c) + I[g(a + b, c + d) - g(a, c)]$.

f is a neutrosophic complex inner product for the following reasons:

- 1) $f(a + bI, a + bI) = g(a, a) + I[g(a + b, a + b) - g(a, a)]$.

If $f(a + bI, a + bI) = 0$, then $g(a, a) = 0 \Rightarrow a = 0$, $g(a + b, a + b) = 0 \Rightarrow a + b = 0 \Rightarrow b = 0$

Thus $a + bI = 0$.

On other hand, $g(a, a), g(a + b, a + b)$ are two real positive numbers, hence:

$g(a, a) + I[g(a + b, a + b) - g(a, a)] \geq 0$ with respect to the neutrosophic partial order relation defined in [].

- 2) $f(a + bI, c + dI) = g(a, c) + I[g(a + b, c + d) - g(a, c)] = \overline{g(c, a)} + I[\overline{g(c + d, a + b)} - \overline{g(c, a)}] = \overline{f(c + dI, a + bI)}$.

- 3) Let $\lambda = p + qI, \mu = m + nI$; $p, q, m, n \in C$ be two neutrosophic complex scalars, hence

$$\begin{aligned} f(\lambda(a + bI), \mu(c + dI)) &= f(pa + [(p + q)(a + b) - pa]I, mc + [(m + n)(c + d) - mc]I) \\ &= g(pa, mc) + I[g((p + q)(a + b), (m + n)(c + d)) - g(pa, mc)] = p\bar{m}g(a, c) + I[(p + q)(\bar{m} + \bar{n})g(a + b, c + d) - p\bar{m}g(a, c)] \\ &= (p + qI)(\bar{m} + \bar{n}I)f(a + bI, c + dI). \end{aligned}$$

- 4) $f([a + bI] + [c + dI], m + nI) = f((a + c) + (b + d)I, m + nI)$
 $= g(a + c, m) + I[g(a + c + b + d, m + n) - g(a + c, m)]$
 $= g(a, m) + g(c, m) + I[g(a + b, m + n) + g(c + d, m + n) - g(a, m) - g(c, m)]$
 $= f(a + bI, m + nI) + f(c + dI, m + nI).$

Example:

Let $V = C^2$ be the Euclidean complex vector space over C .

V has an inner product defined as follows:

$g: C^2 \times C^2 \rightarrow C$ such that $g((x, y), (z, t)) = x\bar{z} + y\bar{t}$; $x, y, z, t \in C$

The corresponding neutrosophic complex inner product generated by g can be obtained as follows:

$f: C^2(I) \times C^2(I) \rightarrow C(I)$ such that:

Doi : <https://doi.org/10.54216/GJMSA.020104>

Received: March 03, 2022 Accepted: July 18, 2022

$$\begin{aligned} f((a + bI, c + dI), (x + yI, z + tI)) &= f((a, c) + (b, d)I, (x, z) + (y, t)I) \\ &= g((a, c), (x, z)) + I[g((a + b, c + d), (x + y, z + t)) - g((a, c), (x, z))] \\ &= a\bar{x} + c\bar{z} + I[(a + b).(\overline{x + y}) + (c + d).(\overline{z + t}) - a\bar{x} - c\bar{z}] \end{aligned}$$

Let $X = (1 + iI, 2 - i + I), Y = (1 + i + (1 - i)I, (3i)I) \in C(I)$, hence

$$\begin{aligned} f(X, Y) &= f((1, 2 - i) + I(i, 1), (1 + i, 0) + I(1 - i, 3i)) = \\ &= g((1, 2 - i), (1 + i, 0)) + I[g((1 + i, 3 - i), (2, 3i)) - g((1, 2 - i), (1 + i, 0))] = \\ &= 1(1 + i) + (2 - i)\bar{0} + I[(1 + i).2 + (3 - i).(3i) - 1(1 + i) - (2 - i)\bar{0}] = \\ &= 1 - i + I[2 + 2i - 9i - 3 - 1 + i] = 1 - i + I[2 + 2i - 9i - 3 - 1 + i] = 1 - i + I[-2 - 6i]. \end{aligned}$$

Definition:

Let $V(I)$ be a neutrosophic complex vector space over $C(I)$ with a neutrosophic complex inner product f . We say:

- 1) $a + bI \perp c + dI$ if and only if $f(a + bI, a + bI) = 0$.
- 2) The norm of $a + bI$ is defined as follows:

$$\|a + bI\| = \sqrt{f(a + bI, a + bI)}.$$

Theorem :

Let $V(I)$ be a neutrosophic complex inner product space over $C(I)$, with a neutrosophic complex inner product f generated by a classical complex inner product $g: V \times V \rightarrow C$, we have:

- 1) $a + bI \perp c + dI$ if and only if $a \perp c$ and $a + b \perp c + d$.
- 2) $\|a + bI\| = \|a\| + I(\|a + b\| - \|a\|)$.
- 3) $\|(a + bI) + (c + dI)\| \leq \|a + bI\| + \|c + dI\|$.
- 4) $\|\lambda(a + bI)\| = |\lambda|\|a + bI\|$; $a, b, c, d \in V$ and $\lambda \in C(I)$.

Proof.

- 1) $f(a + bI, a + bI) = g(a, c) + I[g(a + b, c + d) - g(a, c)]$
 $f(a + bI, a + bI) = 0$ if and only if $g(a, c) = g(a + b, c + d) = 0$, thus $a \perp c$ and $a + b \perp c + d$.
- 2) $\|a + bI\| = \sqrt{f(a + bI, a + bI)} = \sqrt{g(a, a) + I[g(a + b, a + b) - g(a, a)]}$
 $= \sqrt{g(a, a)} + I[\sqrt{g(a + b, a + b)} - \sqrt{g(a, a)}] = \|a\| + I(\|a + b\| - \|a\|)$.
- 3) $\|(a + bI) + (c + dI)\| = \|(a + c) + (b + d)I\| = \|a + c\| + I(\|a + c + b + d\| - \|a + c\|)$
 $\leq \|a\| + \|c\| + I(\|a + b\| + \|c + d\| - \|a\| - \|c\|) \leq \|a\| + I(\|a + b\| - \|a\|) + \|c\| + I(\|c + d\| - \|c\|) \leq$
 $\|a + bI\| + \|c + dI\|$.
- 4) $\|\lambda(a + bI)\|^2 = f(\lambda(a + bI), \lambda(a + bI)) = \lambda\bar{\lambda}(a + bI)(\overline{a + bI}) = (|\lambda|\|a + bI\|)^2$. By taking the square root of the two sides, we get the desired result.

Theorem :

Let V be a vector space over C , with $g: V \times V \rightarrow C$ as a complex inner product, let $V(I)$ be the corresponding neutrosophic complex inner product space over $C(I)$, with f as a neutrosophic complex inner product generated by g , hence $V(I)$ is a semi isomorphic to V .

Proof.

Define $T: V(I) \rightarrow V \times V$ such that $T(a + bI) = (a, a + b)$; $a, b \in V$.

Consider the AH-isometry $\emptyset: C(I) \rightarrow C \times C$; $\emptyset(x + yI) = (x, x + y)$; $x, y \in C$.

\emptyset is a ring isomorphism.

Now, we must prove that T is a semi isomorphism.

$\forall a + bI, c + dI \in V(I)$, it is clear that:

a neutrosophic complex inner product for the following reasons:

$$T[(a + bI) + (c + dI)] = T(a + bI) + T(c + dI).$$

Also, it is easy to check that T is a bijection.

Now, we must prove that $T[(x + yI). (a + bI)] = \emptyset(x + yI). T(a + bI)$ for all $x, y \in C, a, b \in V$.

$$T[(x + yI). (a + bI)] = T[xa + I(xb + ya + yb)] = (xa, xa + xb + ya + yb) =$$

$$(x, x + y). (a, a + b) = \emptyset(x + yI). T(a + bI), \text{ thus } T \text{ is a semi isomorphism.}$$

According to the previous theorem, we get the following result.

If $V(I)$ is a complex neutrosophic inner product space with f as a neutrosophic complex inner product generated by a classical complex inner product $g: V \times V \rightarrow C$, hence $V(I) \cong_s V \times V$.

The converse of this result is hard to prove.

If $V(I)$ is a complex neutrosophic inner product space with $f: V(I) \times V(I) \rightarrow C(I)$, then can we find a classical complex inner product $g: V \times V \rightarrow C$ such that f generated by g and $V(I) \cong_s V \times V$?

To solve the inverse problem, it is sufficient to find a classical complex inner product $g: V \times V \rightarrow C$ such that f is generated by g .

Theorem :

Let $V(I)$ be a complex neutrosophic inner product space over $\mathcal{C}(I)$ with $f: V(I) \times V(I) \rightarrow \mathcal{C}(I)$ as complex neutrosophic inner product space, then V is a complex inner product space over \mathcal{C} .

Proof

Define $g: V \times V \rightarrow \mathcal{C}$; $g(a, c) = f(a + 0I, c + 0I)$ we have:

$$a) \quad g(a, a) = 0 \Rightarrow f(a, a) = 0 \Rightarrow a = 0.$$

$$g(a, a) = f(a, a) \geq 0 \in \mathcal{R}.$$

$$b) \quad g(a + b, c) = f(a + b + 0I, c + 0I) = f(a + 0I, c + 0I) + f(b + 0I, c + 0I) \\ = g(a, c) + g(b, c).$$

$$c) \quad g(\lambda a, \mu c) = f(\lambda a + 0I, \mu c + 0I) = \lambda \bar{\mu} f(a + 0I, c + 0I) = \lambda \bar{\mu} g(a, c)$$

Thus g is a complex inner product on the classical vector space V .

Remark.

From the previous theorems, we get the following result.

According complex inner product space $V(I)$ is semi isomorphic to the direct product of the corresponding classical inner product space V .

References

- [1] Abobala, M., Ziena, B.M., Doewes, R., and Hussein, Z., "The Representation Of Refined Neutrosophic Matrices By Refined Neutrosophic Linear Functions", International Journal Of Neutrosophic Science, 2022.
- [2] Olgun, N., Hatip, A., Bal, M., and Abobala, M., " A Novel Approach To Necessary and Sufficient Conditions For The Diagonalization of Refined Neutrosophic Matrices", International Journal of Neutrosophic Science, Vol. 16, pp. 72-79, 2021.
- [3] E. Adeleke. A. Agboola. , and F. Smarandache. Refined Neutrosophic Rings II. International Journal of Neutrosophic Science, Vol. 2, 2020. pp. 89-94.
- [4] E. Adeleke. A. Agboola. , and F. Smarandache. Refined Neutrosophic Rings I. International Journal of Neutrosophic Science, Vol. 2, 2020, pp. 77-81.
- [5] V. Kandasamy and F. Smarandache. Some Neutrosophic Algebraic Structures and Neutrosophic N-Algebraic Structures", Hexis, Phonex, Arizona, pp.125-240, 2006.
- [6] A. Hatip and N.Olgun. On Refined Neutrosophic R-Module. International Journal of Neutrosophic Science, Vol. 7, 2020, pp.87-96.
- [7] M. Ibrahim. A. Agboola, B.Badmus and S. Akinleye. On refined Neutrosophic Vector Spaces . International Journal of Neutrosophic Science, Vol. 7, 2020, pp. 97-109.
- [8] Abobala, M., Hatip, A., Bal,M., " A Study Of Some Neutrosophic Clean Rings", International journal of neutrosophic science, 2022.
- [9] Abobala, M., "On The Characterization of Maximal and Minimal Ideals In Several Neutrosophic Rings", Neutrosophic Sets and Systems, Vol. 45, 2021.
- [10] Abobala, M., and Hatip, A., "An Algebraic Approach To Neutrosophic Euclidean Geometry", Neutrosophic Sets and Systems, Vol. 43, 2021.
- [11] Smarandache, F., "Neutrosophic Set a Generalization of the Intuitionistic Fuzzy Sets", Inter. J. Pure Appl. Math., pp. 287-297. 2005.
- [12] Smarandache, F., " A Unifying Field in Logics: Neutrosophic Logic, Neutrosophy, Neutrosophic Set, Neutrosophic Probability", American Research Press. Rehoboth, 2003.
- [13] Abobala, M., Bal, M., Aswad, M., "A Short Note On Some Novel Applications of Semi Module Homomorphisms", International journal of neutrosophic science, 2022.
- [14] Abobala, M., "Neutrosophic Real Inner Product Spaces", Neutrosophic Sets and Systems, Vol. 43, 2021.
- [15] Abobala, M., Bal, M., Aswad, M., "A Short Note On Some Novel Applications of Semi Module Homomorphisms", International journal of neutrosophic science, 2022.