

RESEARCH

Open Access



Nonlinear operators between neutrosophic normed spaces and Fréchet differentiation

Vakeel A. Khan^{1*} and Mohammad Daud Khan¹

*Correspondence:

vakhanmaths@gmail.com

¹Department of Mathematics,
Aligarh Muslim University, Aligarh,
202002, India

Abstract

The article focuses on the introduction of neutrosophic continuity and neutrosophic boundedness, which is a fair extension of intuitionistic fuzzy continuity and intuitionistic fuzzy boundedness, respectively. The article further advances to illustrate the Fréchet derivative of nonlinear operators between neutrosophic normed spaces (NNS). Examples have been provided in compliance with the theory with the aid of some standard sequence spaces.

Keywords: Bounded Linear operator; Fréchet derivative; Continuity; Neutrosophic normed space

1 Introduction

The researchers persistently come across distinct mathematical problems that quite often fail to get solved or analyzed by the classical theory of nonlinear analysis. A handful of the arduous problems can be reduced to operator equations to analyze them, and the Fréchet derivative plays a crucial role in solving such problems. The extension of crisp sets was put forth by Zadeh [1], wherein each element had a degree of membership indicating the extent to which an element belongs in a set. This notion is a strong mathematical tool to deal with the complexity of uncertainty in the form of vagueness in several problems arising in the area of science and engineering. It has valuable applications in areas such as population dynamics [2], computer programming [3], chaos control [4, 5]. As an extension of fuzzy sets, Atanassov [6] proposed the concept of intuitionistic fuzzy sets in 1986, which incorporated the degree of non-membership and hesitant function along with the degree of membership. A few papers depicting convergence in the setting of intuitionistic fuzzy sets were defined by Mursaleen et al. [7, 8], where he analyzed the statistical and ideal convergence in intuitionistic fuzzy topological space. Recently, Khan et al. [9] studied the continuous and bounded linear operators in neutrosophic normed spaces.

In the ultimate presence of quantum gravitational effects, the theory of the Heisenberg uncertainty principle must be generalized on the basis of fuzzy structure spacetime. These facts were mainly encouraged by string theory and non-commutative geometry. Quantum gravity robustly explained the spacetime points in a fuzzy way. So, the contrariety of determining the exact location of articles gives spacetime a vague structure [4, 10, 11]. Due to this ambiguous structure, the position space depiction of quantum mechanics may break

© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

down. Therefore, in this manner, a generalized normed space of quasi-position eigenfunction is specifically required [12]. It is monitored that in the quantum gravity regime, the most basic idea of spacetime in self-identity causes diffusion on the wave packet profile. The formed fact originates in the quantum fluctuation of spacetime, best described as fuzzy spacetime.

Smarandache et al. [13] projected another idea, which can be said to be a neutrosophic set by counting a transitional membership function, which was expected to be a formal setting trying to compute the truth, indeterminacy, and falsehood. Later on in [14], he furthermore studied the contrariness between intuitionistic fuzzy set and neutrosophic set, as well as the relations between these two sets [10]. In the generated neutrosophic set, the components T (truth), S (indeterminacy), and U (falsity) have values in between $]0^-, 1^+]$.

Bera and Mahapatra [15, 16] imported the neutrosophic soft normed linear space and defined convexity, metric, and Cauchy sequence on it. Recently, Kirişci & Şimşek [12] examined the statistical convergence on NNS , while Khan et al. [17] evaluated some results of NNS via the Fibonacci matrix. By reviewing the literature, we can conclude that there has been exponential growth in the study of neutrosophic theory in dissimilar fields by various researchers. There are a number of conditions where the norm of a vector is impossible to get, and the concept of neutrosophic norm [18–20] looks more authentic in such cases. These theories can best deal with situations by modeling the inexactness through the neutrosophic norm.

This presented paper resolves and studies the Fréchet derivative of nonlinear operators between NNS , which may produce a helpful functional tool to explain the operator equations. Likewise, particular and universal discordance analysis of solutions of operator mathematical equations are determinate over this concept. So, here we have tried a proper approach of nonlinear functional analysis of operator equations by applying Fréchet derivative.

2 Preliminaries

Definition 2.1 ([6]) Let $F \neq \phi$ be the intuitionistic fuzzy set, and $F \subseteq Y$ is an ordered triplet defined by

$$F = \{ \langle \eta, T(\eta), U(\eta) \rangle : \eta \in Y \},$$

where $T(\eta), U(\eta) : Y \rightarrow [0, 1]$ represent the degree of membership and degree of non-membership, respectively, in such a way that

$$0 \leq T(\eta) + U(\eta) \leq 1$$

$1 - T(\eta) - U(\eta)$ is also said to be degree of hesitancy. The intuitionistic fuzzy components $T(\eta)$, $U(\eta)$ and degree of hesitancy depend on each other.

Definition 2.2 ([14]) Let $A \neq \phi$, $A \subseteq Y$. Then,

$$A_{NS} = \{ \langle \eta, T(\eta), S(\eta), U(\eta) \rangle : \eta \in Y \},$$

where $\mathbf{T}(\eta), \mathbf{S}(\eta), \mathbf{U}(\eta) : Y \rightarrow [0, 1]$ represent the degree of membership, degree of indeterminacy, and degree of nonmembership, respectively, in such a way that

$$0 \leq \mathbf{T}(\eta) + \mathbf{S}(\eta) + \mathbf{U}(\eta) \leq 3.$$

The neutrosophic components $\mathbf{T}(\eta)$, $\mathbf{S}(\eta)$ and $\mathbf{U}(\eta)$ are independent of each other.

Definition 2.3 ([21]) A continuous t -norm is a binary operation $\star : [0, 1] \times [0, 1] \rightarrow [0, 1]$ with the following conditions: (i) \star is associative and commutative; (ii) \star is continuous; (iii) $\rho_1 \star 1 = \rho_1$, $\forall \rho_1 \in [0, 1]$; (iv) $\rho_1 \star \rho_2 \leq \rho_3 \star \rho_4$ whenever $\rho_1 \leq \rho_3$ and $\rho_2 \leq \rho_4$, for each $\rho_1, \rho_2, \rho_3, \rho_4 \in [0, 1]$.

Definition 2.4 ([21]) A continuous t -conorm is a binary operation $\circ : [0, 1] \times [0, 1] \rightarrow [0, 1]$ with the following conditions (i) \circ is associative and commutative; (ii) \circ is continuous; (iii) $\rho_1 \circ 0 = \rho_1$, $\forall \rho_1 \in [0, 1]$; (iv) $\rho_1 \circ \rho_2 \leq \rho_3 \circ \rho_4$ whenever $\rho_1 \leq \rho_3$ and $\rho_2 \leq \rho_4$, for each $\rho_1, \rho_2, \rho_3, \rho_4 \in [0, 1]$.

Definition 2.5 ([22]) Consider Y be a linear space and $\mathcal{M} = \{\langle \eta, \mathbf{T}(\eta), \mathbf{S}(\eta), \mathbf{U}(\eta) \rangle : \eta \in Y\}$ be a normed space in such a way that $\mathcal{M} : Y \times \mathbb{R}^+ \rightarrow [0, 1]$. Let \bullet and \star be continuous t -norm and continuous t -conorm, respectively. Then, the four-tuple $(Y, \mathcal{M}, \bullet, \star)$ is called neutrosophic normed space (NNS) if it satisfies the following axioms, $\forall \eta, y, z \in Y$ and $d, t > 0$;

- (i) $0 \leq \mathbf{T}(\eta, t), \mathbf{S}(\eta, t), \mathbf{U}(\eta, t) \leq 1$,
- (ii) $0 \leq \mathbf{T}(\eta, t) + \mathbf{S}(\eta, t) + \mathbf{U}(\eta, t) \leq 3$,
- (iii) $\mathbf{T}(\eta, t) = 0$ for $t \leq 0$,
- (iv) $\mathbf{T}(\eta, t) = 1$ for $t > 0$ iff $\eta = 0$
- (v) $\mathbf{T}(c\eta, t) = \mathbf{T}(\eta, \frac{t}{|c|}) \forall c \neq 0, t > 0$,
- (vi) $\mathbf{T}(\eta, t) \bullet \mathbf{T}(y, d) \leq \mathbf{T}(\eta + y, t + d)$
- (vii) $\mathbf{T}(\eta, \bullet)$ is continuous nondecreasing function for $t > 0$, $\lim_{t \rightarrow \infty} \mathbf{T}(\eta, t) = 1$
- (viii) $\mathbf{S}(\eta, t) = 1$ for $t \leq 0$,
- (ix) $\mathbf{S}(\eta, t) = 0$ for $t > 0$ iff $\eta = 0$
- (x) $\mathbf{S}(c\eta, t) = \mathbf{S}(\eta, \frac{t}{|c|}) \forall c \neq 0$
- (xi) $\mathbf{S}(\eta, t) \star \mathbf{S}(y, d) \geq \mathbf{S}(\eta + y, t + d)$,
- (xii) $\mathbf{S}(\eta, \star)$ is continuous nonincreasing function, $\lim_{t \rightarrow \infty} \mathbf{S}(\eta, \star) = 0$,
- (xiii) $\mathbf{U}(\eta, t) = 1$ for $t \leq 0$,
- (xiv) $\mathbf{U}(\eta, t) = 0$ for $t > 0$ iff $\eta = 0$,
- (xv) $\mathbf{U}(c\eta, t) = \mathbf{U}(\eta, \frac{t}{|c|}) \forall c \neq 0$
- (xvi) $\mathbf{U}(\eta, t) \star \mathbf{U}(y, d) \geq \mathbf{U}(\eta + y, t + d)$,
- (xvii) $\mathbf{U}(\eta, \star)$ is continuous nonincreasing function, $\lim_{t \rightarrow \infty} \mathbf{U}(\eta, t) = 0$.

In this case, \mathcal{M} is called neutrosophic norm (NN).

Since all the components lie between $[0, 1]$, therefore, if the indeterminacy is zero, then still neutrosophic components are more flexible and more general than fuzzy and intuitionistic fuzzy components. If all the neutrosophic components on a vector space satisfy all conditions of Definition 2.5, then we say that $(Y, \mathcal{M}, \bullet, \star)$ is NNS, where $\mathcal{M} = \{\langle \eta, \mathbf{T}(\eta), \mathbf{S}(\eta), \mathbf{U}(\eta) \rangle : \eta \in Y\}$.

Neutrosophic normed space is more general than norm space. This is illustrated by proper example

Let $(Y, \mathcal{M}, \bullet, \star)$ be NNS. Assume that $\eta \bullet y = \eta y$ and $\eta \star y = \eta + y - \eta y$ for all $\eta, y \in Y$ and $t > 0$ with the condition

$$\mathbf{T}(\eta, t) > 0 \quad \text{and} \quad \mathbf{S}(\eta, t) < 1, \quad \mathbf{U}(\eta, t) < 1 \quad \implies \quad \eta = 0 \quad \text{for all } t > 0.$$

Let $\|\eta\|_\beta = \inf\{t > 0 : \mathbf{T}(\eta, t) \geq \beta \text{ and } \mathbf{S}(\eta, t) \leq 1 - \beta, \mathbf{U}(\eta, t) \leq 1 - \beta\}$, $\forall \beta \in (0, 1)$. Then, $\{\|\cdot\|_\beta : \beta \in (0, 1)\}$ is an ascending family of norms on Y . These norms are said to be β -norms on Y compatible to neutrosophic norm $(\mathbf{T}, \mathbf{S}, \mathbf{U})$.

3 Neutrosophic continuity

Definition 3.1 ([23]) Let $(Y, \mathbf{T}_1, \mathbf{S}_1, \mathbf{U}_1, \bullet, \star)$ and $(W, \mathbf{T}', \mathbf{S}', \mathbf{U}', \bullet, \star)$ be two NNS, and consider a mapping h from $(Y, \mathbf{T}_1, \mathbf{S}_1, \mathbf{U}_1, \bullet, \star) \rightarrow (W, \mathbf{T}', \mathbf{S}', \mathbf{U}', \bullet, \star)$. Then,

- (i) ξ is called strong neutrosophic limit of h at some $\eta_0 \in Y$ if for each $\epsilon > 0$, \exists some $\delta = \delta(\epsilon) > 0$ such that

$$\mathbf{T}'(h(\eta) - \xi, \epsilon) \geq \mathbf{T}_1(\eta - \eta_0, \delta),$$

$$\mathbf{S}'(h(\eta) - \xi, \epsilon) \leq \mathbf{S}_1(\eta - \eta_0, \delta),$$

$$\mathbf{U}'(h(\eta) - \xi, \epsilon) \leq \mathbf{U}_1(\eta - \eta_0, \delta).$$

In this case, we write $(\text{Strong Neutrosophic-SN})\text{-}\lim_{\eta \rightarrow \eta_0} h(\eta) = \xi$, which also means that

$$\lim_{\mathbf{T}_1(\eta - \eta_0, t) \rightarrow 1} \mathbf{T}'(h(\eta) - \xi, t) = 1, (SN) \quad \text{and}$$

$$\lim_{\mathbf{S}_1(\eta - \eta_0, t) \rightarrow 0} \mathbf{S}'(h(\eta) - \xi, t) = 0 (SN),$$

$$\lim_{\mathbf{U}_1(\eta - \eta_0, t) \rightarrow 0} \mathbf{U}'(h(\eta) - \xi, t) = 0 (SN)$$

or

$$\begin{cases} \mathbf{T}_1(h(\eta) - \xi, t) = \xi, (SN) \text{ as } \mathbf{T}'(\eta - \eta_0, t) \rightarrow 1, \text{ and } \mathbf{S}_1(h(\eta) - \xi, t) = \xi, (SN) \text{ as} \\ \mathbf{S}'(\eta - \eta_0, t) \rightarrow 0 \text{ } \mathbf{U}_1(h(\eta) - \xi, t) = \xi, (SN) \text{ as } \mathbf{U}'(\eta - \eta_0, t) \rightarrow 0 \end{cases}$$

$\forall t > 0$.

- (ii) ξ is called weak neutrosophic limit of h at some $\eta_0 \in Y$ if for given $\epsilon > 0$ & $\beta \in (0, 1)$, $\delta = \delta(\epsilon, \beta) > 0$ s.t

$$\mathbf{T}_1(\eta - \eta_0, \delta) \geq \beta \quad \implies \quad \mathbf{T}'(h(\eta) - \xi, \epsilon) \quad \text{and}$$

$$\mathbf{S}'(\eta - \eta_0, \delta) \leq 1 - \beta \quad \implies \quad \mathbf{S}_1(h(\eta) - \xi, \epsilon),$$

$$\mathbf{U}'(\eta - \eta_0, \delta) \leq 1 - \beta \quad \implies \quad \mathbf{U}_1(h(\eta) - \xi, \epsilon).$$

In this case, we write $(\text{Weak Neutrosophic-WN})\text{-}\lim_{\eta \rightarrow \eta_0} h(\eta) = \xi$, which also means that

$$\lim_{T_1(\eta - \eta_0, t) \rightarrow 1} T'(h(\eta) - \xi, t) = \xi, (WN) \quad \text{and}$$

$$\lim_{S_1(\eta - \eta_0, t) \rightarrow 0} S'(h(\eta) - \xi, t) = \xi (SN),$$

$$\lim_{U_1(\eta - \eta_0, t) \rightarrow 0} U'(h(\eta) - \xi, t) = \xi (SN).$$

or

$$\begin{cases} T'(h(\eta) - \xi, t) = \xi, (WN) \text{ as } T_1(\eta - \eta_0, t) \rightarrow 1, \text{ and } S'(h(\eta) - \xi, t) = \xi, (WN) \text{ as} \\ S_1(\eta - \eta_0, t) \rightarrow 0 \text{ } U'(h(\eta) - \xi, t) = \xi, (WN) \text{ as } U_1(\eta - \eta_0, t) \rightarrow 0 \end{cases}$$

for all $t > 0$.

Definition 3.2 Let $h_n : (Y, T_1, S_1, U_1, \bullet, \star) \rightarrow (W, T', S', U', \bullet, \star)$ be a sequence of functions. Then Ψ_n is called pointwise neutrosophic convergent on Y to a function Ψ w.r.t (T, S, U) if for each $\eta \in Y$, the sequence $(\Psi_n(\eta))$ is convergent to $\Psi(\eta)$ w.r.t (T, S, U) .

Definition 3.3 ([12]) Suppose $(Y, \mathcal{M}, \bullet, \star)$ is an NNS. A sequence $m = (m_i)$ is said to be a Cauchy sequence w.r.t \mathcal{M} , if for each $\epsilon > 0$ & $t > 0$, $\exists d \in \mathbb{N}$ s.t $T(m_i - m_k, t) > 1 - \epsilon$, $S(m_i - m_k, t) < \epsilon$ and $U(m_i - m_k, t) < \epsilon$ for all $i, k \geq d$.

Definition 3.4 Suppose $(Y, T_1, S_1, U_1, \bullet, \star)$ and $(W, T', S', U', \bullet, \star)$ are two NNS. A mapping h from $(Y, T_1, S_1, U_1, \bullet, \star) \rightarrow (W, T', S', U', \bullet, \star)$ is called neutrosophic continuous at $\rho_0 \in Y$ if for any given $\epsilon > 0$, $\exists t = t(b, \epsilon)$, $d = d(b, \epsilon) \in (0, 1)$ such that $\forall \rho \in Y$ and for all $b \in (0, 1)$,

$$T_1(\rho - \rho_0, t) > 1 - \gamma \implies T'(\Psi(\rho) - \Psi(\rho_0), \epsilon) > 1 - b,$$

$$S_1(\rho - \rho_0, t) < \gamma \implies S'(\Psi(\rho) - \Psi(\rho_0), \epsilon) < b,$$

$$U_1(\rho - \rho_0, t) < \gamma \implies U'(\Psi(\rho) - \Psi(\rho_0), \epsilon) < b.$$

Proposition 3.1 $(SN) - \lim \implies (WN) - \lim$ but the contrary need not hold. Further, $(WN) - \lim = (SN) - \lim$ whenever $(SN) - \lim$ exists.

Proof It is simple to show. Now, we prove that $(WN) - \lim$ does not imply $(SN) - \lim$ in regular. \square

Example 3.1 Consider $Y = W = \mathbb{R}$

$$T_1(\eta, t) = \begin{cases} \frac{t}{t + \|\eta\|} & \text{if } t > \|\eta\|, \\ 0 & \text{otherwise;} \end{cases}$$

$$T'(\eta, t) = \begin{cases} 1, & \text{if } t > \|\eta\|, \\ 0, & \text{if } t \leq \|\eta\|; \end{cases}$$

and

$$S_1(\eta, t) = \begin{cases} \frac{\|\eta\|}{t + \|\eta\|} & \text{if } t > \|\eta\|, \\ 1 & \text{otherwise;} \end{cases}$$

$$S'(\eta, t) = \begin{cases} 1, & \text{if } t \leq \|\eta\|, \\ 0, & \text{if } t > \|\eta\|; \end{cases}$$

$$U_1(\eta, t) = \begin{cases} \frac{\|\eta\|}{t} & \text{if } t > \|\eta\|, \\ 1 & \text{otherwise;} \end{cases}$$

$$U'(\eta, t) = \begin{cases} 1, & \text{if } t \leq \|\eta\|, \\ 0, & \text{if } t > \|\eta\|. \end{cases}$$

Let the function h from $(\mathbb{R}, T_1, S_1, U_1, \bullet, \star)$ onto $(\mathbb{R}, T', S', U', \bullet, \star)$ be defined by $h(\eta) = \eta$. Then $(WN) - \lim_{\eta \rightarrow 0} h(\eta) = 0$. However, $(SN) - \lim_{\eta \rightarrow 0} h(\eta)$ does not exist. Because, for $\|\eta\| = \epsilon$ there is no $\delta > 0$ satisfying the conditions

$$T'(\eta, \epsilon) = 0 \geq T_1(\eta, \delta) = \frac{\|\delta\|}{\delta + \|\eta\|} = \frac{\|\delta\|}{\delta + \|\epsilon\|}$$

and

$$S'(\eta, \epsilon) = 0 \leq S_1(\eta, \delta) = \frac{\|\epsilon\|}{\delta + \|\epsilon\|},$$

$$U'(\eta, \epsilon) = 0 \leq U_1(\eta, \delta) = \frac{\|\epsilon\|}{\delta + \|\epsilon\|}.$$

Now, we assign the strong and weak neutrosophic continuity of mappings between NNS.

Definition 3.5 ([24]) Suppose $(Y, \mathcal{M}, \bullet, \star)$ is an NNS. & $V \subset Y$. Then V is called neutrosophic open subset Y if for every $\eta \in V$, there exists some $t > 0$ and $\beta \in (0, 1)$ such that $B(\eta, \beta, t) \subseteq V$, where

$$B(\eta, \beta, t) := \{y : T(\eta - y, t) > 1 - \beta \text{ and } S(\eta - y, t) < \beta, U(\eta - y, t) < \beta\}.$$

Definition 3.6 Suppose $(Y, T_1, S_1, U_1, \bullet, \star)$ and $(W, T', S', U', \bullet, \star)$ are two NNS. A mapping $h : Y \rightarrow W$. Then h is called

- (i) weakly neutrosophic continuous at $\eta_0 \in Y$ if for given $\epsilon > 0$ and $\beta \in (0, 1) \exists$ some $\delta = \delta(\epsilon, \beta) > 0$ s.t

$$T_1(\eta - \eta_0, \delta) \geq \beta \implies T'(h(\eta) - h(\eta_0), \epsilon) \text{ and}$$

$$S_1(\eta - \eta_0, \delta) \leq 1 - \beta \implies S'(h(\eta) - h(\eta_0), \epsilon),$$

$$U_1(\eta - \eta_0, \delta) \leq 1 - \beta \implies U'(h(\eta) - h(\eta_0), \epsilon),$$

for all $\eta \in Y$.

- (ii) strongly neutrosophic continuous at $\eta_0 \in Y$ if for given $\epsilon > 0$ and $\beta \in (0, 1) \exists$ some $\delta = \delta(\epsilon, \beta) > 0$ such that

$$\begin{aligned} \mathbf{T}'(h(\eta) - h(\eta_0), \epsilon) &\geq \mathbf{T}_1(\eta - \eta_0, \delta) \quad \text{and} \quad \mathbf{S}'(h(\eta) - h(\eta_0), \epsilon) \leq \mathbf{S}_1(\eta - \eta_0, \delta), \\ \mathbf{U}'(h(\eta) - h(\eta_0), \epsilon) &\leq \mathbf{U}_1(\eta - \eta_0, \delta), \end{aligned}$$

for all $\eta \in Y$.

- (iii) Let h be linear. Then h is said be weakly neutrosophic bounded (for short, WNB) on Y if for given $\beta \in (0, 1) \exists$ some, $n_\beta > 0$ such that

$$\begin{aligned} \mathbf{T}_1\left(\eta, \frac{t}{n_\beta}\right) &\geq \beta \quad \implies \quad \mathbf{T}'(h(\eta), t) \geq \beta \quad \text{and} \\ \mathbf{S}_1\left(\eta, \frac{t}{n_\beta}\right) &\leq 1 - \beta \quad \implies \quad \mathbf{S}'(h(\eta), t) \leq 1 - \beta, \\ \mathbf{U}_1\left(\eta, \frac{t}{n_\beta}\right) &\leq 1 - \beta \quad \implies \quad \mathbf{U}'(h(\eta), t) \leq 1 - \beta, \end{aligned}$$

for all $\eta \in Y$ and $t > 0$. Let $E'(Y, W)$ indicate the set of all WNB linear operators.

- (iv) Let h be linear. Then h is called weakly neutrosophic bounded (for short, SNB) on Y if for given $\beta \in (0, 1)$, \exists some, $K > 0$ such that

$$\begin{aligned} \mathbf{T}'(h(\eta), t) &\geq \mathbf{T}_1\left(\eta, \frac{t}{K}\right) \quad \text{and} \quad \mathbf{S}'(h(\eta), t) \leq \mathbf{S}_1\left(\eta, \frac{t}{K}\right), \\ \mathbf{U}'(h(\eta), t) &\leq \mathbf{U}_1\left(\eta, \frac{t}{K}\right), \end{aligned}$$

for all $\eta \in Y$ and $t > 0$. Let $E(Y, W)$ denote the set of all SNB linear operators.

Theorem 3.1 Suppose $(Y, \mathbf{T}_1, \mathbf{S}_1, \mathbf{U}_1, \bullet, \star)$ and $(W, \mathbf{T}', \mathbf{S}', \mathbf{U}', \bullet, \star)$ are two NNS and $h : Y \rightarrow W$ be a linear mapping. Then h is strongly (weakly) neutrosophic continuous if it is strongly (weakly) neutrosophic bounded.

Definition 3.7 Consider $(Y, \mathbf{T}_1, \mathbf{S}_1, \mathbf{U}_1, \bullet, \star)$ be an NNS. Then, a sequence η_p is called

- (i) weakly neutrosophic convergent to $\eta \in Y$ if and only if, for every $\epsilon > 0$ & $\beta \in (0, 1)$, \exists some, $p_0 = p_0(\beta, \epsilon)$ such that

$$\begin{aligned} \mathbf{T}_1(\eta_p - \eta_0, \epsilon) &\geq \beta \quad \& \quad \mathbf{S}_1(\eta_p - \eta_0, \epsilon) \leq 1 - \beta, \\ \mathbf{U}_1(\eta - \eta_0, \epsilon) &\leq 1 - \beta \quad \text{for all } \eta \geq \eta_0. \end{aligned}$$

In this case, we write $\eta_p \xrightarrow{WN} \eta$.

- (ii) strongly neutrosophic convergent to $\eta \in Y$ if and only if, for every $\beta \in (0, 1)$, there exists some $p_0 = p_0(\epsilon)$ such that

$$\mathbf{T}_1(\eta_p - \eta, t) \geq 1 - \beta \quad \text{and} \quad \mathbf{S}_1(\eta_p - \eta_0, t) \leq \beta, \quad \mathbf{U}_1(\eta - \eta_0, t) \leq \beta, \quad \forall t > 0,$$

we write $\eta_p \xrightarrow{SN} \eta$.

Accordingly, we can define an SN(WN)-Cauchy sequence, SN(WN)-closure of a subset and an SN(WN)-complete NNS.

Theorem 3.2 *If a sequence (η_p) is SN-convergent then it is WN-convergent to the same limit, but not conversely.*

Example 3.2 It is easy to show that SN-convergence implies WN-convergence. Following is an example to depict that the converse of the statement may not be true. Consider c_0 , the Banach space of all sequences $\eta = \eta_p$ convergent to zero with the sup-norm $\|\eta\|_\infty = \sup_p \|\eta\|$, and recognize the neutrosophic norm

$$\mathbf{T}_1(\eta, t) = \begin{cases} \frac{t - \|\eta\|_\infty}{t + \|\eta\|_\infty} & \text{if } t > \|\eta\|_\infty, \\ 0 & \text{if } t \leq \|\eta\|_\infty; \end{cases}$$

and

$$\mathbf{S}_1(\eta, t) = \begin{cases} \frac{2\|\eta\|_\infty}{t + \|\eta\|_\infty} & \text{if } t < \|\eta\|_\infty, \\ 1 & \text{if } t \leq \|\eta\|_\infty; \end{cases}$$

$$\mathbf{U}_1(\eta, t) = \begin{cases} \frac{2\|\eta\|_\infty}{t + \|\eta\|_\infty} & \text{if } t < \|\eta\|_\infty, \\ 1 & \text{if } t \leq \|\eta\|_\infty. \end{cases}$$

on Y . We can find β -norms of neutrosophic norm $(\mathbf{T}_1, \mathbf{S}_1, \mathbf{U}_1)$ since it satisfies the condition (1.1.1). Thus,

$$\mathbf{T}_1(\eta, t) \geq \beta \iff \frac{t - \|\eta\|_\infty}{t + \|\eta\|_\infty} \geq \beta \iff \frac{1 + \beta}{1 - \beta} \|\eta\|_\infty \leq t,$$

and

$$\mathbf{S}_1(\eta, t) \leq 1 - \beta \iff \frac{2\|\eta\|_\infty}{t + \|\eta\|_\infty} \leq 1 - \beta \iff \frac{1 + \beta}{1 - \beta} \|\eta\|_\infty \leq t,$$

$$\mathbf{U}_1(\eta, t) \leq 1 - \beta \iff \frac{2\|\eta\|_\infty}{t + \|\eta\|_\infty} \leq 1 - \beta \iff \frac{1 + \beta}{1 - \beta} \|\eta\|_\infty \leq t.$$

This shows that

$$\|\eta\|_\beta = \inf\{t > 0 : \mathbf{T}_1(\eta, t) \geq \beta \text{ and } \mathbf{S}_1(\eta, t) \leq 1 - \beta, \mathbf{U}_1(\eta, t) \leq 1 - \beta\} = \|\eta\|_\infty.$$

Now, we show that the sequence $\eta = (\eta_p) = (\frac{1}{p})_{p=1}^\infty$ is WN-convergent but not SN-convergent. Since each $\|\cdot\|_\beta$ is equivalent to $\|\cdot\|_\infty$, clearly (η_p) is WN-convergent to 0. However, this convergence is not uniform in β . In fact, for given $\epsilon > 0$

$$\|\eta\|_\beta = \frac{1 + \beta}{1 - \beta} \|\eta\|_\infty < \epsilon \iff \frac{1 + \beta}{(1 - \beta)\epsilon} < 1.$$

Since $\|\eta\|_\infty = 1$ for $\eta = (\frac{1}{p})_{p=1}^\infty$. But this is not possible, since $\frac{1 + \beta}{(1 - \beta)\epsilon} \rightarrow \infty$ as $\beta \rightarrow 1$.

3.1 Neutrosophic Fréchet derivative (NFD)

Definition 3.8 Let $(Y, \mathbf{T}_1, \mathbf{S}_1, \mathbf{U}_1, \bullet, \star)$ and $(W, \mathbf{T}', \mathbf{S}', \mathbf{U}', \bullet, \star)$ be two NNS, $H \subseteq Y$ be an neutrosophic open subset, and $h : H \rightarrow W$ probably nonlinear. Then h is called strong(weak) neutrosophic Fréchet differentiable at $\eta_0 \in H$ if there exists a strongly (weakly) neutrosophic bounded linear operator T from $(Y, \mathbf{T}_1, \mathbf{S}_1, \mathbf{U}_1, \bullet, \star)$ to $(W, \mathbf{T}', \mathbf{S}', \mathbf{U}', \bullet, \star)$ such that for every $t > 0$

$$\begin{aligned}\lim_{\mathbf{T}_1(a,t) \rightarrow 1} \mathbf{T}'\left(\frac{h(\eta_0 + a) - h(\eta_0) - Ta}{1 - \mathbf{T}_1(a,t)}, t\right) &= 1 \text{ (SN(WN))}, \\ \lim_{\mathbf{S}_1(a,t) \rightarrow 0} \mathbf{S}'\left(\frac{h(\eta_0 + a) - h(\eta_0) - Ta}{1 - \mathbf{S}_1(a,t)}, t\right) &= 0 \text{ (SN(WN))}, \\ \lim_{\mathbf{U}_1(a,t) \rightarrow 0} \mathbf{U}'\left(\frac{h(\eta_0 + a) - h(\eta_0) - Ta}{1 - \mathbf{U}_1(a,t)}, t\right) &= 0 \text{ (SN(WN))}.\end{aligned}$$

In this condition, the operator T is said to be strong (weak) neutrosophic, or shot, SN(WN)-Fréchet derivative of h at η_0 and is indicated by $\mathcal{D}_{\text{SN(WN)}}h[\eta_0]$ is called SN(WN)-Fréchet differentiable on U if it is SN(WN)-Fréchet differentiable at every point of H . In this condition, $\eta \rightarrow \mathcal{D}_{\text{SN(WN)}}h[\eta]$ is a basis from H to $(\mathcal{F}(Y, W), \mathcal{F}'(Y, W))$, denoted by $\mathcal{D}_{\text{SN(WN)}}h$.

Theorem 3.3 A strongly (weakly) neutrosophic bounded linear operator \mathcal{A} is SN(WN)-Fréchet differentiable at every point η_0 & $\mathcal{D}_{\text{SN(WN)}}\mathcal{A}[\eta_0] = \mathcal{A}$.

Proof This is explicit since

$$\mathbf{T}'\left(\frac{\mathcal{A}(\eta_0 + a) - \mathcal{A}(\eta_0) - \mathcal{A}a}{1 - \mathbf{T}_1(a,t)}, t\right) = \mathbf{T}'(0, t) = 1,$$

and

$$\begin{aligned}\mathbf{S}'\left(\frac{\mathcal{A}(\eta_0 + a) - \mathcal{A}(\eta_0) - \mathcal{A}a}{\mathbf{S}_1(a,t)}, t\right) &= \mathbf{S}'(0, t) = 0, \\ \mathbf{U}'\left(\frac{\mathcal{A}(\eta_0 + a) - \mathcal{A}(\eta_0) - \mathcal{A}a}{\mathbf{U}_1(a,t)}, t\right) &= \mathbf{U}'(0, t) = 0,\end{aligned}$$

$\forall t > 0$. □

Proposition 3.2 If h is SN(WN)-Fréchet differentiable at $\eta_0 \in H$, then it is strong (weak) neutrosophic continuous at η_0 .

Proof We take the following inequalities. For given $t > 0$,

$$\begin{aligned}\mathbf{T}'(h(\eta) - h(\eta_0), t) &= \mathbf{T}'(h(\eta) - h(\eta_0) - Ta + Ta, t) \\ &\geq \mathbf{T}'(h(\eta) - h(\eta_0) - Ta, t(1 - \mathbf{T}_1(a, t))) \bullet (1 - \mathbf{T}'(Ta, t\mathbf{T}_1(a, t))) \\ &= \mathbf{T}'\left(\frac{h(\eta) - h(\eta_0) - Ta}{(1 - \mathbf{T}_1(a, t))}, t\right) \bullet \mathbf{T}'\left(\frac{Ta}{\mathbf{T}_1(a, t)}, t\right),\end{aligned}$$

and

$$\begin{aligned}
 S'(h(\eta) - h(\eta_0), t) &= S'(h(\eta) - h(\eta_0) - Ta + Ta, t) \\
 &\leq S'(h(\eta) - h(\eta_0) - Ta, t(1 - S_1(a, t))) \star (1 - S_1(Ta, tS'(a, t))) \\
 &= S'\left(\frac{h(\eta) - h(\eta_0) - Ta}{S_1(a, t)}, t\right) \star S'\left(\frac{Ta}{1 - S_1(a, t)}, t\right), \\
 U'(h(\eta) - h(\eta_0), t) &= U'(h(\eta) - h(\eta_0) - Ta + Ta, t) \\
 &\leq U'(h(\eta) - h(\eta_0) - Ta, t(1 - U_1(a, t))) \star (1 - U_1(Ta, tU'(a, t))) \\
 &= U'\left(\frac{h(\eta) - h(\eta_0) - Ta}{U_1(a, t)}, t\right) \star U'\left(\frac{Ta}{1 - U_1(a, t)}, t\right).
 \end{aligned}$$

Since h is $SN(WN)$ -Fréchet differentiable at $\eta_0 \in H$, it follows that

$$T'(h(\eta) - h(\eta_0), t) \geq 1 \bullet T'\left(\frac{Ta}{T_1(a, t)}, t\right),$$

and

$$\begin{aligned}
 S'(h(\eta) - h(\eta_0), t) &\leq 0 \star S'\left(\frac{Ta}{1 - S_1(a, t)}, t\right), \\
 U'(h(\eta) - h(\eta_0), t) &\leq 0 \star U'\left(\frac{Ta}{1 - U_1(a, t)}, t\right),
 \end{aligned}$$

where $T = \mathcal{D}_{SN(WN)}h[\eta_0]$. Therefore, h is strong(weak) neutrosophic continuous. \square

Theorem 3.4 Suppose $(Y, T_1, S_1, U_1, \bullet, \star)$ and $(W, T', S', U', \bullet, \star)$ are two NNS, $H \subseteq Y$ be an neutrosophic open subset, and $h : H \rightarrow W$. If h is SN -Fréchet differentiable at some $\eta_0 \in H$, then it is WN -Fréchet differentiable at some η_0 with the same derivative but not conversely. The proof of the above theorem follows directly from the Proposition 3.2. For the converse part, let us consider the following example.

Example 3.3 Let $Y = W = \ell_p$ ($1 \leq p < \infty$), the Banach space of all absolutely p -summable sequences with the norm $\|\eta\|_p = (\sum_n |\eta|^p)^{\frac{1}{p}}$. Define the functions

$$\begin{aligned}
 T_1(\eta, t) &= \begin{cases} \frac{t^2}{t^2 + 2\|\eta\|_p^p} & \text{if } t > 0, \\ 0 & \text{if } t < 0; \end{cases} \\
 T'(\eta, t) &= \begin{cases} 1, & \text{if } t > 0 \text{ and } t^2 > \|\eta\|_p^p, \\ 0, & \text{if } t < 0 \text{ and } t^2 \leq \|\eta\|_p^p; \end{cases}
 \end{aligned}$$

and

$$S_1(\eta, t) = \begin{cases} \frac{2\|\eta\|_p^p}{t^2 + 2\|\eta\|_p^p} & \text{if } t > 0, \\ 0 & \text{if } t < 0; \end{cases}$$

$$S'(\eta, t) = \begin{cases} 0, & \text{if } t > 0 \text{ and } t^2 > \|\eta\|_p^p, \\ 1, & \text{if } t < 0 \text{ and } t^2 \leq \|\eta\|_p^p; \end{cases}$$

$$U_1(\eta, t) = \begin{cases} \frac{2\|\eta\|_p^p}{t^2 + 2\|\eta\|_p^p} & \text{if } t > 0, \\ 0 & \text{if } t < 0; \end{cases}$$

$$U'(\eta, t) = \begin{cases} 0, & \text{if } t > 0 \text{ and } t^2 > \|\eta\|_p^p, \\ 1, & \text{if } t < 0 \text{ and } t^2 \leq \|\eta\|_p^p. \end{cases}$$

Then $(Y, \mathbf{T}_1, \mathbf{S}_1, \mathbf{U}_1)$ and $(W, \mathbf{T}', \mathbf{S}', \mathbf{U}')$ are neutrosophic norms on ℓ_p . Consider the shift operator $G(\eta) = G(\{\eta_1, \eta_2, \dots\}) = \{0, \eta_1, \eta_2, \dots\}$ on ℓ_p . It is simple to see that linear operator G is weakly bounded hence is weakly continuous from $(\ell_p, \mathbf{T}_1, \mathbf{S}_1, \mathbf{U}_1, \bullet, \star)$ into $(\ell_p, \mathbf{T}', \mathbf{S}', \mathbf{U}', \bullet, \star)$ but not strongly continuous; since G is linear, we get from Theorem 3.2 that $G = \mathcal{D}_{WN}G[\eta]$ for all $\eta \in \ell_p$, while $\mathcal{D}_{SN}G[\eta]$ does not exist.

Definition 3.9 A subset \mathcal{B} in $NNS(Y, \mathbf{T}, \mathbf{S}, \mathbf{U}, \bullet, \star, \circ)$ is called SN(WN)-compact if each sequence of elements of \mathcal{B} has an SN(WN)-convergent subsequence.

Definition 3.10 $(Y, \mathbf{T}_1, \mathbf{S}_1, \mathbf{U}_1, \bullet, \star)$ and $(W, \mathbf{T}', \mathbf{S}', \mathbf{U}', \bullet, \star)$ be two NNS and $h: Y \rightarrow W$. Then h is said to be SN(WN)-compact if for every neutrosophic bounded subset \mathcal{B} of Y , the subset $h(\mathcal{B})$ is relatively SN(WN)-compact, i.e., the closure of $h(\mathcal{B})$ is SN(WN)-compact.

Remark 3.1 By the same way as in the proof of [25, Theorem 5], we can prove that h is SN(WN)-compact if it maps every neutrosophic bounded sequence onto a sequence which has an SN(WN)-convergent subsequence. Therefore, an SN-compact operator is WN-compact but not conversely. For example, the identity operator on $(c_0, \mathbf{T}, \mathbf{S}, \mathbf{U}, \bullet, \star)$, in Example 3.3 is not SN-compact while it is WN-compact. Because $(\frac{1}{p})_{p=1}^\infty$ cannot have SN-convergent subsequence.

Theorem 3.5 $(Y, \mathbf{T}_1, \mathbf{S}_1, \mathbf{U}_1, \bullet, \star)$ and $(W, \mathbf{T}', \mathbf{S}', \mathbf{U}', \bullet, \star)$ be two WN-complete NNS and $h: Y \rightarrow W$ be nonlinear WN-compact operator. Let, for some $\eta_0 \in Y$, $\mathcal{D}_{WN}h[\eta_0] = \mathcal{A}$ exist. Then the linear operator \mathcal{A} is also WN-compact.

Proof Suppose $\eta_p \subset (Y, \mathbf{T}_1, \mathbf{S}_1, \mathbf{U}_1, \bullet, \star)$ is an arbitrary neutrosophic bounded sequence. \exists some $t_0 > 0$ and $k \in (0, 1)$ such that $\mathbf{T}_1(\eta_p, \frac{t_0}{2}) \geq 1 - k$ and $\mathbf{S}_1(\eta_p, \frac{t_0}{2}) \leq k$, $\mathbf{U}_1(\eta_p, \frac{t_0}{2}) \leq k$, for every +ve integer p . Let the sequence $(\eta_0 + \eta_p)_{k=1}^\infty$, and let us show that it is neutrosophic bounded. If we take $1 - p_1 = \mathbf{T}_1(\eta_p, \frac{t_0}{2}) \bullet 1 - p$ and $p_1 = \mathbf{S}_1(\eta_p, \frac{t_0}{2}) \star p$, $p_1 = \mathbf{U}_1(\eta_p, \frac{t_0}{2}) \star p$ then

$$\mathbf{T}_1(\eta_0 + \eta_p, t_0) \geq \mathbf{T}_1\left(\eta_0, \frac{t_0}{2}\right) \bullet \mathbf{T}_1\left(\eta_p, \frac{t_0}{2}\right) > \mathbf{T}_1\left(\eta_0, \frac{t_0}{2}\right) \bullet (1 - p) = 1 - p_1,$$

and

$$\mathbf{S}_1(\eta_0 + \eta_p, t_0) \leq \mathbf{S}_1\left(\eta_0, \frac{t_0}{2}\right) \star \mathbf{S}_1\left(\eta_p, \frac{t_0}{2}\right) < \mathbf{S}_1\left(\eta_0, \frac{t_0}{2}\right) \star p = p_1,$$

$$\mathbf{U}_1(\eta_0 + \eta_p, t_0) \leq \mathbf{U}_1\left(\eta_0, \frac{t_0}{2}\right) \star \mathbf{U}_1\left(\eta_p, \frac{t_0}{2}\right) < \mathbf{U}_1\left(\eta_0, \frac{t_0}{2}\right) \star p = p_1$$

for every positive integer p . Rest of the proof can be done on the same lines as in [26]. \square

4 Conclusion

The present paper introduces the Fréchet derivative of nonlinear operators between *NNS* and the boundedness of linear operators between neutrosophic normed spaces. Current work is an increase and extension of the work of Mursaleen et al. [27], i.e., in an *NNS* which is more regular than the *IFNS*. So that, one may await it to be a more practical, modest work in the domain of neutrosophic topology in modeling the inaccuracy and ambiguity of several subjects arising in many fields of engineering, economics, and science.

Acknowledgements

The authors would like to thank the referees and the editor for their careful reading and valuable comments.

Funding

This work is financially supported by Aligarh Muslim University, Aligarh, India.

Availability of data and materials

Not applicable.

Declarations

Competing interests

The authors declare no competing interests.

Author contributions

VAK carried out the mathematical studies, participated in the sequence alignment, and drafted the manuscript. MDK participated in the design of the study and performed the numerical analysis. All authors read and approved the final manuscript.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 28 April 2022 Accepted: 21 June 2022 Published online: 05 December 2022

References

1. Zadeh, L.A.: Fuzzy sets. *Inf. Control* **8**, 338–353 (1965)
2. Barros, L.C., Bassanezi, R.C., Tonelli, P.A.: Fuzzy modelling in population dynamics. *Ecol. Model.* **128**, 27–33 (2000)
3. Giles, R.: A computer program for fuzzy reasoning. *Fuzzy Sets Syst.* **4**, 221–234 (1980)
4. Feng, G., Chen, G.: Adaptive control of discrete-time chaotic systems: a fuzzy control approach. *Chaos Solitons Fractals* **253**, 459–467 (2005)
5. Fradkov, A.L., Evans, R.J.: Control of chaos: methods and applications in engineering. *Chaos Solitons Fractals* **29**, 33–56 (2005)
6. Atanassov, K.: Intuitionistic fuzzy sets. *Fuzzy Sets Syst.* **20**, 87–96 (1986)
7. Mursaleen, M., Mohiuddine, S.A.: Statistical convergence of double sequences in intuitionistic fuzzy normed spaces. *Chaos Solitons Fractals* **41**(5), 2414–2421 (2009)
8. Mursaleen, M., Mohiuddine, S.A., Edely, O.H.H.: On the ideal convergence of double sequences in intuitionistic fuzzy normed spaces. *Comput. Math. Appl.* **59**(2), 603–611 (2010)
9. Khan, V.A., Esi, A., Ahmad, M., Khan, M.D.: Continuous and bounded linear operators in neutrosophic normed spaces. *J. Intell. Fuzzy Syst.* **40**, 11063–11070 (2021). <https://doi.org/10.3233/JIFS-202189>
10. Hong, L., Sun, J.Q.: Bifurcations of fuzzy nonlinear dynamical systems. *Commun. Nonlinear Sci. Numer. Simul.* **1**, 1–12 (2006)
11. Katrasas, A.K.: Fuzzy topological vector spaces. *Fuzzy Sets Syst.* **12**, 143–154 (1984)
12. Kirişçi, M., Şimşek, N.: Neutrosophic normed spaces and statistical convergence. *J. Anal.* **28**, 1059–1073 (2020)
13. Smarandache, F.: A Unifying Field in Logics: Neutrosophic Logic. Neutrosophy, Neutrosophic Set, Neutrosophic Probability: Neutrosophic Logic: Neutrosophy, Neutrosophic Set, Neutrosophic Probability (2003)
14. Smarandache, F.: Neutrosophic set-a generalization of the intuitionistic fuzzy set. *Int. J. Pure Appl. Math.* **24**(3), 287 (2005)
15. Bera, T., Mahapatra, N.K.: On neutrosophic soft linear spaces. *Fuzzy Inf. Eng.* **9**(3), 299–324 (2017)
16. Bera, T., Mahapatra, N.K.: Neutrosophic soft normed linear space. *Neutrosophic Sets Syst.* **23**(1), 1–6 (2018)
17. Khan, V.A., Daud, K., Mobeen, A.: A some results of neutrosophic normed spaces VIA Fibonacci matrix. *UPB Sci. Bull., Ser. A* **83**(2), 1–12 (2021)
18. Park, J.H.: Intuitionistic fuzzy metric space. *Chaos Solitons Fractals* **22**, 1039–1046 (2004)
19. Saadati, R.: A note on some results on the IF-normed spaces. *Chaos Solitons Fractals* **41**, 206–213 (2009)
20. Alimohammady, M., Roohi, M.: Compactness in fuzzy minimal spaces. *Chaos Solitons Fractals* **28**, 906–912 (2006)

21. Saadati, R., Park, J.H.: On the intuitionistic fuzzy topological spaces. *Chaos Solitons Fractals* **27**, 331–344 (2006)
22. Muralikrishna, P., Kumar, D.S.: Neutrosophic approach on normed linear space. *Neutrosophic Sets Syst.* **30**(1), 1–18 (2019)
23. Salama, A.A., Smarandache, F., Kromov, V.: Neutrosophic closed set and neutrosophic continuous functions. *Neutrosophic Sets Syst.* **4**, Article 2 (2014)
24. Kirişçi, M., Şimşek, N.: Neutrosophic metric spaces. [arXiv:1907.00798](https://arxiv.org/abs/1907.00798). arXiv preprint
25. Kaleva, O., Seikkala, S.: On fuzzy metric spaces. *Fuzzy Sets Syst.* **12**, 215–229 (1984)
26. Schweizer, B., Sklar, A.: Statistical metric spaces. *Pac. J. Math.* **10**, 313–334 (1960)
27. Mursaleen, M., Mohiuddine, S.A.: Nonlinear operators between intuitionistic fuzzy normed spaces and Fréchet derivative. *Chaos Solitons Fractals* **42**, 1010–1015 (2009)

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)

Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

onlineservice@springernature.com