



ICMME 2021



INTERNATIONAL CONFERENCE ON MATHEMATICS
AND MATHEMATICS EDUCATION

MATHEMATICS *in the footsteps of Gazi*

GAZİ UNIVERSITY - MATHEMATICIANS ASSOCIATION

BOOK OF ABSTRACTS

16-18 SEPTEMBER 2021

www.matder.org.tr / www.theicmme.org



International Conference on Mathematics and Mathematics Education
(ICMME - 2021)

Gazi University, Ankara, Turkey, 16-18 September 2021

©All rights reserved. Throughout this book, or part, without the permission of the author, electronic, mechanical, photocopying or reproduced in any recording system, transmitted, stored.

All kinds of information in this book is the responsibility of the authors.

Editors

Nurettin DOĞAN
Selçuk University, Turkey

Mustafa ÖZKAN
Gazi University, Turkey

Assistant Editors

Güler Başak ÖZNUR
Gazi University, Turkey

Elif Ece DEMİR
Gazi University, Turkey

Gökçe EFEOĞLU
Gazi University, Turkey

PREFACE

International Conference on Mathematics and Mathematics Education (ICMME-2021) was held on 16-18 September 2021 in Ankara, Turkey, entirely online via the zoom platform.

MATDER-Association of Mathematicians is an association founded in 1995 by mathematicians in Turkey. Up to now 14 national and 2 international mathematics symposium were organized by MATDER.

These meetings have been one of the main national symposiums. Since the talks in the meetings covers almost all areas of mathematics, mathematics education and engineering mathematics, the conferences have been well attended by mathematicians from academia, Ministry of Education and engineers as well. The last five conferences have been held in Konya (ICMME-2019), Ordu (ICMME-2018), Şanlıurfa (ICMME-2017), Elazığ (ICMME-2016) and Niğde (2015). This year ICMME-2021 has been held at Gazi University in Ankara/Turkey on 16-18 September 2021 as an international conference.

The main aim of this conference is to contribute to the development of mathematical sciences, mathematical education, and their applications and to bring together the members of the mathematics community, interdisciplinary researchers, educators, mathematicians, and statisticians from all over the world. The conference will present new results and future challenges, in series of invited and short talks, poster presentations, workshops, and exhibitions. All presented paper's abstracts will be published in the conference proceeding. Moreover, selected and peer review articles will be published in the following journals:

- Turkish Journal of Mathematics & Computer Science (TJMCS)
- MATDER Matematik Eğitim Dergisi

This conference is organised by MATDER-Association of Mathematicians and Gazi University.

On Behalf of The Organizing Committee
Hasan Hüseyin SAYAN

COMMITTEES

HONORARY AND ADVISORY BOARD

Prof. Dr. Musa YILDIZ	Rector of Gazi University, Turkey
Prof. Dr. Ömer AKIN	Honorary chairman of MATDER

SYMPOSIUM PRESIDENTS

H. Hüseyin SAYAN	Chairman of the Board, Turkey
Naim TUĞLU	Head of Mathematics Department, Gazi University, Turkey

ORGANIZING COMMITTEE

H. Hüseyin SAYAN (President)	Conference Chairman-MATDER President, Gazi University, Turkey
Naim TUĞLU (President)	Conference Chairman , Gazi University, Turkey
Mustafa ÖZKAN (Vice President)	Vice Conference Chairman, MATDER, Gazi University, Turkey
Nurettin DOĞAN (Vice President)	Vice Conference Chairman, MATDER, Selçuk University, Turkey
Abdülhamit KÜÇÜKASLAN	MATDER, Pamukkale University, Turkey
Ali ÖZTURAN	MATDER, Turkey
Aydın İZGİ	Harran University, Turkey
Başak KARPUZ	Dokuz Eylül University, Turkey
Cengiz ÇINAR	MATDER, Gazi University, Turkey
Devrim ÇAKMAK	Gazi University, Turkey
Ercan ÇELİK	Atatürk University, Turkey
Fatma AYAZ	MATDER, Gazi University, Turkey
Gülây KORU YÜCEKAYA	Selçuk University, Turkey
Hacı Mehmet BAŞKONUŞ	Harran University, Turkey
Hasan BULUT	Fırat University, Turkey

Hüseyin DEMİR	Samsun University, Turkey
Hüseyin YILDIRIM	Kahramanmaraş Sütçü İmam University, Turkey
İsmail EKİNCİOĞLU	Kütahya Dumlupınar University, Turkey
İsmet ALTINTAŞ	Kırgızistan-Türkiye Manas University, Turkey
Jehad Alzabut	Prince Sultan University, Turkey
Kemal AYDIN	Selçuk University, Turkey
M.Tamer KOŞAN	Gazi University, Turkey
Matthias Heil	University of Manchester, Turkey
Mohamed İbrahim Abbas	Alexandria University
Mohammad Esmael Samei	Bu-Ali Sina University, Iran
Mustafa KANDEMİR	Amasya University, Turkey
Nilay Akgönüllü PİRİM	MATDER, Turkey
Ömer ÖZEN	MATDER, Turkey
Sare ŞENGÜL	Marmara University, Turkey
Tahir FEYZİOĞLU	MATDER, Turkey
Truong Cong Quynh	The University of Danang-University of Science and Education
Yasin DEMİRCİ	MATDER, Turkey
Yusif S. Gasimov	Azerbaijan University, Azerbaijan
Zakia HAMMOUCH	Moulay Ismail University, Morocco
Ziya Ersin ARSLAN	MATDER, Turkey

LOCAL ORGANIZING COMMITTEE

Anıl ALTINKAYA	Gazi University, Turkey
Bilge Zehra SERGİ	Gazi University, Turkey
Çetin Cemal ÖZEKEN	Gazi University, Turkey
Duygu Gülşah BAŞARAN	Gazi University, Turkey
Elif Ece DEMİR	Gazi University, Turkey
Emre SEVGİ	Gazi University, Turkey
Fatma TOKMAK FEN	Gazi University, Turkey

Gökçe EFEOĞLU	Gazi University, Turkey
Gökhan MUTLU	Gazi University, Turkey
Güler Başak ÖZNUR	Gazi University, Turkey
Meryem KAYA	Gazi University, Turkey
Mustafa Fahri AKTAŞ	Gazi University, Turkey
Neslihan BİRİCİK	Gazi University, Turkey
Neslihan YILDIRIM	Gazi University, Turkey
Rümeysa GÖLCÜ ALTINKAYA	Gazi University, Turkey
Serdal YAZICI	Gazi University, Turkey
Sezin ÇİT	Gazi University, Turkey
Şule Yüksel GÜNGÖR	Gazi University, Turkey

SCIENTIFIC COMMITTEES

MATHEMATICS

A. Duran TÜRKOĞLU	Gazi University, Turkey
Abdullah MAĞDEN	Atatürk University, Turkey
Adil MISIR	Gazi University, Turkey
Aleksandr KOZLOV	Polotsk State University, Belarus
Ali DENİZ	Uşak University, Turkey
Ali Reza ASHRAFİ	University of Kashan, Iran
Alireza KHAŞTAN	Institute for Advanced Studies in Basic Sciences, Iran
Alper Cihan KONYALIOĞLU	Atatürk University, Turkey
Ayhan ERDEM	Gazi University, Turkey
Ayhan ŞERBETÇİ	Ankara University, Turkey
Aynur ARIKAN	Gazi University, Turkey
Aynur Keskin KAYMAKCI	Selçuk University, Turkey
Aysel VANLI	Gazi University, Turkey
Aytekin Bayram ÇIBIK	Gazi University, Turkey
Azhar Ali ZAFAR	University Lahore, Pakistan
Bahri TURAN	Gazi University, Turkey
Başak KARPUZ	Dokuz Eylül University, Turkey

Bayram ÇEKİM	Gazi University, Turkey
Bayram ŞAHİN	Ege University, Turkey
Birol ALTIN	Gazi University, Turkey
Cemil TUNÇ	Yüzüncüyıl University, Turkey
Cenap ÖZEL	King Abdulaziz University, Saudi Arabia
Cengiz ÇINAR	Gazi University, Turkey
Cüneyt ÇEVİK	Gazi University, Turkey
Çetin VURAL	Gazi University, Turkey
Dursun TAŞCI	Gazi University, Turkey
Ebenezer BONYAH	University of Education Winneba, Ghana
EL HASSAN EL KINANI	Moulay Ismail University, Morocco
Engin ÖZKAN	Erzincan University, Turkey
Ercan ALTINIŞIK	Gazi University, Turkey
Ercan ÇELİK	Atatürk University, Turkey
Erol KILIÇ	İnönü University, Turkey
Esma YILDIZ ÖZKAN	Gazi University, Turkey
Esra ERKUŞ DUMAN	Gazi University, Turkey
Esra KIR ARPAT	Gazi University, Turkey
F. Nejat EKMEKÇİ	Ankara University, Turkey
Fatma AYAZ	Gazi University, Turkey
Fatma TOKMAK FEN	Gazi University, Turkey
Ferhan ŞOLA ERDURAN	Gazi University, Turkey
Fikret GÖLGELEYEN	Bülent Ecevit University, Turkey
Fikriye Nuray YILMAZ	Gazi University, Turkey
Gennadii DEMIDENKO	Sobolev Institute of Mathematics, Russia
Gül ÖZKAN KIZILIRMAK	Gazi University, Turkey
Gülray KORU YÜCEKAYA	Selçuk University, Turkey
Gürhan İÇÖZ	Gazi University, Turkey
Hacı Mehmet BAŞKONUŞ	Harran University, Turkey
Hakan EFE	Gazi University, Turkey
Hammad KHALİL	University of Malakand, Pakistan

Hasan BULUT	Fırat University, Turkey
Hasan Hilmi HACISALİHOĞLU	President Of TWMS, Turkey
Hasan ÖĞÜNMEZ	Afyon Kocatepe University, Turkey
Hatice Gül İLARSLAN	Gazi University, Turkey
Hüseyin DEMİR	Ondokuz Mayıs University, Turkey
Hüseyin YILDIRIM	Kahramanmaraş Sütçü İmam University, Turkey
Ilhame AMIRALI	Düzce University, Turkey
Inessa MATVEEVA	Sobolev Institute of Mathematics, Russia
İbrahim YALÇINKAYA	Necmettin Erbakan University, Turkey
İsmail EKİNCİOĞLU	Dumlupınar University, Turkey
İsmet ALTINTAŞ	Kırğızistan-Türkiye Manas University, Kyrgyzstan
İsmet ALTINTAŞ	Sakarya University, Turkey
İsmet YÜKSEL	Gazi University, Turkey
Jan ZEMLICKA	Charles University, Czechia
Jehad ALZABUT	Prince Sultan University, Saudia Arabia
Kazım İLARSLAN	Kırıkkale University, Turkey
Kemal AYDIN	Selçuk University, Turkey
Keziban ORBAY	Amasya University, Turkey
Luca GRILLI	Foggia University, Italy
Luis Manuel SÁNCHEZ RUIZ	Universitat Politècnica de València, Spain
M.Tamer KOŞAN	Gazi University, Turkey
Mahmut ERGÜT	Namık Kemal University, Turkey
Martin BOHNER	Missouri University Of Science And Technology, USA
Mediha ÖRKÇÜ	Gazi University, Turkey
Mehmet MERDAN	Gümüşhane University, Turkey
Mehmet YILDIRIM	Kırıkkale University, Turkey
Meryem KAYA	Gazi University, Turkey
Mohamed İbrahim ABBAS	Alexandria University, Egypt
Mohammad Esmael SAMEI	Bu-Ali Sina University, Iran
Mustafa ÇALIŞKAN	Gazi University, Turkey
Mustafa Fahri AKTAŞ	Gazi University, Turkey

Mustafa ÖZKAN	Gazi University, Turkey
Mustafa YILDIZ	Bülent Ecevit University, Turkey
Müzeyyen SANGURLU SEZEN	Gazi University, Turkey
Naim TUĞLU	Gazi University, Turkey
Necdet BİLDİK	Celal Bayar University, Turkey
Nur ALAM	Pabna University of Science & Technology, Bangladesh
Nurhayat İSPİR	Gazi University, Turkey
Ogün DOĞRU	Gazi University, Turkey
Oktay MUHTAROĞLU	Gaziosmanpaşa University, Turkey
Ömer TARAKÇI	Atatürk University, Turkey
Saeid JAFARI	College of Vestsjaelland South Denmark, Denmark
Snezhana GOCHEVA-ILIEVA	University of Plovdiv Paisii Hilendarski, Bulgaria
Şerife BÜYÜKKÖSE	Gazi University, Turkey
Truong Cong QUYNH	The University of Danang-University of Science and Education, Vietnamese
Ülkü DİNLEMEZ KANTAR	Gazi University, Turkey
Vagif GULIYEV	Kütahya Dumlupınar University, Turkey
Yusif S. GASIMOV	Azerbaijan University, Azerbaijan
Yusuf GÜREFE	Uşak University, Turkey
Yusuf YAYLI	Ankara University, Turkey
Zafar AHSAN	Aligargah Muslim University, India
Zakia HAMMOUCH	Moulay Ismail University, Morocco

MATHEMATICS EDUCATION

Ali ERASLAN	Samsun Ondokuz Mayıs University, Turkey
Ahmet DOĞAN	Uşak University, Turkey
Abdulkadir TUNA	Kastamonu University, Turkey
Cemil İNAN	Artuklu University, Turkey
Şenol DOST	Hacettepe University, Turkey
Hasan ES	Gazi Üniversitesi, Turkey
Hatice AKKOÇ	Marmara University, Turkey
Hülya GÜR	Balıkesir University, Turkey

Melek MASAL	Sakarya University, Turkey
Murat ALTUN	Uludağ University, Turkey
Murat BEKAR	Gazi University, Turkey
Murat PEKER	Afyon Kocatepe University, Turkey
Mustafa DOĞAN	Selçuk University, Turkey
Mustafa KANDEMİR	Amasya University, Turkey
Nurullah YAZICI	Karamanoğlu Mehmetbey University, Turkey
Osman GÜRSOY	Maltepe University, Turkey
Ömer DERELİ	Necmettin Erbakan University, Turkey
Ömer Faruk TEMİZER	Fırat University, Turkey
Özge YİĞİTCAN NAYIR	Başkent University, Turkey
Safure BULUT	Middle East Technical University, Turkey
Sare ŞENGÜL	Marmara University, Turkey
Sebahat YETİM	Gazi University, Turkey
Şeref MİRASYEDİOĞLU	Başkent University, Turkey
Sibel YEŞİLDERE İMRE	Dokuz Eylül University, Turkey
Süha YILMAZ	Dokuz Eylül University, Turkey
Tunay BİLGİN	Van Yüzüncü Yıl University, Turkey
Yasemin KATRANCI	Kocaeli University, Turkey
Yüksel DEDE	Gazi University, Turkey
Zaki F.A. El-RAHEM	Alexandria University, Egypt

CONTENTS

INVITED SPEAKERS	1
AN ASSESSMENT ON THE PLACE OF MATHEMATICAL KNOWLEDGE BETWEEN FORM AND METAPHYSICS.....	2
An efficient algorithm to solve nonlinear reaction-diffusion problems with non-integer order derivatives.....	4
Flows past cylinders: Are the transitions between different flow regimes caused by a continuous evolution or by bifurcations?	5
Sayı Duyusunun Gelişimi ve Matematiksel Yeterlikler	6
Self-small products of abelian groups.....	7
Time-dependent source identification problem for Schrodinger differential and difference equations	8
ABSTRACTS OF ORAL PRESENTATIONS.....	9
ALGEBRA AND NUMBER THEORY.....	10
A Generalization of Gaussian Pell-Lucas Polynomials	11
A Note on Sum Formulas of Generalized Hexanacci Numbers: Closed Forms of the Sum Formulas $\sum_{k=0}^n kx^k W_k$ and $\sum_{k=1}^n kx^k W_{-k}$	13
A Note on Sum Formulas of Generalized Pentanacci Sequence: Closed Forms of the Sum Formulas $\sum_{k=0}^n kx^k W_k$ and $\sum_{k=1}^n kx^k W_{-k}$	15
Complex (Gaussian) q -Fibonacci and Complex (Gaussian) q -Lucas Quaternions.....	16
Groups Whose Proper Factor Groups Are Fitting	19
LINEAR MAPS PRESERVING DRAZIN INVERSES OF	21
MATRICES OVER LOCAL RINGS	21
N-Barely Transitive Groups	22
NC-Barely Transitive Groups	24
On The Catalan Transforms of The Incomplete Jacobsthal Numbers.....	26
On the Finitary Permutation Groups.....	28

SIP-direct-injective Modules and Rings.....	29
Some Notes on Soft Modules in Soft Sets.....	30
Some properties of Apostol Bernoulli Fibonacci and Apostol Euler Fibonacci Polynomials.....	32
The Universal Difference Property for Generalized Splines over Principal Ideal Domains.....	35
On semi-Nekrasov matrices	36
ANALYSIS.....	38
An Extension of the Adams-type Theorem to Vanishing Generalized Weighted Morrey Spaces.....	39
Approximation by Some Kantorovich Type Max-Product Operators	41
APPROXIMATION PROPERTIES OF GENERALIZED JAIN OPERATORS OF TWO VARIABLES	43
Generalized Fractional Integral Inequalities on Convex Functions.....	45
Matrix Transformations and Some Algebraic and Topological Properties Related to the New Catalan Double Sequence Space	47
On Absolute Fibonacci Series Space And Matrix Operators	49
On the Absolute Euler Summability Factors	50
On The Multilinear Riesz Bessel Transforms On Morrey Spaces	51
Rate of Convergence by Bivariate	53
Some Mean Ergodic Type Theorems.....	55
Some Operator Identities on the Umbral Algebra.....	57
Some Results Based on (p, r) –Compact Sets in Banach Spaces	59
Some Results Related to New Jordan Totient Double Sequence Spaces.....	61
Spanne-type Result for the Generalized Fractional Maximal Operator in the Vanishing Generalized Weighted Local Morrey Spaces.....	63
Upper and lower θ_p-continuous multifunctions.....	65
APPLIED MATHEMATICS.....	66

A Mathematical Investigation of COVID-19 in Turkey Using a Susceptible Infectious Recovered (SIR) Type Model	67
A NOTE ON MONOTONE TECHNIQUE FOR A CLASS OF FRACTIONAL INTEGRO-DIFFERENTIAL EQUATIONS	69
A Slough/Wound Interaction Model Accounting for Cellular Diffusion.....	71
Analysis of Continuous Data Assimilation Scheme for the Navier-Stokes Equations Using Variational Multiscale Method	73
Analysis of Covid-19 epidemic in Turkey	75
Decay Estimate for the Klein-Gordon Equation in Anti de Sitter Spacetime	77
Dynamical Behavior of Rational Difference Equation	79
Existence Of Solutions For A Third-Order Boundary Value Problem With Integral Boundary Conditions On Infinite Intervals	81
Fitted Numerical Method for a Volterra Delay Integro-Differential Equation.....	83
Fuzzy Similarity Of Subalpine, Steppe And Segetal Vegetation For Black Sea Region In Turkey	85
Global Existence and Uniqueness of The Inviscid Velocity-Vorticity Model of The g-Navier-Stokes Equations.....	87
Hermite-Hadamard -Fejér type inequalities for higher order convex functions and quadrature formulae	89
Implicit Exponential Finite Difference Method for Numerical Solutions of Modified Burgers Equation.....	91
Modified Finite Difference Method for solution of two-interval boundary value problems with transition conditions	92
New Fractional Operators Including Wright Function in Their Kernels	94
Nonexistence of Global Solutions for a Hyperbolic-Type Wave Equation With Delay	96
Numerical Analysis of Singularly Perturbed Problems with Integral Boundary Conditions	97
Numerical Integration Of System Of Differential Equations By Iterative Decreasing Dimension Algorithm	99
On Controlling Chaos for Multi-Step Iteration Process and Its Special Iterations in Discrete Dynamical Systems.....	101

On Some Applications of Korovkin Type Theorem	103
On stability of the bacterial infection model through incommensurate fractional-order system	105
Parametric Identification in ODE Models as a Tool for Solving Real-world Problems.....	107
Several Weighted Hardy type Inequalities with Robin Boundary Conditions..	109
Some Further Properties of BLUPs in SUR Models.....	111
Some Remarks on Predictors in Reduced Linear Models.....	113
Sub Pico-Second Optical Solitons in Birefringent Fibers with Kaup-Newell Equation by Two Strategic Integration Technologies	115
The Asymptotic Behavior for Generalized λ - Bernstein-Stancu Type Operators.....	117
The Marshall–Olkin transmuted Rayleigh distribution. Statistical properties and different methods of estimation	119
The Poisson Topp Leone generalized exponential distribution	121
GEOMETRY	123
An Extension of Smarandache Ruled Surfaces According.....	124
Bertrand Curves and B-Lift Curves in Lorentzian 3-Space	126
Generating Circle-Foliated Transition Surfaces in Lorentz Space	127
On Some Closed Sets and Circles of the Added Sierpinski Triangle.....	128
Some Notes on $N(k)$ -contact Metric Manifolds.....	130
Some Special Smarandache Ruled Surfaces According.....	132
The properties of projective, concircular and conharmonic curvature tensor fields on the complex Sasakian manifold.....	134
MATHEMATICS EDUCATION	135
A Disease Modeling Approach to Simulate the Spread of Math Anxiety Between Students.....	136
An Application On Comparison of Student Achievements in Face-to-Face and Online Education Models in University Mathematics Education	138

Behavioural Analysis of a Predator-Prey Model	140
Development of Preservice Mathematics Teachers' TPACK through Digital Storytelling	142
Examining the Effects of Pre-school Education on Mathematics Achievement at Different Socioeconomic Levels: The Case of TIMSS 2019 Turkey	144
On Stability Analysis and Flip Bifurcation of a Discrete-Time Predator-Prey System with Allee Effect and Prey Immigration	147
ON THE DEVELOPMENT OF PLANNING ABILITY IN PROBLEM SOLVING FROM GRADES 5 THROUGH 7	149
Pre-service and In-service Teachers' Proof Schemes and Their Opinions on Mathematical Proof	150
Preservice High School Mathematics Teachers' Experiences on their Fieldwork During the COVID-19 Pandemic	153
The Effects Of Activities Carried Out With Web 2.0 Tools On Some Field Competencies Of Mathematics Teacher Candidates	155
The Effectiveness of the Subject Area Textbook Review Course Amid the COVID-19 Pandemic	157
The Field Experiences on the Online Application of the Case Study Method in Mathematics Lesson	159
The Investigation of Studies on Case Study Method in Education	161
The Opinions of Secondary School Students' Parents about Mathematics Course in Distance Education	163
Translation of the Digital Game Critter Corral into Turkish and Its Evaluation in Terms of Mathematics Education Standards and Appropriateness for Children	165
LISTS OF PARTICIPANTS OF ICMME-2021	167

INVITED SPEAKERS

AN ASSESSMENT ON THE PLACE OF MATHEMATICAL KNOWLEDGE BETWEEN FORM AND METAPHYSICS

Vatan KARAKAYA¹

1.Ahi Evran University, Mathematics Department, vkarakaya@ahievran.edu.tr

ABSTRACT

It is known that the source of knowledge, which is the subject of thinking, is the realm of concrete or intangible existence. For this reason, the issue of how knowledge is produced from existence or how existence is the subject of knowledge has been one of the main subjects of philosophy throughout the history of thought. The approach of philosophy on this subject has led to finding the unchanging principles of existence, that is to seek universal truths. These orientations and pursuits have become an alliance in the fact that the knowledge of God is unchangeable and imperative, especially Aristotle and then Ibn Sînâ. The fact that the knowledge of God is logically necessary and unchangeable rather than ontological has necessitated the search for unchangeable principles on existence. Immutable principles on existence are possible through the forms possessed by the being. The lexical meaning of the word form is "shape", but it has been described as "the nature of something, the substance that makes it that thing" [1]. Depending on the philosophical definition of the form, the generally accepted definition given by Ibn Sînâ is "the substance that takes the thing it is from into the realm of action" [2]. According to Aristotle, who accepts the principle of existence as matter (heyula) and form, existence consists of the form realized in matter [3]. The first matter, which is considered as a potential force and a possibility for the emergence of existence, comes into existence only after it takes a form. The entity that takes on a form becomes available for classification as the same genus, the same species or the same chapter, and becomes perceptible by being divided into categories. Man has divided the realm of forms into different groups by using the cognitive features he has from his creation. The first group is the physical world of objects, in which the entity is open to sensory description and definitions of physical objects can be made with the forms obtained from physical objects. According to Aristotle's view adopted in Islamic

Peripatetic philosophy, in order to understand what a thing is physically, it is necessary to know the reason for it and go as far as the first causes. In that case, four reasons must be sought for the realization of all kinds of phenomena and events that occur in the universe. These are matter, form, agent and objective causes. For obtaining information from non-physical entities, the form reason constitutes an important basis. The most general field of knowledge that is not physical and uses form is mathematical knowledge. Mathematical objects have form but are non-physical entities closed to the senses. In this study, how the being becomes the subject of mathematics by gaining form and the relationship between mathematical entities and the form will be examined. How mathematical entities correspond to counting and measuring through form and how we do mathematics through this relationship will be evaluated both in terms of classical philosophy and through modern approaches. Due to the dependence of mathematical and physical knowledge on logically necessary existence knowledge, the possibilities of metaphysical knowledge will also be briefly mentioned. In particular, it will be investigated whether the quantitative approach of Descartes in natural philosophy and the commitment to intuition in Kant's philosophy have a relationship with corporeal form ([4],[5]).

Key Words: Knowledge, Mathematical object, Mathematical form

An efficient algorithm to solve nonlinear reaction-diffusion problems with non-integer order derivatives

Zakia HAMMOUCH¹

1. Moulay Ismail University Morocco, Department of Mathematics,
hammouch.zakia@gmail.com

ABSTRACT

In this talk present a numerical analysis of two types of nonlinear reaction-diffusion problems with periodic conditions and initial conditions, where the order of differentiation is arbitrary. Two iterative schemes are constructed using the upper and lower solutions method, which converges monotonically towards a maximum solution, or a minimum solution of the problem considered when the mesh decreases to zero, depending on whether the initial iteration is an upper solution or a lower solution. Besides, a result for comparing the different monotonic sequences is stated proved and theoretical results are supported by some numerical simulations.

Key Words: Caputo derivative; Conformable derivative; Monotone iterative method; Nonlinear parabolic PDE, Upper and lower solutions.

Flows past cylinders: Are the transitions between different flow regimes caused by a continuous evolution or by bifurcations?

Matthias HEIL¹

¹The University of Manchester, Department of Mathematics, matthias.heil@manchester.ac.uk

ABSTRACT

Solutions to the Navier-Stokes equations often go through a sequence of distinct regimes, with the flow field becoming more "complicated" as the Reynolds number increases. These changes may occur via (i) bifurcations of the underlying solutions of the Navier-Stokes equations, or (ii) a continuous evolution of the "complicated" flow field (with quantifiable, discrete changes to its topology).

We analyse the interplay between these two, in principle distinct, mechanisms in the context of flows past circular cylinders. If the cylinder is stationary the flow undergoes a Hopf bifurcation at a Reynolds number of approximately 46, resulting in the formation of the famous von Kármán vortex street -- a time-periodic flow in which vortices are shed downstream. While this suggests that the change to the flow topology arises via mechanism

(i) we show that the transition from steady to time-periodic flow (through the Hopf bifurcation) and the formation of individual vortices are in fact distinct events that occur at slightly different Reynolds numbers.

When the cylinder performs forced oscillations transverse to the flow direction, the vortex-shedding pattern becomes significantly more complex, leading to the formation of so-called "exotic wakes" whose character is controlled by the Reynolds number as well as the period and amplitude of the cylinder's motion. While it has generally been assumed that the transition between different wake patterns in response to changes in the amplitude occurs via mechanism

(ii) we show that they are actually associated with a spatio-temporal symmetry-breaking bifurcation of the time-periodic flow.

Sayı Duyusunun Gelişimi ve Matematiksel Yeterlikler

Erdinç Çakıroğlu¹

1Middle East Technical University, Department of Mathematics and Science Education,
erdinc@metu.edu.tr

ABSTRACT

Bireylerde sayı duyusu erken çocukluk, hatta bebeklik döneminden itibaren başlar ve okul hayatı boyunca gelişir. İlkokuldan itibaren sayı duyusunun matematik derslerinde desteklenmesi ve geliştirilmesi öğretim programının temel amaçlardan birisidir. Fakat ilkokulun ilk yıllarında matematik öğretimindeki eksiklikler sonraki yıllarda karşılaştığımız matematik öğrenme sorunlarının temelini oluşturmaktadır. Matematikte işlemleri doğru yapabilmek ve kavramlara hakim olmak önemli olsa da matematikte yetkin bireyler yetiştirebilmek için yeterli değildir. Matematikte sağlam bir bilgi alt yapısı oluşturabilmek için bireylerin kazanması gereken birtakım yeterlikler bulunmaktadır. NRC (2001) tarafından önerilen bu yeterlikler (1) kavramsal anlama, (2) işlemsel akıcılık, (3) stratejik yetkinlik, (4) uyarlanabilir muhakeme ve (5) üretken eğilim olmak üzere beş ana başlık altında tanımlanmaktadır. Matematik derslerinde bu 5 yeterliğin birbirine paralel olarak eş zamanlı ele alınması ve öğrencilerin bu yeterlikleri kazanarak matematiği öğrenmesi daha güçlü bir matematiksel temel oluşmasına yardımcı olmaktadır. İlk bakışta bu yeterlikleri öğrencilere kazandırmak ütöpik gibi görünse de kazanılan her bir yeterliğin diğerini de beslediği ve bunların sınıflarda uygulanabilir olduğu çeşitli araştırmalarda gösterilmiştir. Bu çalışmada matematiksel yeterliklerin çocuklarda sayı duyusunun gelişimine nasıl katkı sağladığı hem kuramsal hem de uygulamalı olarak açıklanacaktır. Ayrıca her bir yeterliğin diğerlerinin gelişimine nasıl katkı sağladığı araştırma bulgularından örneklerle paylaşılacaktır. Ayrıca konuyla ilgili bir TÜBİTAK projesinin araştırma çıktılarından örnekler de paylaşılacaktır.

SELF-SMALL PRODUCTS OF ABELIAN GROUPS

Jan Zemlicka¹

¹Charles University, zemlicka@karlin.mff.cuni.cz

ABSTRACT

Let A and B be two abelian groups. The group A is called B -small if the covariant functor $\text{Hom}(A, -)$ commutes with all direct sums of the form $B(k)$ and A is a self-small group provided it is A -small. The main aim of the talk is to characterize self-small products applying developed closure properties of the classes of relatively small groups. In particular, we show that a product of a system of abelian groups is self small if and only if it relatively small over a direct sum of the system.

As a consequence of the theory of relatively small groups and the well-known fact that powers Z_k of the group Z of all integers is slender for any nonmeasurable cardinal k , we characterize self-small products of finitely generated abelian groups. Namely, the product M of finitely generated groups is self-small if and only if either M is isomorphic to power Z_k for some cardinal k , or M is isomorphic to a direct sum of a finitely generated free group F and finite abelian p -groups for each prime number p .

Finally, we also discuss possible application of the developed tools for description of self-compact objects in context of general additive and abelian categories.

Time-dependent source identification problem for Schrodinger differential and difference equations

Allaberen ASHYRALYEV¹

¹Near East University, Faculty of Science and Literature, allaberen.ashyralyev@neu.edu.tr

ABSTRACT

Source identification problems(SIPs) have the significant role in natural science, applied sciences, engineering, quantum mechanics, diffusion equations, heat equations (see, e.g., [1]-[4]). This is a discuss, the time-dependent SIP for the Schrödinger equation

$$\begin{cases} i \frac{du(t)}{dt} + Au(t) = p(t)q + f(t), t \in (0, T) \\ u(0) = \varphi, B[u(t)] = \tau(t), t \in [0, T] \end{cases} \quad (1)$$

In a Hilbert space H with the self-adjoint positive definite operator A with the dense domain $D(A)$ in H is studied. Here, $B: H \rightarrow \mathbb{R}$ is a given linear bounded functional and $\tau(t): [0, T] \rightarrow \mathbb{R}$ is a given smooth function and $q \in D(A), Bq \neq 0$.

Single-step absolute stable difference schemes for the numerical solution of time-dependent SIP are presented. The main theorems on stability estimates for the solutions of these problems are established. In practice, the stability estimates for solution of four problems for Schrödinger differential and difference equations are proved. Numerical results and explanatory illustrations are presented show the validation of the theoretical results.

ABSTRACTS OF ORAL PRESENTATIONS

ALGEBRA AND NUMBER THEORY

A Generalization of Gaussian Pell-Lucas Polynomials

Engin Özkan¹ and Mine Uysal²

¹ Department of Mathematics, Faculty of Arts and Sciences,
Erzincan Binali Yıldırım University, Erzincan, 24100, Turkey
eozkan@erzincan.edu.tr

² Graduate School of Natural and Applied Sciences,
Erzincan Binali Yıldırım University, Erzincan, 24100, Turkey.
mine.uysal@erzincan.edu.tr

ABSTRACT

Number sequences and their polynomials have attracted the attention of many scientists for many years, as they find application in nature and in many sciences. Of course, the Fibonacci numbers are the best known of the sequences of numbers [3-5,7]. Many generalizations of number sequences were then described and studied [3-7]. One of the most important of these generalizations is those about Gaussian. Özkan et al. defined Gauss Fibonacci polynomials, Gauss Lucas polynomials and gave their applications in [4].

Now, let us give some basic definitions for this paper in this section. We know well that the Pell numbers P_n are defined by

$$P_n = 2P_{n-1} + P_{n-2}, \text{ for } n \geq 3$$

with $P_1 = 1$ and $P_2 = 2$. [2]

Similarly, the Pell-Lucas numbers L_n ,

$$Q_n = 2Q_{n-1} + Q_{n-2}, \text{ for } n \geq 3$$

with $Q_1 = 1$ and $Q_2 = 3$. [2]

Pell polynomials are defined by respectively,

$$P_{n+2}(x) = 2xP_{n+1}(x) + P_n(x)$$

with $P_0(x) = 0$ and $P_1(x) = 1$. [2]

Pell-Lucas polynomials are defined by respectively,

$$Q_{n+2}(x) = 2xQ_{n+1}(x) + Q_n(x)$$

with $Q_0(x) = 2$ and $Q_1(x) = 2x$. [2]

In [1], the authors introduced the Gaussian Pell polynomials and examined their properties. Later, Gaussian Pell-Lucas polynomials are defined in [8] and given their properties.

One of the latest works in this area is [7] where it is introduced d-Fibonacci and d-Lucas polynomials.

We define d -Gaussian Pell-Lucas polynomials. Then we present the sum, generating functions and Binet formulas of these polynomials. We give the matrix representations of d -Gaussian Pell-Lucas polynomials.

Key Words: d -Gaussian Pell-Lucas polynomials, Generating Function, Binet Formula, d -Gaussian Pell-Lucas polynomials matrix.

REFERENCES

- [1] S. Halici and S. Oz, On Gaussian Pell polynomials and their some properties, Palestine Journal of Mathematics, 7(1), (2018), 251-256.
- [2] T. Koshy, Pell and Pell-Lucas numbers with Applications, New York: Springer, 2014.
- [3] E. Özkan, On Truncated Fibonacci sequences, Indian J. Pure of And App. Mathematics 38(4) (2007), 241-251.
- [4] E. Özkan and M. Taştan, On Gauss Fibonacci polynomials, on Gauss Lucas polynomials and their applications, Communications in Algebra 48(3) (2020), 952-960.
- [5] E. Özkan and M. Taştan, On Gauss k-Fibonacci polynomials, Electronic Journal of Mathematical Analysis and Applications 9(1) (2021), 124-130.
- [6] E. Özkan and B. Kuloğlu, On the new Narayana polynomials, The Gauss Narayana numbers and their polynomials, Asian-European Journal of Mathematics 14 (2021), 1-10.
- [7] B. Sadaoui, A. Krelifa, d-Fibonacci and d-Lucas polynomials, Journal of Mathematical Modeling 9(3) (2021), 425-436.
- [8] T. Yağmur, Gaussian Pell-Lucas Polynomials, Communications in Mathematics and Applications, 10(4) (2019), 673-679.

A Note on Sum Formulas of Generalized Hexanacci Numbers: Closed Forms of the Sum Formulas $\sum_{k=0}^n kx^k W_k$ and $\sum_{k=1}^n kx^k W_{-k}$

Yüksel Soykan ¹, Erkan Taşdemir ² and Tülin Erdoğan Taşdemir ³

1 Department of Mathematics, Art and Science Faculty, Zonguldak Bülent Ecevit University,
67100, Zonguldak, Turkey.

yuksel_soykan@hotmail.com

2 Pınarhisar Vocational School, Kırklareli University, 39300, Kırklareli, Turkey

erkantasdemir@hotmail.com

3 Ministry of National Education, Pınarhisar Anatolian High School, Kırklareli, Turkey

tulintasdemir@hotmail.com

ABSTRACT

In this paper, closed forms of the sum formulas $\sum_{k=0}^n kx^k W_k$ and $\sum_{k=1}^n kx^k W_{-k}$ for generalized Hexanacci numbers are presented. As special cases, we give summation formulas of Hexanacci, Hexanacci-Lucas, and other sixth-order recurrence sequences.

Key Words: Hexanacci numbers, Hexanacci-Lucas numbers, sum formulas, summing formulas.

REFERENCES

- [1] G.P.S. RATHORE, O. SIKHWAL AND R. CHOUDHARY, FORMULA FOR FINDING NTH TERM OF FIBONACCI-LIKE SEQUENCE OF HIGHER ORDER, INTERNATIONAL JOURNAL OF MATHEMATICS AND ITS APPLICATIONS, 4 (2-D) (2016), 75-80.
- [2] L.R. NATIVIDAD, ON SOLVING FIBONACCI-LIKE SEQUENCES OF FOURTH, FIFTH AND SIXTH ORDER, INTERNATIONAL JOURNAL OF MATHEMATICS AND COMPUTING, 3(2) (2013), 38-40.
- [3] Y. SOYKAN, ON SUMMING FORMULAS OF GENERALIZED HEXANACCI AND GAUSSIAN GENERALIZED HEXANACCI NUMBERS, ASIAN RESEARCH JOURNAL OF MATHEMATICS, 14(4) (2019), 1-14.
- [4] Y. SOYKAN, A STUDY ON SUM FORMULAS OF GENERALIZED SIXTH-ORDER LINEAR RECURRENCE SEQUENCES, ASIAN JOURNAL OF ADVANCED RESEARCH AND REPORTS, 14(2) (2020), 36-48.

- [5]** Y. SOYKAN, A STUDY ON GENERALIZED (R,S,T,U,V,Y)-NUMBERS, JOURNAL OF PROGRESSIVE RESEARCH IN MATHEMATICS, 17(1) (2020) ,54-72.
- [6]** Y. SOYKAN AND N. ÖZMEN, ON GENERALIZED HEXANACCI AND GAUSSIAN GENERALIZED HEXANACCI NUMBERS, ACCEPTED.

A Note on Sum Formulas of Generalized Pentanacci Sequence: Closed Forms of the Sum Formulas $\sum_{k=0}^n kx^k W_k$ and $\sum_{k=1}^n kx^k W_{-k}$

Yüksel Soykan ¹, Erkan Taşdemir ² and Melih Göcen ¹

¹ Department of Mathematics, Art and Science Faculty, Zonguldak Bülent Ecevit University,
67100, Zonguldak, Turkey,

yuksel_soykan@hotmail.com, gocenm@hotmail.com

² Pınarhisar Vocational School, Kırklareli University, 39300, Kırklareli, Turkey
erkantademir@hotmail.com

ABSTRACT

In this paper, closed forms of the sum formulas $\sum_{k=0}^n kx^k W_k$ and $\sum_{k=1}^n kx^k W_{-k}$ for generalized Pentanacci numbers are presented. As special cases, we give summation formulas of Pentanacci, Pentanacci-Lucas, and other fifth-order recurrence sequences.

Key Words: Pentanacci numbers, Pentanacci-Lucas numbers, sum formulas, summing formulas.

REFERENCES

- [1] G.P.S. Rathore, O. Sikhwal and R. Choudhary, Formula for finding nth Term of Fibonacci-Like Sequence of Higher Order, International Journal of Mathematics and its Applications, 4 (2-D) (2016), 75-80.
- [2] R.S. Melham, Some Analogs of the Identity $F_{2n} + F_{2n+1} = F_{22n+1}$, Fibonacci Quarterly, 305-311, 1999.
- [3] L.R. Natividad, On Solving Fibonacci-Like Sequences of Fourth, Fifth and Sixth Order, International Journal of Mathematics and Computing, 3(2) (2013), 38-40.
- [4] Y. Soykan, On Generalized Pentanacci and Gaussian Generalized Pentanacci Numbers, Asian Research Journal of Mathematics, 16(9) (2020), 102-121.
- [5] Y. Soykan, Properties of Generalized Fifth-Order Pell Numbers, Asian Research Journal of Mathematics, 15(3) (2019), 1-18, 2019.
- [6] Y. Soykan and E.E. Polatlı, A Note on Fifth Order Jacobsthal Numbers, IOSR Journal of Mathematics (IOSR-JM), 17(2) (2021), 1-23.
- [7] Y. Soykan, A Study On Generalized 5-primes Numbers, Journal of Scientific Perspectives, 4(3) (2020), 185-202.

Complex (Gaussian) q -Fibonacci and Complex (Gaussian) q -Lucas Quaternions

Tülay Yağmur

Department of Mathematics, Aksaray University, Aksaray, Turkey
tulayyagmur@aksaray.edu.tr

ABSTRACT

Quaternions, introduced by Irish mathematician William Rowan Hamilton, are a four-dimensional non-commutative algebra over the field of real numbers with basis $\{1, i, j, k\}$ [1]. A quaternion q is of the form

$$q = q_0 + q_1i + q_2j + q_3k,$$

where q_0, q_1, q_2, q_3 are real numbers, and i, j, k are quaternionic units satisfy the rules

$$i^2 = j^2 = k^2 = ijk = -1, \quad ij = k = -ji, \quad jk = i = -jk, \quad ki = j = -ik.$$

Horadam [2] defined the n^{th} Fibonacci and Lucas quaternions as

$$Q_n = F_n + F_{n+1}i + F_{n+2}j + F_{n+3}k$$

and

$$V_n = L_n + L_{n+1}i + L_{n+2}j + L_{n+3}k,$$

respectively, where F_n is the n^{th} Fibonacci number defined by $F_n = F_{n-1} + F_{n-2}$ with $F_0 = 0, F_1 = 1$, L_n is the n^{th} Lucas number defined by $L_n = L_{n-1} + L_{n-2}$ with $L_0 = 2, L_1 = 1$, and i, j, k are quaternionic units. In [3], Halıcı defined the complex Fibonacci quaternions and gave some algebraic properties for these quaternions. Moreover, in [4], Halıcı and Cerda-Morales studied the Gaussian Fibonacci quaternions, and in [5], Halıcı introduced the Gaussian Lucas quaternions.

More recently, Akkuş and Kızılaslan [6] introduced the q -Fibonacci and q -Lucas quaternions as

$$Q_n = \alpha^{n-1}[n]_q + \alpha^n[n+1]_qi + \alpha^{n+1}[n+2]_qj + \alpha^{n+2}[n+3]_qk$$

and

$$V_n = \alpha^n \frac{[2n]_q}{[n]_q} + \alpha^{n+1} \frac{[2n+2]_q}{[n+1]_q}i + \alpha^{n+2} \frac{[2n+4]_q}{[n+2]_q}j + \alpha^{n+3} \frac{[2n+6]_q}{[n+3]_q}k,$$

respectively, where i, j, k are quaternionic units. These quaternions are generalized the Fibonacci and Lucas quaternions by receiving components from the q -integers $[n]_q$ defined by $[n]_q = \frac{1-q^n}{1-q}$ (see, [7]). For $\alpha = \frac{1+\sqrt{5}}{2}$ and $q = -\frac{1}{\alpha^2}$, q -Fibonacci and q -Lucas quaternions are reduced the Fibonacci and Lucas quaternions, respectively.

Inspired from the above mentioned studies, in this study, a new generalization for complex (Gaussian) Fibonacci quaternions and complex (Gaussian) Lucas quaternions called complex (Gaussian) q -Fibonacci quaternions and complex (Gaussian) q -Lucas quaternions are defined, respectively, by

$$\mathbb{C}Q_n = Q_n + i Q_{n+1}$$

and

$$\mathbb{C}V_n = V_n + i V_{n+1},$$

where Q_n and V_n are the n^{th} q -Fibonacci and q -Lucas quaternions, respectively.

Furthermore, Binet's formulas, exponential generating functions, Poisson generating functions, Catalan's identities, Cassini's identities and d'Ocagne's identities for complex (Gaussian) q -Fibonacci quaternions and complex (Gaussian) q -Lucas quaternions are given.

Key Words: Quaternion, q -integer, complex Fibonacci number.

REFERENCES

- [1] W.R. Hamilton, Lectures on Quaternions, Hodges and Smith, Dublin, 1853.
- [2] A.F. Horadam, Complex Fibonacci numbers and Fibonacci quaternions, American Math. Monthly 70 (1963), 289-291.
- [3] S. Halıcı, On complex Fibonacci quaternions, Adv. Appl. Clifford Algebras 23 (2013), 105-112.
- [4] S. Halıcı and G. Cerda-Morales, On quaternion-Gaussian Fibonacci numbers and their properties, An. St. Univ. Ovidius. Constanta 29 (2021), 71-82.
- [5] S. Halıcı, On quaternion-Gaussian Lucas numbers, Math. Meth. Appl. Sci. 44 (2021), 7601-7606.
- [6] İ. Akkuş and G. Kızılaslan, Quaternions: Quantum calculus approach with applications, Kuwait J. Sci. 46 (2019), 1-13.

- [7] V. Kac and P. Cheung, Quantum Calculus Universitext, Springer-Verlag, New York, 2002.

Groups Whose Proper Factor Groups Are Fitting

Selami Ercan

Gazi University, Gazi Faculty of Education, Division of Mathematics Education

ABSTRACT

Let G be a group and N be a normal subgroup of G . The factor-group G/N is said to be a proper factor-group if N is non-trivial. The influence of properties of proper factor-groups on properties of groups was the subject of investigation of many authors. Some the results related to groups with many similar proper factor-groups can be found in [2]. Torsion-free groups with every proper homomorphic image an N_1 -group in [1], where N_1 -group if all subgroup of G are subnormal. N_1 -group are certain generalizations of Fitting group. Fitting group is defined to be the subgroup generated by nilpotent normal subgroups of G . Groups, all proper factor-groups of which are Fitting group, are studied in this study. Therefore, in an investigation concerning all proper factor-groups, it is natural to consider groups which include a non-identity normal nilpotent subgroup, that are groups with a non-identity Fitting subgroup.

In this study, we obtained the following results:

Let G be a group which all proper factor-groups of which are Fitting.

- (i) If G is not Fitting group, then Fitting subgroup of G is abelian,
- (ii) If $\text{Fit}(G)$ is torsion-free, then G is torsion free,
- (iii) If for all $x \in G$, $C_G(\langle x \rangle)$ is non-trivial, then G is Fitting.

Key Words: Fitting, homomorphic image, subnormal subgroup

REFERENCES

- [1] S. ERCAN, TORSION-FREE GROUPS WITH EVERY PROPER IMAGES AN M_1 -GROUP, ALGEBRA AND DISCRETE MATHEMATICS, NUMBER 2 (2004), 56-58.
- [2] L. A. KURDACHENKO, J. OTAL, Y. I. SUBOTTIN, GROUPS WITH PRESCRIBED QUOTIENT GROUPS AND ASSOCIATED MODULE THEORY, WORLD SCIENTIFIC, 2002.
- [3] L. A. KURDACHENKO, Y. I. SUBOTTIN, GROUPS WHOSE PROPER QUOTIENTS ARE HYPERCENTRAL, J. AUSTRAL. MATH. SOC.(SERIESA) 65 (1998), 224-237.
- [4] L. A. KURDACHENKO, P SOULES, GROUPS WITH PROPER HYPERCYCLIC HOMOMORPHIC IMAGES, RICERCHE MATH. 50(1) (2001), 53-65
- [5] L. A. KURDACHENKO, P SOULES, GROUPS WITH HYPERCYCLIC PROPER QUOTIENT GROUPS, UKRAIN. MATH. JOURNAL 55 (2003), 566-575.
- [6] D. J. S. ROBINSON, A COURSE IN THE THEORY OF GROUPS, SPRINGER-VERLAG, NEW-YORK, 1982.
- [7] D. J. S. ROBINSON, FINITENESS CONDITIONS AND GENERALIZED SOLUBLE GROUPS, VOL. 1 AND 2, SPRINGER-VERLAG, BERLIN, 1972.

LINEAR MAPS PRESERVING DRAZIN INVERSES OF MATRICES OVER LOCAL RINGS

Tugce Pekacar Calci¹, **Huanyin Chen**², **Sait Halicioglu**³ and **Guo Shile**⁴

¹ Ankara University, Ankara, Turkey, tcalci@ankara.edu.tr

² Hangzhou Normal University, Hangzhou, China, huanyinchen@aliyun.com

³ Ankara University, Ankara, Turkey, halici@ankara.edu.tr

⁴ Fujian Normal University, Fuqing, China, gsl456@163.com

ABSTRACT

Let be R a local ring and suppose that there exists $a \in F^*$ such that $a^6 \neq 1$ and let $T: M_n(R)$ to $M_m(R)$ be a linear map preserving Drazin inverses. Then we prove that $T = 0$; or $n = m$ and T preserves idempotents. We thereby determine the form of linear maps from $M_n(R)$ to $M_m(R)$ preserving Drazin inverses of matrices.

Key Words: Linear map, Drazin inverse, local ring.

REFERENCES

- [1] N. Boudi and M. Mbekhta, Additive maps preserving strongly generalized inverses, J. Operator Theory 64 (2010), 117--130.
- [2] C. Bu, Linear maps preserving Drazin inverses of matrices over fields, Linear Algebra Appl., 396 (2005), 159--173.
- [3] C. Cao, Linear maps preserving idempotence on matrix modules over local rings, J. Natur. Sci. Heilongjiang Univ. 1 (1989), 1--3.
- [4] C. Cao and X. Zhang, Linear preservers between matrix modules over connected commutative rings, Linear Algebra Appl., 397 (2005), 355--366.
- [5] H. Chen, Rings Related Stable Range Conditions, Series in Algebra 11, World Scientific, Hackensack, NJ, 2011.
- [6] J. Cui, Additive Drazin inverse preservers, Linear Algebra Appl., 426 (2007), 448--453.
- [7] M. Dana and R. Yousefi, Formulas for the Drazin inverse of matrices with new conditions and its applications, Int. J. Appl. Comput. Math., 4 (2018), doi: 10.1007/s40819-017-0459-5.
- [8] L. Guo, J. Chen and H. Zou, Representations for the Drazin inverse of the sum of two matrices and its applications, Bulletin of the Iranian Mathematical Society, 2019, DOI:10.1007/s41980-018-0159-x.

***N*-Barely Transitive Groups**

Oğuz Alkış¹, Ahmet Arıkan² and Aynur Arıkan³

1 Science Faculty, Department of Mathematics, Gazi University, 06500 Teknikokullar, Ankara, Turkey, oguzalks@gmail.com

2 Gazi Education Faculty, Department of Mathematics and Science Education, Gazi University, 06500 Teknikokullar, Ankara, Turkey, arikan@gazi.edu.tr

3 Science Faculty, Department of Mathematics, Gazi University, 06500 Teknikokullar, Ankara, Turkey, yalincak@gazi.edu.tr

ABSTRACT

A group is called barely transitive (**BT**-group) or it has barely transitive permutation representation if it acts on an infinite set transitively and faithfully and all its proper subgroups have finite orbits. In [4,5] Hartley introduced **BT**-groups in connection with groups of Heineken-Mohamed type. A description of non-perfect locally finite **BT**-groups was given in [5] (Love's theorem). Some of the main properties of **BT**-groups are described by Kuzucuoğlu in [6].

We restrict the definition of **BT**-groups to normal subgroups and consider the questions “which properties of **BT**-groups are preserved? Is there such a group which is not a **BT**-group?”. We call this new form **N**-barely transitive (**NBT**-group) i.e., the group **G** acts on an infinite set transitively and faithfully, and all its proper normal subgroups have finite orbits. Equivalently, **G** is an **NBT**-group if **G** possesses a subgroup **H** such that $\text{Core}_G H = \bigcap_{g \in G} H^g = 1$ and $|N : N \cap H| < \infty$ for every proper normal subgroup **N** of **G**. Notice that the subgroup **H** is the stabilizer of a point. Clearly every infinite simple group is an **NBT**-group. So, there are **NBT**-groups of any given infinite cardinality.

A group is called an **FC**-group if every element has finitely many conjugates. If a group in which every proper subgroup is an **FC**-group but itself is not, then it is called a minimal non-**FC**-group (**MNFC**-group). The answer of the question “Does there exist a perfect locally finite **MNFC**-**p**-group?” is still unknown. If there exists such a group, then this group has a quotient which is a **BT**-**p**-group of finitary permutations

on some infinite set by [8]. Recall that a permutation group G is finitary if the support of every element is finite. In [5] it is shown that if there exists a finitary BT -group, then this group is a perfect $MNFC$ - p -group where p is a prime. Therefore, the existence of a finitary BT -group will answer the question positively as it is mentioned in [7]. In this presentation, some of the results on NBT -groups in [1,2] are mentioned and it is given a perfect finitary NBT - p -group for any prime p which is not a BT -group. So, the question is answered positively for NBT -groups.

Key Words: Barely transitive group, N -barely transitive group, minimal non- FC -group, finitary permutation group, perfect group.

REFERENCES

- [1] O. Alkış, A. Arıkan, and A. Arıkan, N -barely transitive permutation groups. Ricerche mat (2021). <https://doi.org/10.1007/s11587-021-00608-x>
- [2] O. Alkış and A. Arıkan, Some properties of NBT -groups, Gazi University Journal of Science (2021), submitted.
- [3] B. Hartley, A note on the normalizer condition. Proc. Camb. Phil. Soc. 74 (1973), 11-15.
- [4] B. Hartley, On the normalizer condition and barely transitive permutation groups. Algebra and Logic 13 (1974), 334-340.
- [5] B. Hartley and M. Kuzucuoğlu, Non-simplicity of locally finite barely transitive groups. Proc. Edin. Math. Soc. 40, (1997), 483-490.
- [6] M. Kuzucuoğlu, Barely transitive permutation groups. Arch. Math. 55 (1990), 521-532.
- [7] M. Kuzucuoğlu, Barely transitive groups. Turk J. Math. 31 (2007), 1-15.
- [8] F. Leinen, A reduction theorem for perfect locally finite minimal non- FC groups. Glasgow Math. J. 41(1999), 81-83.

NC-Barely Transitive Groups

Oğuz Alkış¹, Ahmet Arıkan² and Aynur Arıkan³

1 Science Faculty, Department of Mathematics, Gazi University, 06500 Teknikokullar, Ankara, Turkey, oguzalks@gmail.com

2 Gazi Education Faculty, Department of Mathematics and Science Education, Gazi University, 06500 Teknikokullar, Ankara, Turkey, arikan@gazi.edu.tr

3 Science Faculty, Department of Mathematics, Gazi University, 06500 Teknikokullar, Ankara, Turkey, yalincak@gazi.edu.tr

ABSTRACT

Let G be a group acting transitively and faithfully on an infinite set. If for every proper normal subgroup N and element x in G with $\langle N, x \rangle \neq G$, every orbit of $\langle N, x \rangle$ is finite, then G will be called an NC -barely transitive group ($NCBT$ -group). Then G has a subgroup H of infinite index such that $\text{Core}_G H = 1$ and $|\langle N, x \rangle : \langle N, x \rangle \cap H| < \infty$ for every proper normal subgroup N and x in G with $\langle N, x \rangle \neq G$. Recall that an infinite group is said to be an N -barely transitive group (NBT -group) if it acts transitively and faithfully on an infinite set and every orbit of every proper normal subgroup is finite [1,2]. So, every $NCBT$ -group is an NBT -group. As NBT -groups, the concept of $NCBT$ -group is a generalization of barely transitive groups (BT -groups) in which every proper subgroup has finite orbits (see [3,4] for locally finite BT -groups).

In this presentation, we consider NBT -groups which are $NCBT$ -groups in [1,2]. We give a characterization of NBT -groups with every non-trivial epimorphic image is non-simple. Also, the structure of non-perfect NBT -groups and some results under the restriction of being locally soluble (or locally nilpotent) are given. Furthermore, we provide perfect $NCBT$ -group examples and soluble $NCBT$ -groups examples of derived length 2 in the context of the groups satisfying minimal condition on normal subgroups. Indeed, it is shown that it is possible to construct soluble $NCBT$ -groups of arbitrary derived length [1].

Key Words: Barely transitive group, N -barely transitive group, NC -barely transitive group, finitary permutation group, perfect group, locally soluble group.

REFERENCES

- [1] O. Alkış, A. Arıkan and A. Arıkan, *N*-barely transitive permutation groups. *Ricerche mat* (2021). <https://doi.org/10.1007/s11587-021-00608-x>
- [2] O. Alkış and A. Arıkan, Some properties of *NBT*-groups, *Gazi University Journal of Science* (2021), submitted.
- [3] B. Hartley, On the normalizer condition and barely transitive permutation groups. *Algebra and Logic* 13(1974), 334-340.
- [4] M. Kuzucuoğlu, Barely transitive permutation groups. *Arch. Math.* 55 (1990), 521-532.

On The Catalan Transforms of The Incomplete Jacobsthal Numbers

Engin Özkan¹, Mine Uysal² and Bahar Kuloğlu³

1 Department of Mathematics, Faculty of Arts and Sciences, Erzincan Binali Yıldırım University, Erzincan, 24100, Turkey, ozkan@erzincan.edu.tr

2 Graduate School of Natural and Applied Sciences, Erzincan Binali Yıldırım University, Erzincan, 24100, Turkey, mine.uysal@erzincan.edu.tr

3 Graduate School of Natural and Applied Sciences, Erzincan Binali Yıldırım University, Erzincan, 24100, Turkey, bahar_kuloglu@hotmail.com

ABSTRACT

Number sequences have attracted the attention of many scientists for many years, as they find application in nature and in many sciences [2,4,5,6,7,8]. Number sequences, especially Fibonacci sequences, find application in many departments of mathematics as well as in other branches of science [2,3,4,8].

One of these generalizations is the Jacobsthal numbers. Jacobsthal numbers were first defined by Horodam in 1996 as follows.

For $n \geq 2$, the Jacobsthal numbers J_n are defined by the following recurrence relation

$$J_n = J_{n-1} + 2J_{n-2}$$

with $J_0 = 0$ and $J_1 = 1$.

In the same paper, he gave an explicit form of Jacobsthal numbers as follows

$$J_n = \sum_{r=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n-1-r}{r} 2^r$$

where $\lfloor x \rfloor = \max\{m \in \mathbb{Z} : m \leq x\}$.

The Catalan numbers with general term $C(n)$ are defined in [1] by

$$C(n) = \frac{1}{n+1} \binom{2n}{n}.$$

Generating functions of the Catalan numbers $C(n)$ are given in [1] by

$$c(x) = \frac{1 - \sqrt{1-4x}}{2x}.$$

Philip defined incomplete Fibonacci and Lucas numbers in 1996 [4]. Djordjevic *et al.* introduced incomplete generalized Jacobsthal and Jacobsthal-Lucas numbers in 2005 [2].

In this work, we give the Catalan transform of the incomplete Jacobsthal numbers and find the generating functions of the Catalan transform of these numbers. Also, we apply the Hankel transform to the Catalan transforms of these numbers. We calculate determinants of matrixes formed with $CJ_{k,n}^l$ by using Hankel transform.

Key Words: Binet formula, Generating function, Hankel transform, Catalan transform, Incomplete Jacobsthal numbers.

This work was supported by Erzincan Binali Yıldırım University Scientific Research Projects Coordination Unit. Project Number: FBA-2021-783.

REFERENCES

- [1] P. Barry and A Catalan transform and related transformations on integer sequences, J. Integer Seq. 8(4) (2005), 1-24.
- [2] G. B. Djordjević and H. M. Srivastava, Incomplete generalized Jacobsthal and Jacobsthal-Lucas numbers, Mathematical and Computer Modelling 42(9-10) (2005), 1049-1056.
- [3] S. Falcon, Catalan Transform of the k-Fibonacci sequence, Communications of the Korean Mathematical Society 28(4) (2013), 827–832.
- [4] P. Filipponi, Incomplete Fibonacci and Lucas numbers, Rendiconti del Circolo Matematico di Palermo 45(1) (1996), 37-56.
- [5] Á. Pintér and H. M. Srivastava, Generating functions of the incomplete Fibonacci and Lucas numbers, Rend. Circ. Mat. Palermo 48(2) (1999), 591–596.
- [6] P. M. Rajković, M. D. Petković and P. Barry, The Hankel transform of the sum of consecutive generalized Catalan numbers, Integral Transform Spec. Funt. 18(4) (2007), 285-296.
- [7] M. Taştan and E. Özkan, Catalan Transform of the k-Jacobsthal Sequence, Electronic Journal of Mathematical Analysis and Applications (EJMAA) 8(2) (2020), 70-74.
- [8] M. Taştan and E. Özkan, Catalan transform of the k-Pell, k-Pell–Lucas and modified k-Pell sequence, Notes on Number Theory and Discrete Mathematics 27 (1) (2021), 198-207.

On the Finitary Permutation Groups

Rümeysa S. Altınkaya¹, Aynur Arıkan²

¹ Gazi University, Mathematics Department, Ankara, Turkey, email: rsgolcu@gazi.edu.tr

² Gazi University, Mathematics Department, Ankara, Turkey, email: yalincak@gazi.edu.tr

ABSTRACT

Let G be a group acting on infinity set Ω . The support of x in G is defined by $\text{supp}(x) = \{\alpha \in \Omega \mid \alpha^x \neq \alpha\}$. If $\text{supp}(x)$ is finite for all $x \in G$, then G is called finitary symmetric group on Ω . If there exist an equivalence relation ϱ on Ω such that $\alpha \equiv \beta \pmod{\varrho}$ if and only if $\alpha^x \equiv \beta^x \pmod{\varrho}$ for all $x \in G$, then ϱ is a congruence relation. G is almost primitive on Ω if there is maximal congruence such that $|\varrho|$ is finite; if there is no maximal congruence such that $|\varrho|$ is finite, then G is totally imprimitive. According to the result of Giorgetta [1], if such a group G has finite exponent then all its orbits must be finite. As for the result of Wiegold [3], if such a group G is soluble then all its orbits must be finite.

The main purpose of this study is to generalize these two studies mentioned. It is investigated that all orbits of the group G in Ω are finite for almost primitive and totally imprimitive cases of the group G of finite permutations of a set Ω satisfying some non-trivial laws. For this research, the group G is shown to generates variety \mathfrak{D} of all groups. Also, it is examined that all orbits are finite if the group G is locally solvable group.

Key Words: Finitary permutation groups, almost primitive, totally imprimitive.

REFERENCES

- [1] P. M. Neumann, The lawlessness of groups of finitary permutations, Arch. Math. 26 (1975), 561-566.
- [2] D.Giorgetta, Dr. Phil. Dissertation, E.T.H. Zürich.
- [3] J. Wiegold, Groups of finitary permutations. Arch.Math. 25 (1974), 466-469.
- [4] D.J.S. Robinson, A Course in the Theory of Groups, Springer-Verlag, New York, 1996.

SIP-direct-injective Modules and Rings

Özgür Taşdemir¹

1 Trakya University, Edirne, Turkey, ozgurtasdemir@trakya.edu.tr

ABSTRACT

Recall that a module M over a ring is called a *direct-injective module* (or *C2-module*) if every submodule of M is isomorphic to a direct summand of M , then it is a direct summand of M . A module M is called an *SIP-module* if the intersection of every pair of direct summands of M is again a direct summand of M . In this study, we define that a module M over a ring is called an *SIP-direct-injective module* if the intersection of every pair of direct summands of M is isomorphic to a direct summand of M , then it is a direct summand of M . The class of SIP-direct-injective modules unifies SIP-modules and C2-modules. Various basic properties of these modules are studied.

Key Words: summand intersection property, C2-modules, direct-injective-modules, direct summand, isomorphism.

REFERENCES

- [1] G.V. Wilson, On modules with the summand intersection property, Communications Algebra, 14(1) (1986),21-38.
- [2] S.H. Mohamed, B.J. Müller, Continuous and Discrete Modules, Cambridge Univ. Press, Cambridge, UK, 1990.

Some Notes on Soft Modules in Soft Sets

Filiz Çıtak

Tokat Gaziosmanpaşa University, Faculty of Art and Science, Department of Mathematics,
Tokat, Turkey, filiz.citak@gop.edu.tr

ABSTRACT

There are several theories, such as the probability theory, the theory of fuzzy sets, the theory of rough sets, the theory of vague sets, the theory of interval mathematics, which can be considered as mathematical tools for dealing with uncertainties. The concept of soft sets was introduced by Molodtsov [1] as a new mathematical tool for dealing with uncertainties in 1999. Maji [2] defined operations on soft sets. According to Maji's definition,

Let A be a set of parameters. Let U be an initial universal set and let $P(U)$ be the power set of U . A function F_A is called as a soft set over U , where $F_A: A \rightarrow P(U)$ is a function.

The soft set theory has been developed in many fields as soft decision making, soft fuzzy sets, intuitionistic soft set. Next, many researchers have studied algebraic structures over a soft set.

In this study, the operations of soft sets over a module are investigated. Sum of two soft sets, negative of a soft set and multiplied by a scalar of a soft set over a module are defined. Their properties were examined. Then, a soft module is defined as follows.

A soft module F_M is a soft set F_M over U such that

- 1) $F_M(0_R) = U$
- 2) $F_M(r.a) \supseteq F_M(a)$ for all $r \in R$ and $a \in M$.
- 3) $F_M(a+b) \supseteq F_M(a) \cap F_M(b)$ for all $a, b \in M$.

It is shown that multiplied by a scalar of the soft module is a subset of the soft module. Also, it is proved that sum of the soft module is a soft module. And some basic properties of soft modules are discussed.

Key Words: Soft sets, soft product, soft intersection modules.

REFERENCES

- [1] D. Molodtsov, Soft set theory-first results, Computers and Mathematics with Applications 37 (1999), 19-31.
- [2] P.K. Maji, R. Bismas and A.R. Roy, Soft set theory, Computers and Mathematics with Applications 45 (2003), 555-562.

Some properties of Apostol Bernoulli Fibonacci and Apostol Euler Fibonacci Polynomials

Naim TUĞLU¹, Elif ERCAN²

¹ Gazi University, Faculty of Science, Departments of Mathematics Ankara / TURKEY
,naimtuglu@gazi.edu.tr

² Gazi University, Faculty of Science, Departments of Mathematics Ankara /
TURKEY, elifercan06@gmail.com

ABSTRACT

The generating function for Bernoulli polynomials is defined by Taylor series expansion,

$$\frac{te^{xt}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}$$

where $B_n(x)$ are the Bernoulli polynomials in x , for all $n > 0$.

The Apostol-Bernoulli polynomials are defined by the exponential generating function,

$$\left(\frac{t}{\lambda e^t - 1}\right) e^{xt} = \sum_{n=0}^{\infty} B_n(x, \lambda) \frac{t^n}{n!}$$

A famous and important sequence is the Fibonacci sequence, named after the the Italian mathematician known as Leonardo Pisano. This sequence is defined recursively. The Fibonacci sequence is defined by $F_{n+2} = F_{n+1} + F_n$, when $F_0 = 0, F_1 = 1$

The Golden exponential function is entire analytic function, defined as

$$e_F^{tx} = \sum_{n=0}^{\infty} \frac{t^n}{F_n!}$$

Generating function for Bernoulli-Fibonacci polynomials $B_n^F(x)$ is defined by series expansion,

$$\frac{te_F^{tx}}{e_F^t - 1} = \sum_{n=0}^{\infty} B_n^F(x) \frac{t^n}{F_n!}$$

The generating function for Euler polynomials is defined by Taylor series expansion,

$$\frac{2e^{xt}}{e^t + 1} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}$$

The Apostol-Euler polynomials are defined by the exponential generating function,

$$\left(\frac{2}{\lambda e^t + 1}\right) e^{xt} = \sum_{n=0}^{\infty} E_n(x, \lambda) \frac{t^n}{n!}$$

Generating function for Euler-Fibonacci polynomials $B_n^F(x)$ is defined by series expansion,

$$\frac{2e_F^{tx}}{e_F^t + 1} = \sum_{n=0}^{\infty} E_n^F(x) \frac{t^n}{F_n!}$$

The Golden derivative, applied to Bernoulli-Fibonacci polynomials $B_n^F(x)$ gives Fibonacci numbers, $D_F^x(B_n^F(x)) = F_n B_n^F(x)$

In this study a new F-exponential generating function for Apostol Bernoulli Fibonacci polynomials and various properties of Apostol Bernoulli Fibonacci polynomials are obtained. By identifying Apostol Euler Fibonacci numbers and polynomials are found. In addition, harmonic based F exponential generating function is defined for Apostol Bernoulli Fibonacci numbers and Apostol Euler Fibonacci numbers. We define incomplete Apostol Bernoulli Fibonacci numbers and incomplete Apostol Euler Fibonacci numbers by using Apostol Bernoulli Fibonacci numbers and Apostol Euler Fibonacci numbers.

Key Words: Bernoulli Fibonacci polynomials, Euler Fibonacci polynomials, Apostol polynomials.

REFERENCES

- [1] Q.M.Luo, On the Apostol-Bernoulli polynomials. *Cent. Eur. J. Math.* 2(4) (2004), 509-515.
- [2] Q.M. Luo and H.M. Srivastava, Some generalizations of the Apostol Bernoulli and Apostol-Euler polynomials. *J. Math. Anal. Appl.* 308, no. 1, (2005), 290-302.
- [3] Q.M. Luo and H.M. Srivastava, Some relationships between the Apostol-Bernoulli and Apostol-Euler polynomials. *Comput. Math. Appl.* 51(3-4), (2006), 631-642.
- [4] H.M. Srivastava ve Pinter, Remarks on some relationships between the Bernoulli and Euler polynomials. *Appl. Math. Lett.* 17(4) (2004), 375-380.
- [5] S. Kuş, N. Tuglu ve T. Kim, Bernoulli F-polynomials and Fibo-Bernoulli matrices. *Advances in Difference Equations*, 2019(145).
- [6] M. Özvatan, Generalized Golden-Fibonacci Calculus and Applications, Master Thesis, The Graduate School of Engineering and Sciences of Izmir Institute of Technology, İzmir, 2018.
- [7] T.M. Apostol, On the Lerch Zeta function, *Pacific J. Math.* 1 (1951) 161–167.
- [8] T.M. Apostol, Introduction to Analytic Number Theory, Springer-Verlag, New York, 1976

The Universal Difference Property for Generalized Splines over Principal Ideal Domains

Samet Sarioğlu¹ and Selma Altınok²

¹ Hacettepe University Department of Mathematics, Ankara, Turkey,
ssarioglan@hacettepe.edu.tr

² Hacettepe University Department of Mathematics, Ankara, Turkey,
sbhupal@hacettepe.edu.tr

ABSTRACT

We study generalized splines over principal ideal domains and investigate the universal difference property (UDP). The UDP is introduced by K. Anders and her students [3]. They showed that the UDP holds for cycles, paths and trees. They also proved that the UDP holds for theta graphs over Prüfer domains and uniserial rings.

We prove that the UDP holds for any graph over principal ideal domain by using the existence of flow-up classes over PID's.

Key Words: Generalized splines, universal difference property, principal ideal domains.

REFERENCES

- [1] S. Altınok and S. Sarioğlu, Flow-up bases for generalized spline modules on arbitrary graphs, *Journal of Algebra and its Applications* (2021), doi: 10.1142/S0219498821501802.
- [2] K. Anders, A. Crans, B. Foster-Greenwood, B. Mellor and J. Tymoczko, Graphs admitting only constant splines, *Pacific J. Math.* 304 (2020), no. 2, 385-400.
- [3] K. Anders, D. Arreola, L. Asencio, C. Ireland, L. Smith, Graph splines and the universal difference property, *Joint Mathematics Meeting AMS Special Session* (2020).
- [4] S. Gilbert, J. Tymoczko and S. Viel, Generalized splines on arbitrary graphs, *Pacific J. Math.* 281 (2016), no. 2, 333-364.
- [5] J. Tymoczko, An introduction to equivariant cohomology and homology, following Goresky, Kottwitz, and MacPherson, *Snowbird lectures in algebraic geometry*, *Contemp. Math.* (2005), vol. 388, Amer. Math. Soc., Providence, RI, pp. 169-188.

On semi-Nekrasov matrices

Maja Nedović¹

¹ Department for Fundamental Sciences,
Faculty of Technical Sciences,
University of Novi Sad, Serbia
maja.nedovic@uns.ac.rs

ABSTRACT

Nekrasov matrices represent a special subclass in the class of non-singular H-matrices. In this paper, lower semi-Nekrasov matrix class, a generalization of Nekrasov matrix class, is considered. The relation of semi-Nekrasov matrices to semi-strictly diagonally dominant matrices is established. For matrices of this type, an upper bound for the maximum norm of the inverse matrix is proposed. New bound can be applied to Nekrasov matrices as well and numerical examples show that, in some cases, new norm bound for the inverse matrix is tighter than some already known bounds defined for Nekrasov matrices. The corresponding class of block matrices is also considered, as the subclass in the class of block-H matrices. Two different types of block-generalizations are given. Starting from the original matrix and the fixed partition of the index set, we observe two different ways to define the comparison matrix that is of a smaller dimension and using its entries we define upper bounds for the norm of the inverse matrix in the block case as well. As the norm bounds for the inverse can be used in bounding the condition number or in estimating error bounds in linear complementarity problems, we discussed possibilities for applications of the presented results.

Key Words: Nekrasov matrices, maximum norm bound, block matrices.

REFERENCES

- [1] R. Beauwens, Semistrict diagonal dominance, SIAM J. Numer. Anal. 13 (1976) 109–112.
- [2] Lj. Cvetković, K. Doroslovački, Max norm estimation for the inverse of block matrices, Appl. Math. Comput. 242 (2014) 694–706.

- [3] L.Yu. Kolotilina, On bounding inverse to Nekrasov matrices in the infinity norm, Zap.Nauchn.Sem.POMI. 419 (2013) 111–120.
- [4] W. Li, On Nekrasov matrices, Linear Algebra Appl. 281 (1998) 87–96.
- [5] M. Nedović, Norm bounds for the inverse for generalized Nekrasov matrices in point-wise and block case, Filomat Vol.35 No 8 (2021)
- [6] M. Nedović, Lj. Cvetković, Norm bounds for the inverse and error bounds for linear complementarity problems for $\{P1, P2\}$ -Nekrasov matrices, Filomat Vol 35 No 1 (2021)
- [7] P.N. Shivakumar et al., On two-sided bounds related to weakly diagonally dominant M-matrices with application to digital dynamics, SIAM J. Matrix Anal. Appl. 17 (2) (1996) 298–312.
- [8] T. Szulc, Lj. Cvetković, M. Nedović, Scaling technique for Partition - Nekrasov matrices, Appl.Math.Comput. Vol 271C (2015) 201–208.

ANALYSIS

An Extension of the Adams-type Theorem to Vanishing Generalized Weighted Morrey Spaces

Abdulhamit Kucukaslan

Faculty of Applied Sciences, Pamukkale University, 20680, Denizli, Turkey
kucukaslan@pau.edu.tr

ABSTRACT

Morrey spaces were introduced by Morrey in [3]. These spaces appeared to be useful in the study of local behavior properties of the solutions of second order elliptic PDEs. The vanishing Morrey space of the classical Morrey spaces was introduced by Vitanza in [7] and applied there to obtain a regularity result for elliptic PDEs. Later in [8] Vitanza proved an existence theorem for a Dirichlet problem and a regularity result assuming that the partial derivatives of the coefficients of the highest and lower order terms belong to vanishing Morrey spaces depending on the dimension. Persson et al. [4] showed the commutators of Hardy operators on vanishing Morrey spaces. Also Ragusa [5] obtained a sufficient condition for commutators of fractional integral operators to belong to vanishing Morrey spaces.

The vanishing generalized Morrey space and vanishing generalized local Morrey space was introduced by Samko in [6]. The boundedness of the multi-dimensional Hardy type operators, maximal, potential and singular operators in these spaces were proved in [6]. Kucukaslan et al. [1] proved the Spanne-type and Adams-type boundedness of generalized fractional integral operators on vanishing generalized local Morrey spaces. Guliyev et al. [2] proved the commutators of Riesz potential operator in the vanishing generalized weighted Morrey spaces with variable exponent.

The generalized fractional maximal operators M_p was initially investigated in by Nakai in 1994. Nakai introduced the generalized Morrey spaces and proved the boundedness of the generalized fractional integral operator in these spaces. Nowadays many authors have been culminating important observations about the generalized fractional maximal operators M_p especially in connection with Morrey-type spaces.

In this talk, we generalize the Adams-type boundedness of the generalized fractional maximal operator M_ρ from the vanishing generalized weighted Morrey spaces $VM_{p,\varphi^p}(\omega)$ to the vanishing generalized weighted Morrey spaces $VM_{q,\varphi^q}(\omega)$ with the weight function ω , and from the vanishing generalized weighted Morrey spaces $VM_{1,\varphi}(\omega)$ to the vanishing generalized weighted weak Morrey spaces $VWM_{q,\varphi^q}(\omega)$. The all weight functions belong to the Muckenhoupt-Weeden classes $A_{p,q}$.

Key Words: Generalized fractional maximal operator, Vanishing generalized weighted Morrey space, Muckenhoupt-Weeden classes.

REFERENCES

- [1] A. Kucukaslan, S. G. Hasanov and C. Aykol, Generalized fractional integral operators on vanishing generalized local Morrey spaces, *Int. J. of Math. Anal.*, 11(6) (2017), 277--291.
- [2] V.S. Guliyev, J.J. Hasanov and X.A. Badalov, Commutators of Riesz potential in the vanishing generalized weighted Morrey spaces with variable exponent, *Math. Inequal. Appl.*, 22(1) (2019), 331-351.
- [3] C. B. Morrey, On the solutions of quasi-linear elliptic partial differential equations, *Trans. Amer. Math. Soc.*, 43 (1938), 126-166.
- [4] L. E. Persson, M. A. Ragusa, N. Samko, P. Wall, Commutators of Hardy operators in vanishing Morrey spaces, *AIP Conf. Proc.* 1493, 859, 2012.
- [5] M. A. Ragusa, Commutators of fractional integral operators on vanishing-Morrey spaces, *J. Global Optim.* 40(1-3) (2008), 361-368.
- [6] N. Samko, Weighted Hardy operators in the local generalized vanishing Morrey spaces, *Positivity* 17 (2013), 683-706.
- [7] C. Vitanza, Functions with vanishing Morrey norm and elliptic partial differential equations, *Proceedings of Methods of Real Analysis and Partial Differential Equations*, Capri, Springer (1990), 147-150.
- [8] C. Vitanza, Regularity results for a class of elliptic equations with coefficients in Morrey spaces, *Ricerche di Matematica* 42(2) (1993), 265-281.

Approximation by Some Kantorovich Type Max-Product Operators

Sule Yüksel Güngör¹

1 Gazi University, Faculty of Sciences, Department of Mathematics, 06500, Ankara, Turkey
sulegungor@gazi.edu.tr

ABSTRACT

In order to draw attention to a possible interesting new direction in the constructive approximation theory, an open problem was presented by S.G. Gal in [1]. According to this problem a sequence of discrete max-product approximation operators $L_n^{(M)}$ attached to $f: I \subseteq \mathbb{R} \rightarrow \mathbb{R}^+$ with the operations “V” (maximum) and “.” (product), can be defined by

$$L_n^{(M)}(f)(x) = \frac{\bigvee_{k \in I_n} p_{n,k}(x) \cdot f(x_{n,k})}{\bigvee_{k \in I_n} p_{n,k}(x)}, x \in I, n \in \mathbb{N}$$

where $p_{n,k}(x)$ are various kinds of function basis on I , I_n are finite or infinite families of indices and $\{x_{n,k}: k \in I_n\}$ represents a division of I . These operators are nonlinear positive operators having pseudo-linearity property. Moreover, the max-product operators are subadditive, positive homogenous and monotone. In [2], to each max-product operator $L_n^{(M)}$, the authors defined its Kantorovich variant by

$$LK_n^{(M)}(f)(x) = \frac{\bigvee_{k \in I_n} p_{n,k}(x) \cdot \left(\frac{1}{(x_{n,k+1} - x_{n,k})} \int_{x_{n,k}}^{x_{n,k+1}} f(t) dt \right)}{\bigvee_{k \in I_n} p_{n,k}(x)}, x \in I, n \in \mathbb{N}$$

with $\{x_{n,k}: k \in I_n\}$ a division of the finite or infinite interval I . They studied uniform and pointwise approximation properties, shape preserving properties and localization results for certain well-known operators.

Based on this study we introduce Kantorovich variants of some max-product operators and give quantitative estimates with the help of the quantitative estimates of the original operators. Compared with the approximation order by the original operators, it follows that the approximation order by its Kantorovich variants is no worse. Also, we investigate shape preserving properties for Kantorovich type operators.

Key Words: Max-product operators, max-product operators of Kantorovich type, shape preserving properties.

REFERENCES

- [1] S.G. Gal, Shape-Preserving Approximation by Real and Complex Polynomials, Birkhäuser Boston, Inc., Boston, MA, 2008.
- [2] L. Coroianu and S.G. Gal, Approximation by max-product operators of Kantorovich type, Stud. Univ. Babeş-Bolyai Math. 64 (2019), 207-223.
- [3] Ş.Y. Güngör and N. İspir, Approximation by Bernstein-Chlodowsky operators of max-product kind, Mathematical Communications, 23 (2018), 205-225.
- [4] Ş.Y. Güngör and N. İspir, Quantitative estimates for generalized Szász operators of max-product kind, Results. Math. 70 (2016), 447-456.

APPROXIMATION PROPERTIES OF GENERALIZED JAIN OPERATORS OF TWO VARIABLES

Gürhan SOYLU

Gazi University Graduate School Of Natural And Applied Sciences

grhnsyl@gmail.com

ABSTRACT

In this study, The deals with the approximation processes of the bivariate Jain operators defined in this paper. We also obtain some rates of not only classical uniform convergence a Vorovskaya type result is also proved for these operators.

In the first chapter, the approximation properties of Jain operators are studied. In the second chapter, the continuity of Jain operators and the approximation speed using functions in the Lipschitz class are given. In the third chapter, the approximation properties of bivariate generalized Jain operators are studied. In the fourth chapter, the approximation speed using continuity of bivariate generalized Jain operators and functions in the Lipschitz class is given. In the fifth chapter, the Voronovskaya Asymptotic Approach of Jain Operators is presented. Finally, in the sixth chapter, the Voronovskaya asymptotic approach of bivariate generalized Jain operators is obtained.

Key Words: Jain operators of the variables, linear positive operators, Lipschitz class, modulus of continuity.

REFERENCES

- [1] O. Doğru, R. N. Mohapatra and M. Örkücü, Approximation Properties of Generalized Jain Operators, Filomat, 30:9 (2016), 2359-2366.
- [2] P. Patel and V. N. Mishra, On Generalized Szász-Mirakyan operators, arxiv preprint arxiv: 1508.07896.

- [3] C. Muraru, O. Doğru and A. Gülsün, A-Statistical L_p approximation properties of an integral variant of a general positive linear operators, Tusi Mathematical Research Group, 11:761 (2020), 761-779.
- [4] O. Agratini, Approximation properties of a class of linear operators, Math. Method. Appl. Sci., 36 (2013), 2353-2358.
- [5] F. Dirik and K. Demirci, Szasz-Mirakjan type operators of two variables providing a better estimation on $[0,1] \times [0,1]$, Mat. Vesnik, 63 (2011), 59-66.
- [6] G.C. Jain, Approximation of functions by a new class of linear operators, J. Aust. Math. Soc. 13 (1972), 271-276.
- [7] V. Gupta and G.C. Greubel, Moment estimations of new Szász. Mirakyan. Durrmeyer operators, Applied Mathematics and Computation 271 (2015), 540-547.
- [8] Volkov V I, On the convergence of sequences of linear positive operators in the space of Continuous functions of two variables, (Russian) Dokl. Akad. Nauk SSSR (N.S.) 115 (1957), 17-19.

Generalized Fractional Integral Inequalities on Convex Functions

Abdullah Akkurt¹, Fatih Korkmaz² and Hüseyin Yıldırım³

^{1,2,3}Department of Mathematics, Faculty of Science and Arts, Kahramanmaraş Sütçü İmam University, 46100, Kahramanmaraş, Turkey
abdullahmat@gmail.com, faith.efk@gmail.com, hyildir@ksu.edu.tr

ABSTRACT

The Fractional calculus technique is an important subject that has been known for approximately 300 years and is expected to lead to beneficial results. We can understand the importance of fractional derivatives and integrals from the various branches of science to which they have been applied and the useful results that have emerged. Although there are many definitions of the fractional calculus technique, the most well-known form is Riemann-Liouville derivatives and integrals. For this, please see the following references [6, 9].

The Hermite-Hadamard inequality is indispensable for the theory of inequality, especially for convex functions. The number of Hermite-Hadamard type inequalities for fractional derivatives and integrals has increased considerably in recent years. For this, please see the following references [1-5, 7, 8, 10, 11].

In this paper, we have obtained some new fractional integral inequalities for two synchronous functions, where generalized Riemann-Liouville fractional integrals are used. In obtaining these fractional integral inequalities, we extended them with some new parameters. We show that these inequalities we found support the results available in the literature for the specific values.

Key Words: Fractional Integral, Convex Functions, Riemann-Liouville Fractional Integral.

REFERENCES

- [1] A. AKKURT and H. YILDIRIM, On Some Fractional Integral Inequalities of Hermite-Hadamard type for r -Preinvex Functions, Khayyam J. Math. 2(2) (2016), 120–127, DOI: 10.22034/kjm.2016.40640.
- [2] M.Z. SARIKAYA, A. AKKURT, H. BUDAK, M.E. YILDIRIM and H. YILDIRIM. Hermite-Hadamard's inequalities for conformable fractional integrals. An International Journal of Optimization and Control: Theories & Applications (IJOCTA), 9(1) (2019), 49–59.

- [3] S. BELARBI and Z. DAHMANI, On some new fractional integral inequalities, *Journal of Inequalities in Pure and Applied Mathematics*, 10(3) (2009), Article 86, 5 pp.
- [4] P. L. BUTZER, A. A. KILBAS and J.J. TRUJILLO, Compositions of Hadamard-type fractional integration operators and the semi-group property, *Journal of Mathematical Analysis and Applications*, 269 (2002), 387-400.
- [5] P.L. CHEBYSHEV, Sur les expressions approximatives des integrales definies par les autres prises entre les mêmes limites, *Proc. Math. Soc. Charkov*, 2 (1882), 93–98.
- [6] U.N. KATUGAMPOLA, New Approach to a generalized fractional integral, *Appl. Math. Comput.* 218(3) (2011), 860-865.
- [7] S.M. MALAMUD, Some complements to the Jenson and Chebyshev inequalities and a problem of W. Walter, *Proc. Amer. Math. Soc.*, 129(9) (2001), 2671-2678.
- [8] B.G. PACHPATTE, A note on Chebyshev-Grüss type inequalities for differential functions, *Tamsui Oxford Journal of Mathematical Sciences*, 22(1) (2006), 29-36.
- [9] S.G. SAMKO, A.A. KILBAS and O.I. MARICHEV, *Fractional Integrals and Derivatives, Theory and Applications*, Gordon and Breach, Yverdon et alibi, 1993.
- [10] M.Z. SARIKAYA, E. SET, H. YALDIZ, and N. BAŞAK, Hermite-Hadamard's inequalities for fractional integrals and related fractional inequalities, *Mathematical and Computer Modelling*, Volume 57, Issues 9–10, May 2013, Pages 2403-2407.
- [11] M.Z. SARIKAYA and H. OGUNMEZ, On new inequalities via Riemann-Liouville Fractional Integration, *Abstract and Applied Analysis*, vol. 2012, Article ID 428983, 10 pages, 2012.

Matrix Transformations and Some Algebraic and Topological Properties Related to the New Catalan Double Sequence Space

Sezer Erdem¹ and Serkan Demiriz²

1 Battalgazi Farabi Anatolian Imam Hatip High School, Malatya, Turkey
sezererdem8344@gmail.com

2 Department of Mathematics, Gaziosmanpasa University, Tokat, Turkey
serkandemiriz@gmail.com

ABSTRACT

In mathematics, one of the fascinating number sequence is the integer sequence $1, 1, 2, 5, 14, 42, 132, 429, \dots$ of Catalan numbers. More recently, a new conservative matrix have been defined by means of the fascinating sequence of Catalan numbers and have been studied the matrix domains of this newly introduced matrix in the classical sequence spaces in [5]. After that, this matrix and its domains have been studied in the papers [1,6]. Each story on single sequence and single sequence space has been experienced over double sequence and double sequence space. In this present paper, firstly we define a new double sequence space by using the 4 dimensional Catalan matrix derived by the sequences of the Catalan numbers and we prove that this newly described double sequence space is a Banach space with its norm. Then, we give a inclusion relation including this newly defined double sequence space. Moreover, we calculate the α -, β (bp) - and γ -duals of this space. Finally, we characterize some new 4 dimensional matrix transformation classes and we complete this work with some significant results.

Key Words: 4 dimensional Catalan matrix, double sequence space, α -, β (bp) - and γ -duals, matrix transformation.

REFERENCES

- [1] P. Z. Alp, A new paranormed sequence space defined by Catalan conservative matrix, *Math Meth Appl Sci.* (2020);1-8.
- [2] S. Demiriz and S. Erdem, Domain of Binomial Matrix in Some Spaces of Double Sequences, *Punjab Univ. J. Math.*, 52(11) (2020), 65-79.

- [3] S. Erdem and S. Demiriz, Almost Convergence and 4-Dimensional Binomial Matrix, Konuralp J. Math., 8(2) (2020), 329-336.
- [4] S. Erdem and S. Demiriz, A New RH-Regular Matrix Derived by Jordan's Function and Its Domains on Some Double Sequence Spaces, Journal of Function Spaces, vol. 2021, Article ID 5594751, (2021) 9 pages.
- [5] M. İlkhani, A new conservative matrix derived by Catalan numbers and its matrix domain in the spaces c and c_0 , Linear and Multilinear Algebra, 68:2, (2020), 417-434.
- [6] M. İ. Kara and E. E. Kara, Matrix transformations and compact operators on Catalan sequence spaces, J. Math. Anal. Appl. 498 (2021).
- [7] O. Tuğ, Four-dimensional generalized difference matrix and some double sequence spaces. J. Inequal. Appl., 2017(1), 149 (2017).
- [8] M. Yeşilkayağıl and F. Başar, Domain of Euler mean in the space of absolutely p -summable double sequences with $0 < p < \infty$, Anal. Theory Appl., **34**, (2018), No:3, 241-252.

On Absolute Fibonacci Series Space And Matrix Operators

Fadime Gökçe¹

¹ University of Pamukkale, Department of Statistics, fgokce@pau.edu.tr

ABSTRACT

In a more recent paper, the series space $|F_\theta|(p)$ which is defined as the domain of a matrix corresponding to the absolute Fibonacci summability in the Maddox's space $l(p)$ has been introduced and studied by Gökçe and Sarıgöl [1]. In this study, certain characterizations of matrix operators from the paranormed space $|F_\theta|(p)$ to the classical sequence spaces c, c_0, l_∞ are obtained. Also, we show that the matrix operators between the absolute Fibonacci series space and the spaces c, c_0, l_∞ are bounded operators.

Key Words: Absolute summability, Fibonacci numbers, matrix transformation, bounded linear operators.

REFERENCES

- [1] F. Gökçe and M.A. Sarıgöl, Series spaces derived from absolute Fibonacci summability and matrix transformations, Boll. Unione Mat. Ital., 13(1) (2019), 29-38.
- [2] K.G. Grosse – Erdmann, Matrix transformations between the sequence spaces of Maddox, J. Math. Anal. Appl. 180(1) (1993), 223-238.
- [3] E.E. Kara, Some topological and geometrical properties of new Banach sequence spaces, J. Inequal. Appl. (1) (2013), 38.
- [4] T. Koshy, Fibonacci and Lucas numbers with applications, 51 John Wiley and Sons, 2001.

On the Absolute Euler Summability Factors

Fadime Gökçe¹

¹ University of Pamukkale, Department of Statistics, fgokce@pau.edu.tr

ABSTRACT

By (A, B) , it is denoted the set of all sequences λ such that the series $\sum a_n \lambda_n$ is summable by the summability method B whenever the series $\sum a_n$ is summable by the summability method A . In the present study, applying the main theorems in [2] to summability factors, we give necessary and sufficient conditions on the sequence λ for $\lambda \in (|E^r, \theta|(p), |E^r, \mu|)$ and $\lambda \in (|E^r, \theta|(p), |E^r, \mu|(q))$ where $|E^r, \theta|(p), |E^r, \mu|(q)$ are the absolute Euler summability methods and θ, μ are any sequences of non-negative numbers. Finally, we obtain some results as a special case.

Key Words: Absolute summability, Euler matrix, summability factor, matrix transformations

REFERENCES

- [1] F. Gökçe and M.A. Sarıgöl, On absolute Euler spaces and related matrix operators. Proc. Nat. Acad. Sci. India Sect. A, 90(5) (2020), 769-775.
- [2] F. Gökçe and M.A. Sarıgöl, Generalization of the space $l(p)$ derived by absolute Euler summability and matrix operators, J. Inequal. Appl., (1) 2018, (2018).
- [3] K.G. Grosse – Erdmann, Matrix transformations between the sequence spaces of Maddox, J. Math. Anal. Appl. 180(1) (1993), 223-238.

On The Multilinear Riesz Bessel Transforms On Morrey Spaces

Ismail Ekincioglu¹, Cansu Keskin² and A. Züleyha Ucer

¹ Dumlupinar University, Departments of Mathematics, ismail.ekincioglu@dpu.edu.tr

² Dumlupinar University, Departments of Mathematics, cansu.keskin@dpu.edu.tr

³ Dumlupinar University, Departments of Mathematics, zuleyhauker@outlook.com

ABSTRACT

A multilinear B-maximal operator that acts on the product of m Lebesgue spaces and is smaller than the m -fold product of the B-maximal function is studied. The operator is used to obtain a precise control on multilinear singular integral operators of the Riesz Bessel transforms. A natural variant of the operator which is useful to control certain commutators of multilinear Riesz Bessel transforms with BMO functions is then considered. Finally, the Riesz Bessel transform generated by generalized translate operator on Morrey spaces are studied.

Key Words: B-maximal operator, Riesz Bessel transforms, BMO.

REFERENCES

- [1] A.P. Calderon and A. Zygmund, On the existence of certain singular integrals, Acta Math. 88 (1952), 85–139.
- [2] M. Christ and R. Fefferman, A note on weighted norm inequalities for the Hardy-Littlewood maximal operator, Proc. Amer. Math. Soc. 87 no. 3 (1983), 447–448.
- [3] R.R. Coifman and C. Fefferman, Weighted norm inequalities for maximal functions and singular integrals, Studia Math. 51 (1974), 241–250.
- [4] R.R. Coifman and Y. Meyer, On commutators of singular integrals and bilinear singular integrals, Trans. Amer. Math. Soc. 212 (1975), 315–331.
- [5] G. Di Fazio and M.A. Ragusa, Interior estimates in Morrey spaces for strong solutions to non-divergence form equations with discontinuous coefficients, J. of Functional Analysis 112 (1993), 241–256.
- [6] C. Fefferman and E.M. Stein, Some maximal inequalities, Amer. J. Math., 93 (1971), 107–115.
- [7] C. Fefferman and E.M. Stein, H^p spaces of several variables, Acta Math., 129 (1972), 137–193.

- [8] R.A. Fefferman, Multiparameter Calderón-Zygmund theory, Harmonic analysis and partial differential equations (Chicago, IL, 1996), 207–221, Chicago Lectures in Math., Univ. Chicago Press, Chicago, IL, 1999.
- [9] S.H. Ferguson and M.T. Lacey, A characterization of product BMO by commutators, Acta Math. 189 no. 2 (2002), 143–160.
- [10] J. García-Cuerva and J.L. Rubio de Francia, Weighted norm inequalities and related topics, North Holland, Amsterdam, 1985.
- [11] L. Grafakos and R.H. Torres, Multilinear Calderón-Zygmund theory, Adv. Math. 165 (2002), no. 1, 124–164.
- [12] L. Grafakos and R.H. Torres, Maximal operator and weighted norm inequalities for multilinear singular integrals, Indiana Univ. Math. J., 51 no. 5 (2002), 1261–1276.
- [13] L. Grafakos and R.H. Torres, On multilinear singular integrals of Calderón-Zygmund type, Proceedings of the 6th International Conference on Harmonic Analysis and Partial Differential Equations (El Escorial, 2000). Publ. Mat. 2002, Vol. Extra, 57–91.
- [14] M. Lacey, S. Petermichl, J. Pipher and B. Wick, Multiparameter Riesz Commutators, preprint.
- [15] M. Lacey and C. Thiele, L^p estimates on the bilinear Hilbert transform for $2 < p < \infty$, Ann. of Math. (2) 146 no. 3 (1997), 693–724.

Rate of Convergence by Bivariate Generalized Kantorovich Type Sampling Series

Serkan Ayan¹, Nurhayat İspir²

1 Bursa Technical University, Faculty of Engineering and Natural Sciences, Department of
Mathematics 16290 Bursa

serkan.ayan@btu.edu.tr

2 Gazi University, Science Faculty, Department of Mathematics 06500 Ankara

nispir@gazi.edu.tr

ABSTRACT

One of the crucial subfields of electrical engineering is Signal Processing, which focuses on analysing, modifying and synthesizing signals such as sound and images. The analysis of Kantorovich type sampling series in the one-dimensional case has an important place in terms of its application to this theory. In particular, the multivariate versions of this type sampling series include Image Processing applications and allow us to examine the reconstruction of the image. In terms of constructing generalized Kantorovich type sampling operators, it is important to consider an average of f function on a small interval including k/n values instead of using $f(k/n)$ sample values. Therefore, the reduction of the so-called "time-jitter" errors in the neighbourhood of a point rather than a point is studied using Kantorovich type sampling operators.

In [1], a generalized class of Kantorovich type sampling series is studied in the case of one variable and some pointwise convergence theorems are given. In addition, the Voronovskaja type asymptotic formula is obtained. In this study, inspired by the operators in [1], bivariate generalized Kantorovich type sampling series are discussed. Then the degree of approximation in terms of full and partial modulus of continuity is given. Furthermore, the generalized Boolean Sum (GBS) operators of these generalized Kantorovich type sampling series are introduced, a result regarding the rate of convergence to Bögel continuous functions is obtained.

Key Words: Kantorovich type sampling operators, GBS operators, mixed modulus of smoothness.

REFERENCES

- [1] C. Bordaro, I. Mantellini, On Convergence Properties for a Class of Kantorovich Discrete Operators, Numer. Funct. Anal. Optim. 33.4 (2012), 374-396.
- [2] S.K. Angamuthu, D. Ponnaian, Approximation by generalized bivariate Kantorovich sampling type series, J. Anal 2 (2019), 429-449.
- [3] P.N. Agrawal, N. Ispir, M. Sidharth, Quantitative estimates of generalized Boolean sum operators of blending type, Numer. Funct. Anal. Optim. 39.3 (2018), 295-307.
- [4] A.S. Kumar, B. Shivam, Inverse approximation and GBS Kantorovich type sampling series, RACSAM, 114.2 (2020), 1-15.

Some Mean Ergodic Type Theorems

Gencay Oğuz¹

1 Ankara University, Faculty of Science, Department of Mathematics, Tandoğan 06100
Ankara, Turkey, gencayyoguzz@gmail.com

ABSTRACT

This talk is based on a joint paper with C.Orhan [5]. A bounded linear operator T on a Banach space X is called mean ergodic if its averages $M_n(T) := \frac{1}{n} \sum_{k=1}^n T^k$ converge for every $x \in X$. The ergodic theorem asserts that if $T: X \rightarrow X$ is a bounded linear operator on a reflexive Banach space whose iterates T^j form a bounded sequence of bounded linear operators, then the sequence $\{M_n(T)\}$ converges strongly to a projection onto the kernel of the operator $I - T$.

Cohen [1] determined a class of regular infinite matrices $A = (a_{nk})$ for which

$$L_n := \sum_{k=1}^{\infty} a_{nk} T^k$$

converges strongly to an element invariant under T . He proved that such a sequence $\{L_n\}$ is strongly convergent provided that $\{L_n x: n \in \mathbb{N}\}$ is weakly compact and $\lim_{j \rightarrow \infty} \sum_{k=j}^{\infty} |a_{n,k+1} - a_{nk}| = 0$, uniformly in n . Recall that, if the matrix $A = (a_{nk})$ maps convergent sequences into the convergent sequences leaving the limit invariant, then A is called a regular matrix. It seems that Cohen's result provides a generalization of the mean ergodic theorems given by J. Von Neumann [4], F. Riesz [6] and K. Yosida [8].

In this talk, we will call an operator $T \in B(X)$ an A -mean ergodic operator if the limit of $\{L_n x\}$ exists. Then we study A -mean ergodic type theorems when $A = (a_{nk})$ is a regular infinite matrix satisfying Cohen's uniformity condition. In particular we get an ergodic decomposition. Using this we also give necessary and sufficient conditions in order that $\{T^n x\}$ is convergent.

Key Words: Ergodic theorem, Mean ergodic theorem, Ergodic decomposition, Regular matrix, Bounded linear operator

REFERENCES

- [1] L.W. Cohen, On the mean ergodic theorem. Ann. Math. 41, no:3 (1940), 505-509.

- [2] H. Mustafayev, Some convergence theorems in Fourier algebras, Bull. Austr. Math. Soc. 96 (2017), 487-495.
- [3] H. Mustafayev, Convergence of iterates of convolution operators in L_p spaces, Bull. Sci. Math. 152 (2019), 61-92
- [4] J. von Neumann, Proof of the quasi-ergodic hypothesis, Proc. Nat. Acad. Sci. USA 18 (1932), 70-82.
- [5] G. Oğuz and C. Orhan, Some mean and uniform ergodic type theorems, (submitted).
- [6] F. Riesz, Some mean ergodic theorems, J. Lond. Math. Soc. 13 (1938), 274.
- [7] T. Yoshimoto, Ergodic theorems and summability methods, Quart. J. Math. 38(3) (1987), 367-379.
- [8] K. Yosida, Mean ergodic theorem in Banach space, Proc. Imp. Acad. Tokyo, 14 (1938), 292-294.

Some Operator Identities on the Umbral Algebra

Rahime Dere Paçin¹

¹ Department of Mathematics and Science Education, Faculty of Education, Alanya Alaaddin Keykubat University Alanya/Antalya TURKEY rahimedere@gmail.com

ABSTRACT

Let \mathcal{P} be the algebra of polynomials in the single variable x over the field complex numbers. Let \mathcal{P}^* be the vector space of all linear functionals on \mathcal{P} . We use the notation

$$\langle L | p(x) \rangle$$

to denote the action of a linear functional L on a polynomial $p(x)$.

Let \mathcal{F} denote the algebra of formal power series in the variable t over \mathbb{C} . \mathcal{F} will denote both the algebra of formal power series in t and the vector space of all linear functionals on \mathcal{P} . Then we have automatically defined an algebra structure on the vector space of all linear functionals on \mathcal{P} , namely, the algebra of formal power series. This shall call \mathcal{F} the umbral algebra.

An element of \mathcal{F} plays three roles in the umbral calculus. It is a formal power series, a linear functional and a linear operator [4].

One can get interesting identities by using actions of some linear functionals and linear operators on some special polynomials, such as Sheffer polynomials.

In this work, we study some operators by using the methods of the umbral algebra. We give the actions of this operators on some special polynomials. In this way, we obtain various properties of these polynomials.

Key Words: Umbral algebra, linear operators, special polynomials.

REFERENCES

- [1] R. Dere, Y. Simsek and H. M. Srivastava, A unified presentation of three families of generalized Apostol type polynomials based upon the theory of the umbral calculus and the umbral algebra, Journal of Number Theory 133 (2013), 3245-3263.
- [2] R. Dere and Y. Simsek, Hermite Base Bernoulli Type Polynomials on the Umbral Algebra, Russian Journal of Mathematical Physics 22(1) (2015), 1-5.

- [3]** D. S. Kim, T. Kim, D. V. Dolgy and S.-H. Rim, Some new identities of Bernoulli, Euler and Hermite polynomials arising from umbral calculus, *Advances in Difference Equations*, 73 (2013).
- [4]** S. Roman, *The Umbral Calculus*, Dover Publ. Inc. New York, 2005.

Some Results Based on (p, r) –Compact Sets in Banach Spaces

Ayşegül Keten Çopur ¹ and Fatma Özceran ²

1 Necmettin Erbakan University, Faculty of Science, Department of Mathematics and Computer Science, Konya, Turkey, aketen@erbakan.edu.tr

2 Necmettin Erbakan University, The Graduate School of Natural and Applied Science, Department of Mathematics, Konya, Turkey, ftmzcrn95@gmail.com

ABSTRACT

Grothendieck [1] characterized relatively compact subsets of a Banach space as sets contained in the closed convex hull of a sequence converging to zero in the Banach space. This result is well known in the literature as Grothendieck's compactness criterion.

Inspired by this result of Grothendieck [1], which characterizes relatively compact sets in Banach spaces, Sinha and Karn [2] introduced p –compactness notion ($1 \leq p \leq \infty$), which is a stronger form of the compactness.

Later, Ain et al. [3] introduced (p, r) –compactness notion ($1 \leq p \leq \infty, 1 \leq r \leq p^*$), which is a stronger form of the p –compactness. We denote that if $p = \infty$ (thus $r = 1$), then the (p, r) –compactness coincides with the compactness and if $r = p^*$, then the (p, r) –compactness is precisely the p –compactness [3].

The approximation property, which is one of the important properties of Banach spaces theory, has been systematically studied by Grothendieck [1]. Grothendieck obtained various characterizations of the approximation property, defined some versions of the approximation property and examined the relationships between them (see [4,5]).

The p –compactness concept leaded naturally p –approximation property concept. Sinha and Karn [2] defined the p –approximation property concept by replacing compact sets in definition of the approximation property with p –compact sets. The p –compact set and the p –approximation property concepts defined by Sinha and Karn [2] have attracted great attention and been studied by many scientists. We can mention the works done by [5-8].

In this study, some results in the literature based on p –compact sets such as factorizations of p –compact operators, p –approximation property and the representation of a dual space are considered in terms of (p, r) –compact sets.

Key Words: approximation property, p –compact set, (p, r) –compact set.

REFERENCES

- [1] A. Grothendieck, Produits tensoriels topologiques et espaces nucléaires, Memoirs of the American Mathematical Society, 16 (1955).
- [2] D.P. Sinha and A.K. Karn, Compact operators whose adjoints factor through subspaces of ℓ_p , Studia Mathematica, 150 (1) (2002) 17-33.
- [3] K. Ain, R. Lillemets and E. Oja, Compact operators which are defined by ℓ_p –spaces, Quaestiones Mathematicae, 35 (2) (2012) 145–159.
- [4] J. Lindenstrauss and L. Tzafriri, Classical Banach spaces I, sequences spaces, Springer-Verlag, Berlin (1977).
- [5] P.G. Casazza, Approximation properties, in: W.B. Johnson, J. Lindenstrauss (Eds.), Handbook of the Geometry of Banach Spaces, Elsevier, Amsterdam, vol. 1 (2001), 271-316.
- [6] J.M. Delgado, E. Oja, C. Piñeiro and E. Serrano, The p –approximation property in terms of density of finite rank operators, Journal of Mathematical Analysis and Applications, 354 (1) (2009) 159-164.
- [7] Y.S. Choi and J.M. Kim, The dual space of $(\mathcal{L}(X, Y), \tau_p)$ and the p –approximation property, Journal of Functional Analysis, 259 (9) (2010), 2437–2454.
- [8] D. Galicier, S. Lassalle and P. Turco, The ideal of p –compact operators: a tensor product, approach, Studia Mathematica, 211 (2012), 269–286.

Some Results Related to New Jordan Totient Double Sequence Spaces

Sezer Erdem¹ and Serkan Demiriz²

1 Battalgazi Farabi Anatolian Imam Hatip High School, Malatya, Turkey
sezererdem8344@gmail.com

2 Department of Mathematics, Gaziosmanpasa University, Tokat, Turkey
serkandemiriz@gmail.com

ABSTRACT

The Jordan totient function $J_t: N \times N \rightarrow N, k \mapsto J_t(k)$ is described as the number of t -tuples of positive integers all less than or equal to k that form a coprime with $(t+1)$ -tuples together with k , where $k, t \in N$ and $N = \{1, 2, 3, \dots\}$. J_t is multiplicative and furthermore it is an arithmetic function and it is one of many generalizations of the famous Euler totient function ϕ . The 2 dimensional Jordan totient matrix and its domain on the space of l_p of absolutely p -summable single sequences are described and examined by İkhan et al. [5]. After that, compact operators on the matrix domain of the Jordan totient matrix have been studied in [6]. 4 dimensional Jordan totient matrix and its associated double sequence spaces have been examined in [4]. In the present paper, first of all we define two new double sequence spaces by using the 4 dimensional Jordan totient matrix derived by the Jordan totient function and we show that this newly described double sequence spaces are Banach spaces with their norms. Then, we give a inclusion relation including this spaces. Moreover, we compute the α -, β (bp) - and γ -duals and finally, we characterize some new 4 dimensional matrix transformation classes and complete this work with some significant results.

Key Words: Jordan totient function, RH-regular matrix, matrix domain, double sequence space, α -, β (bp) - and γ -duals, matrix transformations.

REFERENCES

- [1] C.R. Adams, On non-factorable transformations of double sequences, Proc. Natl. Acad. Sci. USA, 19(5) (1933), 564-567.
- [2] S. Demiriz and S. Erdem, Domain of Binomial Matrix in Some Spaces of Double Sequences, Punjab Univ. J. Math., 52(11) (2020), 65-79.

- [3] S. Erdem and S. Demiriz, Almost Convergence and 4-Dimensional Binomial Matrix, Konuralp J. Math., 8(2) (2020), 329-336.
- [4] S. Erdem and S. Demiriz, A New RH-Regular Matrix Derived by Jordan's Function and Its Domains on Some Double Sequence Spaces, Journal of Function Spaces, vol. 2021, Article ID 5594751, (2021) 9 pages.
- [5] M. İlkhan, N. Şimşek and E. E. Kara, A new regular infinite matrix defined by Jordan totient function and its matrix domain in l_p , Math Meth Appl Sci. 2020;1-12.
- [6] M. İlkhan, E. E. Kara and F. Usta, Compact Operators on the Jordan Totient Sequence Spaces, Mathematical Methods in the Applied Sciences 44(2), (2020).
- [7] O. Tuğ, Four-dimensional generalized difference matrix and some double sequence spaces. J. Inequal. Appl., 2017(1), 149 (2017).
- [8] M.Yeşilkayagil and F. Başar, Domain of Euler mean in the space of absolutely p-summable double sequences with $0 < p < \infty$, Anal. Theory Appl., 34, (2018), No:3, 241-252.

Spanne-type Result for the Generalized Fractional Maximal Operator in the Vanishing Generalized Weighted Local Morrey Spaces

Abdulhamit Kucukaslan

Faculty of Applied Sciences, Pamukkale University, 20680, Denizli, Turkey
kucukaslan@pau.edu.tr

ABSTRACT

Morrey spaces were introduced by Morrey in [3]. These spaces appeared to be useful in the study of local behavior properties of the solutions of second order elliptic PDEs. The vanishing Morrey space of the classical Morrey spaces was introduced by Vitanza in [7] and applied there to obtain a regularity result for elliptic PDEs. Later in [8] Vitanza proved an existence theorem for a Dirichlet problem, under weaker assumptions than those introduced by Miranda in [2] and a regularity result assuming that the partial derivatives of the coefficients of the highest and lower order terms belong to vanishing Morrey spaces depending on the dimension. Persson et al. [4] showed the commutators of Hardy operators on vanishing Morrey spaces. Also Ragusa [5] obtained a sufficient condition for commutators of fractional integral operators to belong to vanishing Morrey spaces.

The vanishing generalized Morrey space and vanishing generalized local Morrey space was introduced by Samko in [6]. The boundedness of the multi-dimensional Hardy type operators, maximal, potential and singular operators in these spaces were proved in [6]. Kucukaslan et al. [1] proved the Spanne-type and Adams-type boundedness of generalized fractional integral operators on vanishing generalized local Morrey spaces.

The generalized fractional maximal operators M_p was initially investigated in by Nakai in 1994. Nakai introduced the generalized Morrey spaces and proved the boundedness of the generalized fractional integral operator in these spaces. Nowadays many authors have been culminating important observations about the generalized fractional maximal operators M_p especially in connection with Morrey-type spaces.

In this talk, we prove the Spanne-type boundedness of the generalized fractional maximal operator M_ρ from the vanishing generalized weighted local Morrey spaces $VLM_{p,\varphi_1}(\omega^p)$ to another vanishing generalized weighted local Morrey spaces $VLM_{q,\varphi_2}(\omega^q)$ with ω^q , and from the vanishing generalized weighted local Morrey spaces $VLM_{1,\varphi_1}(\omega)$ to the vanishing generalized weighted weak local Morrey spaces $VWLM_{q,\varphi_2}(\omega^q)$ with ω . Also, we get a new theorem for the Spanne-type result of the generalized fractional maximal operator M_ρ from the vanishing generalized local Morrey spaces VLM_{p,φ_1} to vanishing generalized local Morrey spaces VLM_{q,φ_2} , including weak estimates. The all weight functions belong to Muckenhoupt-Weeden classes $A_{p,q}$.

Key Words: Generalized fractional maximal operator, Vanishing generalized weighted local Morrey space, Vanishing generalized Morrey space, Muckenhoupt-Weeden classes.

REFERENCES

- [1] A. Kucukaslan, S. G. Hasanov, C. Aykol, Generalized fractional integral operators on vanishing generalized local Morrey spaces, *Int. J. of Math. Anal.*, 11(6) (2017), 277--291.
- [2] C. Miranda, Sulle equazioni ellittiche del secondo ordine di tipo non variazionale, a coefficienti discontinui, *Ann. Math. Pura E Appl.*, 63(4) (1963), 353-386.
- [3] C. B. Morrey, On the solutions of quasi-linear elliptic partial differential equations, *Trans. Amer. Math. Soc.*, 43 (1938), 126-166.
- [4] L. E. Persson, M. A. Ragusa, N. Samko, P. Wall, Commutators of Hardy operators in vanishing Morrey spaces, *AIP Conf. Proc.* 1493, 859, 2012.
- [5] M. A. Ragusa, Commutators of fractional integral operators on vanishing-Morrey spaces, *J. Global Optim.* 40(1-3) (2008), 361-368.
- [6] N. Samko, Weighted Hardy operators in the local generalized vanishing Morrey spaces, *Positivity* 17 (2013), 683-706.
- [7] C. Vitanza, Functions with vanishing Morrey norm and elliptic partial differential equations, *Proceedings of Methods of Real Analysis and Partial Differential Equations*, Capri, Springer (1990), 147-150.
- [8] C. Vitanza, Regularity results for a class of elliptic equations with coefficients in Morrey spaces, *Ricerche di Matematica* 42(2) (1993), 265-281.

Upper and lower θ_p -continuous multifunctions

Aynur KESKİN KAYMAKCI ¹, Ayşe Nazlı ÜRESİN ²

¹ Selcuk University Faculty of Sciences, Department of Mathematics, Campus, 42030,
Konya/Turkey, akeskin@selcuk.edu.tr

² Ahep University, Cikilli, Saraybeleni Street, 07400, Nr:7, Alanya, Antalya/Turkey,
nazliuresin@gmail.com

ABSTRACT

By a multifunction $F:X \rightarrow Y$, we mean a point-to-set correspondence from X into Y . The aim of this presentation is to introduce and study the notion of upper (resp. lower) θ_p -continuous multifunctions. Basic properties of upper (resp. lower) θ_p -continuous multifunctions and relationships between upper (resp. lower) θ_p -continuous multifunctions and graphs are also investigated.

Key Words: preopen sets, preclosure, multifunctions, upper (resp. lower) θ_p -continuous multifunctions.

REFERENCES

- [1] T. Banzaru, Multifunctions and M-product spaces (Romanian). Bull Stiin. Teh. Inst Politeh Timisoara Ser Mat Fiz Mer Teor Apl, 17(31) (1972), 17-23.
- [2] T. Banzaru, On the upper semicontinuity of the upper topological limit for multifunction nets. Semin Mat Fiz Inst Politeh Timisoara (1983), 59-64.
- [3] C. Berge, Espaces topologiques fonctions multivoques. Paris: Dunod; 1959.
- [4] M. Caldas, S. Mafari, T. Noiri and M. Simões, A new generalization of contra-continuity via Levine's g-closed sets. Chaos, Solitons & Fractals 32 (2007), 1597-1603.
- [5] S.H. Cho, A note on strongly θ -precontinuous functions. Acta Math. Hungar, 101(1-2) (2003), 173-178.
- [6] J. Dontchev, M. Ganster and T. Noiri, On p-closed spaces. Internat. J. Math. Math. Sci, 24 (2000), 203-212.
- [7] S.N. El-Deeb, I.A. Hasanein, A.S. Mashhour and T. Noiri, On p-regular spaces. Bull Math Soc Sci Math R. S. Roumanie 27(75) (1983), 311-315.
- [8] Mashhour AS, Abd El-Monsef ME, El-Deeb SN. On precontinuous and weak precontinuous mappings. Proc Math Phsy Soc Egypt 1982;53:47-53.

APPLIED MATHEMATICS

A Mathematical Investigation of COVID-19 in Turkey Using a Susceptible Infectious Recovered (SIR) Type Model

Agâh Akin¹, Harun Baldemir² and Ömer Akin³

1 Bitlis Tatvan State Hospital, 13200, Bitlis, Turkey
agh_akn@hotmail.com

2 Department of Mathematics, Faculty of Science, Cankiri Karatekin University, 18100, Cankiri
Turkey

HarunBaldemir@karatekin.edu.tr

3 Department of Mathematics, Faculty of Sciences and Arts, TOBB Economics and Technology
University, 06650, Ankara, Turkey
omerakin@etu.edu.tr

ABSTRACT

At the end of 2019, a novel coronavirus (abbreviated as COVID-19) was identified in Wuhan, China. It rapidly spreads, resulting in an epidemic throughout China, followed by an increasing number of cases in other countries throughout the World. The World Health Organization (WHO) declared COVID-19 a global pandemic on 11 March 2020 when there were more than 100.000 reported cases and 4.000 deaths across 114 countries.

In [1] we applied the Fuzzyfied Richards Growth Model to understand the dynamic behaviour of the COVID-19 based on the real data and predicted possible future scenarios applying fuzzy approaches for some countries around the world including China, the United States, the top five countries with the highest population in Europe and Turkey.

In this joint work, we investigate the dynamic behaviour of COVID-19 in Turkey using the generalized SEIR (Susceptible Exposed Infectious Recovered) Model [2] that includes seven different states namely susceptible, insusceptible, exposed, infectious, quarantined, recovered and death. After that, by getting help from [1] we extend the generalized SEIR Model to the fuzzy environment. Finally we try to obtain lower and upper solutions by using Zadeh's extension principle [3] in order to approximate the spread of COVID19 among people in Turkey.

Key Words: COVID19, SIR Model, Zadeh's Extension Principle, Fuzzy Environment.

REFERENCES

- [1]** H. Baldemir, A. Akin and Ö. Akin, Fuzzy Modelling of Covid-19 in Turkey and Some Countries in The World. Turkish Journal of Mathematics and Computer Science, 12(2) (2020), pp.136-150.
- [2]** L. Peng, W. Yang, D. Zhang, C. Zhuge and L. Hong, 2020. Epidemic analysis of COVID-19 in China by dynamical modeling. arXiv preprint arXiv:2002.06563.
- [3]** L.A. Zadeh, The concept of a linguistic variable and its application to approximate reasoning-III. Information sciences, 9(1) (1975), pp.43-80.

A NOTE ON MONOTONE TECHNIQUE FOR A CLASS OF FRACTIONAL INTEGRO-DIFFERENTIAL EQUATIONS

Ali YAKAR¹, Hadi KUTLAY²

¹Department of Mathematics, Tokat Gaziosmanpaşa University, Tokat, Turkey,
aliyakar@gmail.com

²Department of Mathematics, Tokat Gaziosmanpaşa University, Tokat, Turkey,
hkutlay.tokat@gmail.com

ABSTRACT

In this paper, we consider the following nonlinear Caputo fractional integro-differential equation

$${}^c D^{q_1} u(t) = F(t, u(t), {}^c I^{q_2} u(t))$$

with boundary condition

$$g(u(0), u(T)) = 0$$

where $F \in C[J \times R \times R_+, R]$, $J = [0, T]$ and $0 < q_2 \leq q_1 < 1$.

We investigate the existence of the solutions of the Caputo fractional integro-differential equation by applying monotone iterative technique with the method of upper and lower solutions. The monotone iterative technique, with the method of upper and lower solutions, offers monotone sequences that converge uniformly and monotonically to minimal and maximal solutions of the problem considered. Since each member of such sequence is the solution of the linear initial value problem of Caputo fractional integro-differential equations which can be explicitly computed, the advantage and the importance of this technique needs no special emphasis. The monotone iterative technique can be applied to many nonlinear fractional differential equations. Moreover, this method can successfully be employed to generate two sided pointwise bounds on solutions of the linear initial value problem of Caputo fractional integro-differential equations, from which qualitative and quantitative behaviour can be investigated. Furthermore, if uniqueness conditions are satisfied the minimal and maximal solutions will converge to the unique solution of the nonlinear Caputo fractional integro-differential equation.

Key Words: Caputo Derivative, Integro-differential equation, Monotone Iterative Technique, Maximal and Minimal Solutions

REFERENCES

- [1] A. A. Kilbas, H. M. Srivatsava and J. J. Trujillo, Theory and Applications of Fractional Differential Equations, Elsevier, Amsterdam, 2006.
- [2] V. Lakshmikantham, S. Leela and Vasundhara Devi, J. Theory of Fractional Dynamic Systems, Cambridge Academic Publishers, Cambridge, 2009.
- [3] G. S. Ladde, V. Lakshmikantham and A. S. Vatsala, Monotone Iterative Techniques for Nonlinear Differential Equations, Pitman Publishing Company, Boston, 1985.
- [4] C. Yakar and A. Yakar, Monotone iterative techniques for fractional order differential equations with initial time difference, Hacettepe Journal of Mathematics and Statistics 40(2) (2011), 331-340.
- [5] A. Yakar and H. Kutlay, Monotone iterative technique via initial time different coupled lower and upper solutions for fractional differential equations. Filomat, 31(4) (2017), 1031-1039.
- [6] A. Yakar and H. Kutlay ,A note on comparison results for fractional differential equations. In AIP Conference Proceedings (Vol. 1676, No. 1, p. 020064), 2015.
- [7] J. D. Ramirez and A. S. Vatsala, "Monotone method for nonlinear Caputo fractional boundary value problems," Dynamic Systems and Applications, vol. 20, no. 1 (2011), pp. 73–88.
- [8] J. Vasundhara Devi, Ch. V. Sreedhar, Monotone iterative method for Caputo fractional integro-differential equations, European Journal of Pure and Applied Mathematics, Vol. 9, No. 4 (2016), 346-359.

A Slough/Wound Interaction Model Accounting for Cellular Diffusion

Rıdvan Yaprak¹, Erhan Coşkun²

1 Karadeniz Technical University, Faculty of Science, Department of Math,
ridvanyaprak@ktu.edu.tr

2 Karadeniz Technical University, Faculty of Science, Department of Math,
erhan@ktu.edu.tr

ABSTRACT

In this study we consider Slough/Wound interaction model, called SWODE, a nonlinear Ordinary Differential Equation, ODE system, developed by Jones et. al in 2003 [1] that describes evolution of wound and slough area.

We first nondimensionalize the model and estimate a larger domain of attraction for the zero equilibrium solution with an appropriately formulated Liapunov function. Thus, we determine the region where the wound heals and slough tissue disappears in terms of nondimensional model parameters.

Next, we extend the SWODE model to account for one dimensional cellular diffusion resulting in a model, called SWPDE1, a one dimensional nonlinear system of Partial Differential Equations. Evolution of wound and slough interactions is described and equilibrium solutions with lower and upper solutions are determined. The uniqueness of corresponding stationary system is also handled numerically and qualitative behaviour of equilibrium solutions are analysed.

Finally, we further extend SWPDE1 to account for two dimensional cellular diffusion, resulting in a model, SWPDE2, a two dimensional time-dependent nonlinear PDE system. In this model, we consider rectangular wound and slough within a rectangular region and investigate wound-slough interaction as the healing process goes on. Method of lines is used to obtain a nonlinear system of ODEs and MATLAB ODE solvers [2] are used to integrate the resulting system.

Key Words: Wound healing, nonlinear ordinary differential equation system, nonlinear partial differential equation system, ODE solvers of MATLAB.

REFERENCES

- [1] M.A. Jones, B. Song and D.M. Thomas, Controlling Wound Healing Through Debridement, *Mathematical and Computer Modelling* 40 (2004), 1057-1064.
- [2] L.F. Shampine, M.W. Reichelt, The Matlab ODE Suite, *SIAM Journal on Scientific Computing*, 18(1997), 1-22.

ANALYSIS OF CONTINUOUS DATA ASSIMILATION SCHEME FOR THE NAVIER-STOKES EQUATIONS USING VARIATIONAL MULTISCALE METHOD

Gülnur Haçat¹, Mine Akbaş² and Aytekin Çibık³

¹ Department of Mathematics, Gazi University, 06550, Ankara, gulnur.hacat@gazi.edu.tr

² Department of Mathematics, Duzce University, 81620, Düzce, mineakbas@duzce.edu.tr

³ Department of Mathematics, Gazi University, 06550, Ankara, abayram @gazi.edu.tr

ABSTRACT

In this study, we analyse a continuous data assimilation (CDA) [1] scheme which enables us to combine an observable data with a numerical method to obtain better solutions in which these solutions are also closely similar to the current state of the system. The scheme applied on a Navier-Stokes system which is discretized with two-step Backward Differentiation (BDF2) in time and finite element in space. Also the nonlinear term arises in the system is treated with extrapolation techniques. In order to improve the accuracy and prevent some non-physical oscillations due to the effect of small viscosity and the dominance of convection, a projection based variational multiscale method (VMS) [2] was also cast on the system. The corresponding CDA scheme with VMS is given by,

$$\begin{aligned} & \left(\frac{3y_h^{n+1} - 4y_h^n + y_h^{n-1}}{2\Delta t}, v_h \right) + \text{Re}^{-1}(\nabla y_h^{n+1}, \nabla v_h) + b(2y_h^n - y_h^{n-1}, y_h^{n+1}, v_h) \\ & - (p_h^{n+1}, \nabla \cdot v_h) + (\sigma(I - P_H)\nabla y_h^{n+1}, (I - P_H)\nabla v_h) + \mu(I_H(y_h^{n+1} - u^{n+1}), v_h) = (f^{n+1}, v_h) \\ & (\nabla \cdot y_h^{n+1}, q_h) = 0 \end{aligned}$$

where μ is the nudging parameter, I_H is the interpolation operator, σ in VMS term is the eddy viscosity parameter.

The nudging term μ pushes the large spatial scales of the approximating solution toward those of the reference solution while the viscosity stabilizes and dissipates the fine spatial scales and any spillover into the fine scales caused by the nudging term. That's why we added this term to the equation we studied. In the rest of the study, the long-time stability and long-time convergence analyses of the

scheme are presented in detail and several numerical tests are given in order to support theoretical findings and demonstrate the promise of the method.

Key Words: Continuous data assimilation, Navier-Stokes equations, variational multiscale method.

REFERENCES

- [1] A. Larios and Y. Pei, "Nonlinear continuous data assimilation". arXiv:1703.03546, 2017.
- [2] T. Hughes, L. Mazzei, and K. Jansen, "Large eddy simulation and the variational multiscale method", Comput. Vis. Sci., 3 (2000), pp. 47–59.

Analysis of Covid-19 epidemic in Turkey using Generalized SEIR Model

Ersan ERDEM¹, Gülnur ÇELİK KIZILKAN²

1 Necmettin Erbakan University, Institute of Science Department of Mathematics, Konya
ersanerdem.ee@gmail.com

2 Necmettin Erbakan University, Faculty of Science Department of Mathematics-Computer,
Konya, gckizilkan@erbakan.edu.tr

ABSTRACT

A new type of coronavirus called SARS-CoV-2, or Covid-19 was first identified in Wuhan city, Hubei Province of China at the end of 2019. In the following weeks, Covid-19 has rapidly spread to all over China and to many countries. So, WHO has declared the disease as a pandemic on 11 March 2020. Covid-19 is a respiratory tract disease which the human-to-human transmissible, can cause symptoms including fever, difficulty in breathing, dry cough. Many countries have taken several strict precautions such as limiting travel among cities, closing public area, quarantining cities to be able to control the contagious disease.

Covid-19 has become doubtlessly one of humanity's biggest epidemic problems. This epidemic has generated significant social, economic, and health problems. And many people died. Therefore, it has gained importance predicting how to progress of epidemic. Mathematical models which is based on a series of dynamic mathematical equation play an important role in epidemiology. The SIR model which is the well-known basic infectious disease model in the literature, was defined in 1927 by Kermack and McKendrick (Kermack and McKendrick 1927). Later, by adding new compartments to the SIR model, some infectious disease models such as SEIR, SEIRD which can simulate real life a little better were obtained (see, for instance, Li et al 1999, Weitz and Dushoff 2015 etc). Many study on the generalized version of the classical SEIR model known as SEIRDP model have been made after Covid-19 epidemic outbreak (see, for instance, Godio et al 2020, Afonso et al 2020, Peng et al 2020).

In this study, we have tried to apply to SEIRDP infectious disease model the spreading of Covid-19 in Turkey. For this, we have obtained real data that is publicly available at official web site of Turkey Government Ministry of Health by using web scraping program that we have written with Python programming language (Turkey Government Ministry of Health 2021, Python 2021). We have tried to analyze epidemic and to determine parameters of that model using nonlinear least squares method. We have aimed to be able to evaluate the daily number of infectious cases, recovered cases and deaths, the consequences of the precautions taken.

Key Words: Covid-19, epidemic models, SIR, generalized SEIR, python.

REFERENCES

- [1] W. O. Kermack and A. G. McKendrick ,A contribution to the mathematical theory of epidemics. Proc. royal society london.Ser. A, Containing papers a mathematical physical character 115, (1927), 700–721.
- [2] Y. Li Michael, R. John, Graef, Liancheng Wang Janos Karsai, Global dynamics of a SEIR model with varying total population size, Mathematical Biosciens, 160 (1999),191-213.
- [3] A. Godio, F. Pace and A. Vergnano, SEIR Modeling of the Italian Epidemic of SARS-CoV-2, preprints, doi:10.20944/preprints202004.0073.v1, (2020).
- [4] S.M.S Afonso, J.S. Azevedo and M.H.G. Silva, Epidemic analysis of Covid-19 in Brazil by a generalized SEIR model, (2020).
- [5] Weitz J, Dushoff J. Modeling post-death transmission of ebola: challenges for inference and opportunities for control. Scientific reports, 5 (2015), 8751.
- [6] L. Peng, W Yang, D Zhang, C. Zhuge, L. Hong, Epidemic analysis of COVID-19 in China by dynamical modeling; Epidemiology, 2020.
- [7] Python, <https://www.python.org/>, Access date: 20.08.2021.
- [8] Turkey Government Ministry of Health <https://covid19.saglik.gov.tr/EN-69532/general-coronavirus-table.html>, Access date: 20.08.2021.

Decay Estimate for the Klein-Gordon Equation in Anti de Sitter Spacetime

Muhammet Yazıcı

Department of Mathematics, Karadeniz Technical University, Trabzon, 61080, Turkey,
m.yazici@ktu.edu.tr

ABSTRACT

The Klein-Gordon equation arises in relativistic physics such as cosmology, general relativity and quantum field theory. In this study, we consider the following initial value problem for the Klein-Gordon equation in the Anti-de Sitter space time,

$$\begin{aligned}\Phi_{tt} + nH\Phi_t - e^{2Ht}\Phi + m^2\Phi &= 0, & (x, t) \in \mathbb{R}^n \times [0, \infty), \\ \Phi(x, 0) &= \varphi_0(x), & \Phi_t(x, 0) = \varphi_1(x), & x \in \mathbb{R}^n,\end{aligned}$$

where $m > 0$ is the physical mass and H is denoted by Hubble constant. Here, the sign of H is negative which describes the contraction spatially for the model of the universe. If the sign of H is positive, then the so-called de Sitter space time model describes the expansion spatially. When H is zero, the model is called Minkowski space time. The fundamental solution of the initial value problem is shown in [1]. Then, $L^p - L^q$ decay estimate for the solution of the problem is obtained in [2]. Moreover, in [3], L^∞ decay estimate for the solution of the zero initial value problem with source term is obtained. Decay estimates play an important role to prove the global existence of the solutions for the nonlinear partial differential equations. Therefore, we prove the L^∞ decay estimate for the solution of the initial value problem nonzero initial data.

Key Words: Anti-de Sitter spacetime, Klein-Gordon equation, L^∞ estimate.

REFERENCES

- [1] K. Yagdjian and A. Galstian, The Klein-Gordon Equation in Anti-de Sitter Space-time, Rend. Sem. Mat. Univ. Pol. Torino 67(2) (2009), 271-292.
- [2] A. Galstian, $L^p - L^q$ Decay Estimates for the Klein-Gordon Equation in Anti-de Sitter Space-time, Rend. Istit. Mat. Univ. Trieste 42 (2010), 27-50.

- [3]** M. Yazici, A Remark on the Decay Property for the Klein-Gordon Equation in Anti-de Sitter Space Time, New Trends Math. Sci. 5(4) (2017), 142-147.

Dynamical Behavior of Rational Difference Equation

$$x_{n+1} = \frac{x_{n-11}}{\pm 1 \pm x_{n-1} x_{n-3} x_{n-5} x_{n-7} x_{n-9} x_{n-11}}$$

Burak Oğul¹, Dağıstan Şimşek²

1 Department of Management Information Systems, Istanbul Aydın University, Istanbul, Turkey.
burakogul@aydin.edu.tr

2 Department of Engineering Basic Sciences, Konya Technical University, Konya, Turkey.
dsimsek@ktun.edu.tr

ABSTRACT

The study and solution of nonlinear rational recursive sequence of high order is quite challenging and rewarding. In the recent times, nonlinear difference equations have a critical role in the fields of physics, economy, ecology and computational science engineering, etc. Many researchers have investigated the behavior of the solution of nonlinear difference equations. So, recently there has been an increasing interest in the study of qualitative analysis of rational difference equations.

Our aim in this paper is to investigate the behavior of the solution of the following nonlinear difference equations

$$x_{n+1} = \frac{x_{n-11}}{\pm 1 \pm x_{n-1} x_{n-3} x_{n-5} x_{n-7} x_{n-9} x_{n-11}}, n \in \mathbb{N}_0$$

where the initial conditions are arbitrary nonzero positive real numbers. Also, we get explicit forms of the solutions.

Key Words: Difference equations, local stability, recursive sequences.

REFERENCES

- [1] C. Cinar, T. Mansour and I. Yalcinkaya, On the difference equation of higher order, *Utilitas Mathematica*, 92 (2013), 161-166.
- [2] M.A.E. Abdelrahman and O. Moaaz, On the New Class of The Nonlinear Rational Difference Equations, *Electronic Journal of Mathematical Analysis and Applications*, 6 (1) (2018), 117-125.
- [3] M. Ari and A. Gelisken, Periodic and asymptotic behavior of a difference equation. *Asian-European Journal of Mathematics*, 12(6) (2019), 2040004.

- [4] M.R.S. Kulenovic and G. Ladas, Dynamics of second order rational difference equations. Chapman & Hall/CRC, Boca Raton, FL, 2002. With open problems and conjectures.
- [5] R.P. Agarwal, Difference Equations and Inequalities. 1st edition, Marcel Dekker, New York, 1992, 2nd edition, 2000.

Existence Of Solutions For A Third-Order Boundary Value Problem With Integral Boundary Conditions On Infinite Intervals

Fercan Filiz¹, Erbil Çetin²

1 Department of Mathematics, Ege University, Bornova, İzmir 35100 Turkey,
fercanfilizz@gmail.com

2 Department of Mathematics, Ege University, Bornova, İzmir 35100 Turkey,
erbil.cetin@ege.edu.tr

ABSTRACT

This paper deal with the existence of bounded or unbounded solutions for the following third order three-point integral conditional boundary value problem on a half-line:

$$x'''(t) + r(t)g(t, x(t), x'(t), x''(t)) = 0, \quad t \in (0, \infty),$$

$$x(0) = \int_0^\eta x(s)ds, \quad x'(0) = A, \quad x'(\infty) = B,$$

where $\eta \in (0, \infty)$, $r: (0, \infty) \rightarrow (0, \infty)$ and $g: (0, \infty) \times \mathbb{R}^3 \rightarrow \mathbb{R}$ satisfies Nagumo's condition which plays an important role in the nonlinear term depend on the first-order derivative explicitly. This subject "third-order boundary value problem with integral boundary conditions on infinite intervals" is important and also complicated. Our main aim is to show that there is at least one solution to the considering problem. First we create the Green function for our problem. Moreover we define the lower and upper solutions and Nagumo's condition. By applying Schäuder's fixed point theorem and the lower and upper solutions method, we obtain the expected result. Then, we indicate the existence of at least one solution of the problem by determining suitable and sufficient conditions in our problem.

Key Words: Three-point integral conditional boundary value problem, lower and upper solutions, half-line, Nagumo's condition, Schäuder's fixed point theorem.

REFERENCES

- [1] A. Boucherif, Third order boundary value problems with integral boundary conditions, Nonlinear Anal. 70 (2009), 364–371.
- [2] Agarwal RP, Çetin E 2015, Unbounded solutions of third order three point boundary value problems on a half-line, Adv.Nonlinear Analysis, (2015).
- [3] Akcan U, Çetin E 2018, / Filomat 32:1(28) (2018), 341-353
- [4] L. Jackson, K. Schrader 1971, third order differential equations, journal of differential equations 9 (1971), 46-54.
- [5] Y. Wang, W. Ge 2007, Existence of solutions for a third order differential equation with integral boundary conditions, Comput. Math. Appl. 53 (2007), 144–154.

Fitted Numerical Method for a Volterra Delay Integro-Differential Equation

İlham AMIRALI¹, Gabil M. AMIRALIYEV² and Hülya ACAR³

1 Duzce University, Faculty of Science of Mathematics, 81620, Duzce, Turkey,
ailhame@gmail.com

2 Erzincan Binali Yıldırım University, Faculty of Science of Mathematics, 24100, Erzincan,
Turkey, gabilamirali@yahoo.com

3 Duzce University, Faculty of Science of Mathematics, 81620, Duzce, Turkey,
hulyaacar98@gmail.com

ABSTRACT

In this work, the authors propose a finite difference technique for approximating solutions of first-order Volterra delay integro-differential equation. The presented numerical method acquires a second-order convergence in discrete maximum norm. The derived results are numerically validated in test problems to support the theoretical analysis.

Key Words: Volterra integro- differential equation, delay, maximum norm.

REFERENCES

- [1] E. Adel, A. Elsaid and M. El-Agamy, Adaptive finite element method for Fredholm integro-differential equation, Communications on Advanced Computational Science with Applications, 1 (2017), 24–32.
- [2] I. Amirali, Analysis of higher order difference method for a pseudo-parabolic equation with delay, Miskolc Mathematical Notes, 20(2) (2019), 755–766.
- [3] S.G. Gal, Shape-Preserving Approximation by Real and Complex Polynomials, Birkhäuser Boston, Inc., Boston, MA, 2008.
- [4] I. Amirali, S. Cati and G. M. Amiraliyev, Stability inequalities for the delay pseudoparabolic equations, International Journal of Applied Mathematics, 32(2) (2019), 289–294.
- [5] I.G. Amiraliyeva, F. Erdogan and G. M. Amiraliyev, A uniform numerical method for dealing with a singularly perturbed delay initial-value problem, Appl. Math. Letters, 23 (2010), 1221–1225.
- [6] G. M. Amiraliyev, M. E. Durmaz and M. Kudu, Uniform convergence results for singularly perturbed Fredholm integro-differential equation, J. Math. Anal., 9(6) (2018), 55–64.
- [7] G. M. Amiraliyev, M. E. Durmaz and M. Kudu, Fitted second order numerical method for a singularly perturbed Fredholm integro-differential equation, Bull. Belg. Math. Soc. Simon Stevin, 27 (2020), 71–88.
- [8] G. M. Amiraliyev and O. Yapman, " On the Volterra delay integro-differential equation with layer behavior and its numerical solution, Miskolc Mathematical Notes, 20(1) (2019), 75–87.

- [9] G. M. Amiraliyev, O. Yapman and M. Kudu, "A fitted approximate method for a Volterra delay integro-differential equation with initial layer, Hacettepe Journal of Mathematics and Statistics, 48(5) (2019), 1417–1429.

Fuzzy Similarity Of Subalpine, Steppe And Segetal Vegetation For Black Sea Region In Turkey

M. Ümit BINGÖL¹, Sanem AKDENİZ ŞAFAK², Ömer AKIN^{3*}

¹ Ankara University, Department of Biology, Döğol Street, 06100 Beşevler/Ankara/Turkey
mumit1111@hotmail.com,

²Aksaray University, Department of Biology, Aksaray University Campus Road, 68100
Aksaray/Turkey, sanem.akdeniz@hotmail.com, sakdeniz@aksaray.edu.tr,

³TOBB ETU, Department of Mathematics, Söğütözü Street, 06650 Ankara/Turkey
omerakin@etu.edu.tr

ABSTRACT

In this study, the results obtained in subalpine, steppe and segetal vegetation in the research project on plant sociology conducted in Black Sea region (Turkey) has been evaluated with the fuzzy similarity measures which is an important approach in applied mathematics. The similarity relations between the sample areas included in each plant community were calculated and interpreted. The issue of fuzzy similarity of sets and elements in sets is used. This study brings a different perspective to new researches in plant sociology. According to this point of view, the fuzzy similarity among plant associations and relevés came to the fore.

Fuzzy similarity is a highly preferred method in all applied sciences for a better understanding of the subject, with its practicality, time saving and contribution to the economy.

For the sake of mathematics point of view, we give the formulas for the fuzzy similarity among plant communities below:

Fuzzy Similarity of Two Fuzzy Sets:

$$FS (AS_i, AS_j) = \max \{ \min \{ \mu_{AS_i}(r_k), \mu_{AS_j}(r_k) \} \}; k = 1(1)62, i, j = 1(1)5. (1)$$

$$r_k \in AS_i \subset X$$

Fuzzy Similarity of Two Elements:

$$FS_e(r_k, r_m) = \max \{ \min \{ \mu_{AS_i}(r_k), \mu_{AS_i}(r_m) \} \}; k, m = 1(1)30, i = 1(1)3 (2).$$

Key Words: Fuzzy Similarity, Plant Community, Phytosociology, Subalpine, Steppe, Segetal vegetation, Ecology.

REFERENCES

- [1] S. Akdeniz Şafak, Ö. Akın and M.Ü. Bingöl (2019). Using fuzzy logic in nomenclature of plant communities and determining syntaxonomic levels. International Conference on Mathematics and Mathematics Education (ICMME 2019), Selçuk University, Konya, July 11-13, 2019, Mathematics in Konya, Land of Tolerance, Abstract Book, p. 188, ISBN: 978- 605-184-176-2. (Oral Presentation).
- [2] M.Ü. Bingöl, F. Geven and K. Güney (2007) "A Phytoecological and Phytosociological Research on the Sakarat Mountain (Amasya)". Research Project. The Scientific and Technological Research Council of Turkey (TÜBİTAK). TÜBİTAK TOVAG-HD 105O018 (In Turkish).
- [3] Bingöl MÜ, Akdeniz Şafak S, Akın Ö (2020). The Analysis of Plant Communities Belonging to Forest Vegetation of Sakarat Mountain (Amasya) in Fuzzy Similarity Environment. First Online Conference on Modern Fractional Calculus and Its Applications (OCMFCA-2020), December 4-6, 2020, Biruni University, Istanbul, Turkey. Sözlü sunum / Oral Presentation.
- [4] Braun-Blanquet J (1932). Plant Sociology. (transl. by G. D. Fuller and H. S. Conard), Mc Graw Hill, New York and London.
- [5] Braun-Blanquet J, Pavillard J (1925). Vocabulary for plant sociology.
- [6] E. Feoli, G. Ferro and Ganis P (2006). Validation of phytosociological classifications based on a fuzzy set approach. Community Ecology. 7:99-108. <https://doi.org/10.1556/ComEc.7.2006.1.10>
- [7] E. Feoli and L. Orlóci (2011). Can similarity theory contribute to the development of a general theory of the plant community. Community Ecology. 12:135-141.
- [8] Zadeh L.A. (1971) Similarity relations and fuzzy orderings. Information Sciences. 3(2):177-200. [https://doi.org/10.1016/S0020-0255\(71\)80005-1](https://doi.org/10.1016/S0020-0255(71)80005-1).

Global Existence and Uniqueness of The Inviscid Velocity-Vorticity Model of The g-Navier-Stokes Equations

Özge Kazar¹, Meryem Kaya²

¹ Department of Electronics and Automation, Vocational School of Technical Sciences, Hitit University, Çorum, Turkey, ozgekazar@hitit.edu.tr

² Department of Mathematics, Faculty of Science, Gazi University, Ankara, Turkey, meryemk@gazi.edu.tr

ABSTRACT

In this talk we consider a velocity-vorticity formulation for the g-Navier-Stokes equations. The system is performed by entegrating the velocity-pressure system which is involved by using the rotational formulation of the nonlinearity and the vorticity equation for the g-Navier-Stokes equations. In this study we particularly interest the following velocity-vorticity system of the g-Navier-Stokes equations over the two dimensional periodic box $\Omega = [0,1]^2 \subset \mathbb{R}^2$:

$$\begin{aligned} \frac{\partial u}{\partial t} - \nu \Delta_g u + \nu \frac{1}{g} (\nabla g \cdot \nabla) u + w \times u + \nabla P &= f, \\ \frac{\partial w}{\partial t} - \nu \Delta_g w + \nu \frac{1}{g} (\nabla g \cdot \nabla) w + (u \cdot \nabla) w &= \nabla \times f + w \left(\frac{\nabla g}{g} \cdot u \right), \\ \nabla \cdot (gu) &= 0 \quad \nabla \cdot (gw) = 0, \\ u(x, 0) &= u_0 \quad w(x, 0) = w_0 \end{aligned}$$

where $P = p + \frac{1}{2} |u|^2$, p denotes pressure, u represent velocity, w which play the role of vorticity, f is the body (external) forcing on the fluid and $\nu > 0$ kinematic viscosity. We consider this problem under the periodic boundary conditions. We assume u, p and w and the first derivative of u, w to be spatially periodic, i.e.,

$$u(x_1 + 1, x_2) = u(x_1, x_2) = u(x_1, x_2 + 1) \quad (x_1, x_2) \in \mathbb{R}^2.$$

We investigate global existence and uniqueness of the inviscid velocity-vorticity model of the g-Navier-Stokes equations using the classical Picard iteration method. The inviscid velocity-vorticity model of the g-Navier-Stokes is equivalent to the functional differential equation

$$\begin{aligned} \frac{du}{dt} + P_g(w \times u) &= P_g f, \\ \frac{dw}{dt} + B_g(u, w) &= P_g(\nabla \times f) + P_g \left(w \left(\frac{\nabla g}{g} \cdot u \right) \right), \end{aligned}$$

$$\nabla \cdot (gu) = 0 \quad \nabla \cdot (gw) = 0,$$

$$u(x, 0) = u_0 \quad w(x, 0) = w_0$$

where, for simplicity, we assume f to be time independent. We rewrite $B_g = P_g((u \cdot \nabla)w)$ and $P_g: L^2_{\text{ver}}(\Omega, g) \rightarrow H_g(\Omega)$ Helmholtz-Leray orthogonal projection.

Inviscid forms of the models in computational fluid dynamics have been attracted and extensively studied by many researchers.

Key Words: Existence and uniqueness, g-Navier-Stokes equations, inviscid velocity-vorticity model.

REFERENCES

- [1] M., Akbas, L. G. Rebholz, and C., Zervas, Optimal vorticity accuracy in an efficient velocity-vorticity method for the 2D Navier-Stokes equations. *Calcolo*. 55 (2018), no. 1, 1-29.
- [2] Y. Cao, E. M. Lunasin and E. S. Titi, Global well-posedness of the three-dimensional viscous and inviscid simplified Bardina turbulence models. *Communications in Mathematical Sciences*, 4(4) (2006), 823-848.
- [3] T. B. Gatski, Review of incompressible fluid flow computations using the vorticity-velocity formulation, *Applied Numerical Mathematics*. 7 (1991), no. 3, 227-239.
- [4] T. Heister, M. A. Olshanskii, and L. G. Rebholz, Unconditional long-time stability of a velocity- vorticity method for the 2D Navier-Stokes equations, *Numer. Math.* 135 (2017), no. 1, 143-167.
- [5] A. Larios, Y. Pei, and L. Rebholz, Global well-posedness of the velocity-vorticity-Voigt model of the 3D Navier-Stokes equations, *J. Differ. Equ.* 266 (2019), no. 5, 2435-2465.
- [6] A. J., Majda, A. L., Bertozzi, and A. Ogawa, Vorticity and incompressible flow, Cambridge texts in applied mathematics. *Appl. Mech. Rev.*, 55 (2002), no. 4, 229-247.
- [7] J., Roh, g-Navier Stokes Equations PhD Thesis, University of Minnesota, (2001).
- [8] J. Wu, Viscous and inviscid magneto-hydrodynamics equations. *Journal d'analyse Mathématique*, 73(1) (1997), 251-265.

Hermite-Hadamard -Fejér type inequalities for higher order convex functions and quadrature formulae

Josipa Barić ¹, Ljiljanka Kvesić ², Josip Pečarić ³ and Mihaela Ribičić Penava ⁴

1 Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, University of Split, Ruđera Boškovića 32, 21 000 Split, Croatia, jbaric@fesb.hr

2 Faculty of Science and Education, University of Mostar, Matice hrvatske bb, 88000 Mostar, Bosnia and Herzegovina, ljiljanka.kvesic@fpmoz.sum.ba

3 RUDN University, Miklukho-Maklaya str.6, 117198, Moscow, Russia, pecaric@element.hr

4 Department of Mathematics, Josip Juraj Strossmayer University of Osijek, Trg Ljudevita Gaja 6, 31000 Osijek, Croatia, mihaela@mathos.hr

ABSTRACT

The Hermite-Hadamard-Fejér inequalities state:

$$f\left(\frac{a+b}{2}\right) \int_a^b p(x) dx \leq \int_a^b p(x) f(x) dx \leq \frac{f(a) + f(b)}{2} \int_a^b p(x) dx,$$

where $p: [a, b] \rightarrow \mathbb{R}$ is nonnegative, integrable and symmetric about $\frac{a+b}{2}$ and $f: [a, b] \rightarrow \mathbb{R}$ is a convex function. In a special case, for $p(x) \equiv 1$ above inequalities reduce to the Hermite-Hadamard inequalities.

Over the last decades, Hermite-Hadamard type inequalities and Hermite-Hadamard-Fejér type inequalities have been extensively investigated and applied in approximation theory, special means theory, optimisation theory, information theory and numerical analysis.

The main purpose of this talk is to present some new Hermite-Hadamard-Fejér type inequalities for higher order convex functions and the general weighted integral formula involving w-harmonic sequences of functions. Since, a special case of the obtained integral formula is the general weighted Euler integral identity, we also derive Hermite-Hadamard-Fejér type inequalities for the general three, four and five point quadrature formulae of Euler type. Further, we establish Hermite-Hadamard-Fejér type inequalities for the corrected three point formula, the corrected four point formula and the corrected five point quadrature formula of Euler type. These formulae include not only the values of the function in the chosen points, but also the values of the first derivative at the end points of the interval, so have a degree of exactness higher than the adjoint original formulae.

Finally, certain special cases will be presented such as Hermite-Hadamard-Fejér type estimates for Simpson, Maclaurin, corrected Simpson and corrected Maclaurin quadrature rules.

Key Words: Hermite-Hadamard-Fejér inequalities, higher order convex functions, quadrature formulae.

REFERENCES

- [1] J. Barić, Lj. Kvesić, J. Pečarić and M. Ribičić Penava, Fejér type inequalities for higher order convex functions and quadrature formulae, *Aequationes Mathematicae*, (2021), accepted for publication
- [2] S.S. Dragomir and C.E.M. Pearce, *Selected Topics on Hermite–Hadamard Inequalities and Applications*. RGMIA Monographs. Victoria University, Melbourne, 2000.
- [3] I. Franjić and J. Pečarić, Hermite-Hadamard type inequalities for higher order convex functions and various quadrature rules. *Acta Math. Vietnam.* 37 (2012), 109-120.
- [4] S. Kovač and J. Pečarić, Weighted version of general integral formula of Euler type. *Math. Inequal. Appl.* 13 (2010), 579-599.
- [5] M. Klaričić Bakula, J. Pečarić and J. Perić, Extensions of the Hermite-Hadamard inequality with applications. *Math. Inequal. Appl.* 15 (2012), 899–921.

IMPLICIT EXPONENTIAL FINITE DIFFERENCE METHOD FOR NUMERICAL SOLUTIONS OF MODIFIED BURGERS EQUATION

Gonca Çelikten

Department of Mathematics, Faculty of Science and Letters, Kafkas University, Kars, Turkey ,
gnc.cnbk@gmail.com

ABSTRACT

In this paper, implicit exponential finite difference scheme has been used to acquire the numerical solutions of modified Burgers equation. A model problem is used to test the performance of the method. Comparisons of the exact solutions and the numerical solutions acquired by several other methods are present with tables. L_2 and L_∞ error norms are used to demonstrate the correctness of the results. The method is analyzed by von Neumann stability analysis method and it is displayed that the method is unconditionally stable.

Key Words: Modified Burgers equation, implicit exponential finite difference method, von Neumann stability analysis.

REFERENCES

- [1] M. A. Ramadan and T. S. El-Danaf, Numerical treatment for the modified Burgers equation, *Mathematics and Computers in Simulation*, 70 (2005), 90-98.
- [2] A. G. Bratsos and L. A. Petrakis, An explicit numerical scheme for the modified Burgers' equation, *International Journal for Numerical Methods in Biomedical Engineering*, 27 (2011), 232-237.
- [3] A. G. Bratsos, A fourth-order numerical scheme for solving the modified Burgers equation, *Computers and Mathematics with Applications*, 60 (2010), 1393-1400.
- [4] D. Irk, Sextic B-spline collocation method for the modified Burgers' equation, *Kybernetes*, 38 (2009), 1599-1620.
- [5] B. Saka, I. Dag, A numerical study of the Burgers' equation, *Journal of the Franklin Institute*, 345 (2008), 328-348.

Modified Finite Difference Method for solution of two-interval boundary value problems with transition conditions

Semih ÇAVUŞOĞLU¹, Oktay Sh. MUKHTAROV ^{2,3}

1 Institute of Graduate Studies, Tokat Gaziosmanpasa University, Tokat, Turkey,
semihcavusoglu@gmail.com

2 Department of Mathematics, Faculty of Science, Tokat Gaziosmanpasa University, Tokat, Turkey

3 Institute of Mathematics and Mechanics, Azerbaijan National Academy of Sciences, Baku, Azerbaijan,

omukhtarov@yahoo.com

ABSTRACT

A lot of mechanical and physical systems are modeled by ordinary or partial differential equations, for which exact solutions are impossible to find. Many researchers have tried to do this in various semi-analytical, numerical and approximate methods, such as the Finite Element Methods [1], the Adomian Decomposition Methods, the Differential Transform Method [2], the Runge-Kutta Method, the Explicit Euler Method, the Taylor's Expansion Method [3] etc. One of them is the Finite Difference Method [4], which is the simplest scheme and plays an important role in mathematical physics and engineering in all those cases where the differential equation can't be solved by analytical methods.

In this study, we have proposed a new modification of classical Finite Difference Method (FDM) for the solution of boundary value problems which are defined on two disjoint intervals and involved additional transition conditions at an common end of these intervals. The proposed modification of FDM differs from the classical FDM in calculating the iterative terms of numerical solutions. To illustrate the efficiency and reliability of the proposed modification of FDM some examples are solved. The obtained results are compared with those obtained by the standart FDM and by the analytical method. Corresponding graphical illustration are also presented.

Key Words: Finite Difference Method, transition conditions, boundary value problems.

REFERENCES

- [1] A.E. Omari, M. Chourak, S.E. Cherif, C.N. Ugena, M. Rougui and A. Chaaraoui, Numerical modeling of twin tunnels under seismic loading using the Finite Difference Method and Finite Element Method, *Materials Today: Proceedings*, 45 (2021), 45, 7566-7570.
- [2] I. Beroš, N. Hlupic, and D. Basch, Modification of the finite-difference method for solving a special class of nonlinear two-point boundary value problems, *Computer Science*, 16(1) (2021), 487-502.
- [3] P. K. Pandey, Nonstandard finite difference method for the approximate solution of two-point fourth order boundary value problems in ODEs, *Applied Sciences*, 23 (2021), 100-111.
- [4] S. Cavusoglu, O. Sh. Mukhtarov, A new finite difference method for computing approximate solutions of boundary value problems including transition conditions, *Bulletin of the Karaganda university Mathematics series*, 2(102) (2021), DOI 10.31489/2021M2/54-61

New Fractional Operators Including Wright Function in Their Kernels

Enes ATA¹, İ. Onur KIYMAZ²

1 Dept. of Mathematics, Faculty of Arts and Science, University of Ahi Evran, Kırşehir, Turkey,
enesata.tr@gmail.com

2 Dept. of Mathematics, Faculty of Arts and Science, University of Ahi Evran, Kırşehir, Turkey,
iokiymaz@ahievran.edu.tr

ABSTRACT

Fractional Calculus is a field of mathematics which deals with the definitions, properties and various applications of integrals and derivatives of arbitrary order. It started with G. W. Leibniz and has been developed up to nowadays by many mathematicians including P. S. Laplace, J. B. J. Fourier, N. H. Abel, J. Liouville, B. Riemann, A. K. Grünwald, A. V. Letnikov, J. Hadamard, A. Erdelyi and H. Kober. Fractional Calculus has been applied in various areas of applied mathematics, physics, engineering and finance.

Very recently, different types of functions have been used by many researchers in the kernels of integral operators for obtaining new definitions of fractional operators. Some of these operators are known as Caputo-Fabrizio, Losada-Nieto, Atangana-Baleanu and Gomez Aguilar-Atangana.

Motivated by these studies, we first defined two new fractional operators with a Wright function in their kernels. The first fractional operator is like Riemann-Liouville type, and the second one is like Caputo type. Then we obtained their Laplace and inverse Laplace transforms. We also determined fractional derivatives of some elementary functions and examined the relationships of new fractional operators with the other known fractional operators in the literature. Finally, as an example, we solved two differential equations with fractional derivatives obtained using both new fractional derivative operators.

Key Words: Gamma function, Wright function, Laplace transform, Inverse Laplace transform, Riemann-Liouville fractional operator, Caputo fractional operator.

REFERENCES

- [1] A. Atangana and D. Baleanu, New fractional derivatives with non-local and non-singular kernel: Theory and application to heat transfer model, *Thermal Science*, 20 (2) (2016), 763–769.
- [2] E. Ata and İ.O. Kıymaz, A study on certain properties of generalized special functions defined by Fox-Wright function, *Applied Mathematics and Nonlinear Sciences*, 5 (1) (2020), 147-162.
- [3] E. Ata, Generalized beta function defined by Wright function, preprint, (2021), arXiv: 1803.03121v3 [math.CA].
- [4] J. Losada and J.J. Nieto, Properties of a new fractional derivative without singular kernel, *Progr. Fract. Differ. Appl.*, 1 (2) (2015), 87–92.
- [5] J.F. Gomez-Aguilar and A. Atangana, New insight in fractional differentiation: Power, exponential decay and Mittag-Leffler laws and applications, *EPJ Plus*, 132 (13) (2017), 1–21.
- [6] L. Debnath, D. Bhatta, *Integral Transforms and Their Applications*, Third Edition, CRC Press, Boca Raton, London, New York, 2015.
- [7] M. Caputo and M. Fabrizio, A new definition of fractional derivative without singular kernel, *Progr. Fract. Differ. Appl.*, 1 (2) (2015), 73–85.
- [8] X.J. Yang, H.M. Srivastava and A.T. Macchado, A new fractional derivative without singular kernel, *Thermal Science*, 20 (2) (2016), 753–756.

Nonexistence of Global Solutions for a Hyperbolic-Type Wave Equation With Delay

Hazal Yüksekaya¹ and Erhan Pişkin²

¹ Department of mathematics, Dicle University, hazally.kaya@gmail.com

² Department of mathematics, Dicle University, episkin@dicle.edu.tr

ABSTRACT

In this paper, we consider a hyperbolic-type wave equation with delay. Under suitable conditions, the nonexistence of global solutions have been discussed. Generally, time delay effects arise in many applications and practical problems such as physical, chemical, biological, thermal and economic phenomena. Also, delay effects can be a source of instability. It is well known that delay effects may destroy the stabilizing properties of a well-behaved system. There are several examples that illustrate how time delays destabilize some internal or boundary control system.

Key Words: Nonexistence of solutions, hyperbolic-type, wave equation, delay.

REFERENCES

- [1] M. Kafini and S.A. Messaoudi, Local existence and blow-up of positive-initial-energy solutions of a nonlinear wave equation with delay, *Nonlinear Stud.* 27(3) (2020), 865-877.
- [2] S. Nicaise and C. Pignotti, Stabilization of the wave equation with boundary or internal distributed delay, *Differential Integral Equations.* 21 (2008), 935-958.
- [3] S. Nicaise and C. Pignotti, Stability and instability results of the wave equation with a delay term in the boundary or internal feedbacks, *SIAM J. Control Optim.* 45(5) (2006), 1561-1585.

Numerical Analysis of Singularly Perturbed Problems with Integral Boundary Conditions

Derya Arslan¹, Ömer Akın²

1 Bitlis Eren University, Department of Mathematics, 13200, Bitlis, Turkey.
ayredlanu@gmail.com

2 TOBB University, Department of Mathematics, Ankara, Turkey.
omerakin@gmail.com

ABSTRACT

In this research, exponential difference scheme is investigated to solve the singular perturbation problem with integral boundary condition. Some properties of the exact solution and its first-order derivative are obtained. A finite difference scheme is created on this mesh. The stability and convergence analysis of the method are shown as first-order convergent at the discrete maximum norm, independent of the perturbation parameter ε . Numerical datas are revealed by solving an example. These datas are demonstrated on table and figure.

The singularly perturbed problem is a problem involving a small parameter ε . So, there are thin transition layers where the solutions varies very rapidly for small values of ε , while away from layers it behaves regularly and varies slowly. Hence, the presence of small parameter in singularly perturbed problems gives severe difficulties that have to be addressed to ensure accurate numerical solutions [6-10]. To solve these type of problems, primarily there are some approaches namely, finite difference methods, finite element methods such as exponential elements, and fitted mesh methods which use or special piecewise uniform grids which condense in the boundary layers.

The solutions of like this problems include layers which occur in narrow layer regions of the domain. Singular perturbation problems emerge very frequently in fluid mechanics, fluid dynamics, quantum mechanics, elasticity, aerodynamics, meteorology, plasma dynamics, magnetohydrodynamics, rarefied gas dynamics, oceanography and other domains of the great world of fluid motion [1-5].

Key Words: Singular perturbation equation, finite difference scheme, uniform mesh, uniform convergence, integral conditions.

REFERENCES

- [1] R.E. O'Malley, Singular perturbation methods for ordinary differential equations, Springer Verlag, New York, 1991.
- [2] E.P. Doolan, J.J.H. Miller and W.H.A. Schilders, Uniform Numerical Method for Problems with Initial and Boundary Layers, Boole Press, 1980.

- [3] J.J.H. Miller, E. O'Riordan and G.I. Shishkin, Fitted Numerical Methods for Singular Perturbation Problems, World Scientific, Singapore, 1996.
- [4] P.A. Farrell, A.F. Hegarty, J.J.H. Miller, E. O'Riordan and G.I. Shishkin, Robust Computational Techniques for Boundary Layers, Chapman Hall/CRC, New York, 2000.
- [5] H.G. Roos, M. Stynes and L. Tobiska, Robust Numerical Methods Singularly Perturbed Differential Equations, Springer-Verlag, Berlin, 2008.
- [6] L. Bougoffa and A. Khanfer, Existence and uniqueness theorems of second-order equations with integral boundary conditions, 55 (2018), 899-911.
- [7] M. Cakir, A numerical study on the difference solution of singularly perturbed semilinear problem with integral boundary condition. Mathematical Modelling and Analysis, 21 (2016), 644-658.
- [8] D. Arslan, An approximate solution of linear singularly perturbed problem with nonlocal boundary condition, Journal of Mathematical Analysis, 11 (2020), 46-58.
- [9] D. Arslan, An effective numerical method for singularly perturbed nonlocal boundary value problem on Bakhvalov Mesh, Journal of Informatics and Mathematical Sciences, 11 (2019), 253-264.
- [10] R.A. Khan, The generalized method of quasilinearization and nonlinear boundary value problems with integral boundary conditions, Electronic Journal of Qualitative Theory of Differential Equations, 10 (2003), 1-9.

Numerical Integration Of System Of Differential Equations By Iterative Decreasing Dimension Algorithm

Zeliha ÖZDEMİR¹, Kemâl AYDIN²

1 Selcuk University, Graduate School of Natural Sciences, Konya, Turkey,
ozzdemirrzzeliha@gmail.com

2 Selcuk University, Faculty of Science, Department of Mathematics, Konya, Turkey,
kaydin@selcuk.edu.tr

ABSTRACT:

Studying on solution of the system of linear algebraic equations

$$Ax = f \quad (1)$$

is a classical problem which is important not only in linear algebra but also in other branches of science, engineering, economics. In (Wang and Jiang, 2000), a decreasing dimension method (DDM) has been proposed to solve the system (1) where A is $N \times N$ - regular matrix, x and f are N -vectors. In (Zhang, 2002), it has been said that the proposed DDM in (Wang and Jiang, 2000) is same as the well known domain decomposition technique based on a Schur complement type method. So in (Keskin and Aydın, 2007), the authors improved DDM and gave iterative decreasing dimension method (IDDM) which decreases the dimension of the linear systems, one order in every step without any pre-process.

On the other hand; we consider a system of ordinary differential equations

$$\frac{dx}{dt} = \kappa(t, x), \quad t \in (t^0, T], \quad (2)$$

where $\kappa(\cdot, \cdot)$ is continuous with respect to the first argument and Lipschitz continuous with respect to the second argument. The system (2) were converted to type (1) system by Euler's method and Newton's method under suitable conditions (initial value or boundary value conditions), respectively. Thus, the numerical integration of the type (2) system was converted to the type (1) system and investigated by Schur's complement method (Guibert and Tromeur-Dervout, 2008; Burden and Faires, 2010; Falgout et al., 2014; Badia and Olm, 2017).

In this study, the numerical integration of the system type (2) was investigated with modified algorithms by IDDM based on Schur's complement method. Numerical examples are also given using these algorithms

Key Words: Differential equations system, Schur complement method, Iterative decreasing dimension method.

REFERENCES

- [1] H. Wang and J. Jiang, Solution of the system of linear algebraic equations by decreasing dimension, *Appl. Math. Comp.*, 109(1) (2000), 51-57.
- [2] J. Zhang, Comments on "Solution of the system of linear algebraic equations by decreasing dimension". *Appl. Math. Comp.*, 128(1) (2002), 95-98.
- [3] T. Keskin and K. Aydın, Iterative decreasing dimension algorithm, *Comput. Math. Appl.*, 5 (2007), 1153-1158.
- [4] S. Badia and M. Olm, Nonlinear parallel-in-time Schur complement solvers for ordinary differential equations, *J. Comput. Appl. Math.*, 344 (2018), 794-806.
- [5] R. D. Falgout, S. Friedhoff, Tz. V. Kolev, S. P. MacLachlan and J. B. Schroder, Parallel time integration with multigrid. *SIAM J. Sci. Comput.*, 36(6) (2014), 635-661.
- [6] D. Guibert and D. Tromeur-Dervout, A Schur complement method for DAE/ODE systems in multi-domain mechanical design, In book: *Domain Decomposition Methods in Science and Engineering XVII*, 2008, 535-541.
- [7] R. L. Burden and J. D. Faires, *Numerical Analysis*, Ninth Edition, Brooks/Cole, 2010.

On Controlling Chaos for Multi-Step Iteration Process and Its Special Iterations in Discrete Dynamical Systems

Derya Sekman¹ and Vatan Karakaya²

¹ Dept. of Mathematics, Ahi Evran University, Kirsehir, Turkey,
deryasekman@gmail.com

² Dept. of Mathematical Engineering, Yildiz Technical University, Istanbul, Turkey,
vkkaya@yahoo.com

ABSTRACT

Due to the close relationship between chaos and dynamical systems, it is a natural result to consider the iteration processes that are related to the dynamical systems of fixed point theory. From this natural relationship, the control of the chaos that occurs in fixed point iteration dynamics is the main problem of the article. To achieve this goal, analytical solutions have been given to control the chaos that occurs for unstable fixed points of multi-step iteration process. After, it has been shown that there is an effective regime for the parameters of multi-step iteration. It is well known that it occurs the chaos in the any system according to parameters in system. To illustrate this claim, well-known special cases of multi-step iteration process, Noor iteration, Ishikawa iteration, Mann iteration, Krasnoselskij iteration, Picard iteration processes have been introduced. In particular, among these iterations, the Noor iteration process has been studied in detail in terms of controlled chaos. The Lyapunov exponent was used to estimate the stability and instability of fixed points and periods that cause chaos in iteration processes. Finally, with the help of MATLAB program, all these results are shown on logistic system and cubic equation with chaos properties.

This work was supported by the Yildiz Technical University Scientific Research Projects Coordination Unit. Project Number: FDK-2018-3348.

Key Words: Controlling chaos, multi-step iteration process, dynamical systems.

REFERENCES

- [1] Ashish, J. Cao and R. Chugh, Chaotic behavior of logistic map in superior orbit and an improved chaos based traffic control model, Nonlinear Dyn. 94 (2018), 959-975.

- [2] V. Berinde, Controlling chaotic dynamical systems through fixed point iterative techniques, "Vasile Alecsandri" University of Bacau Faculty of Sciences Scientific Studies and Research Series Mathematics and Informatics 19(2) (2009), 47-58.
- [3] V. Berinde, Iterative Approximation of Fixed Point, Springer Verlag, New York, 2007.
- [4] F. Gürsoy, V. Karakaya and B.E. Rhoades, Data dependence results of new multistep and S-iterative schemes for contractive-like operators, Fixed Point Theory and Applications, 76 (2013), 1-12.
- [5] F. Gürsoy, V. Karakaya and B.E. Rhoades, Some convergence and stability results for the Kirk Multistep and Kirk-SP fixed point iterative algorithms, Abstract and Applied Analysis, Article ID 806537 (2014), 1-12.
- [6] W. Huang, Controlling chaos through growth rate adjustment, Discrete Dynamics in Nature and Society 7(3) (2002), 191-199.

On Some Applications of Korovkin Type Theorem for k-Positive Linear Operators

Nida PALAMUT KOŞAR¹,

¹ Department of Mathematics and Science Education, Nizip Education Faculty,
Gaziantep University, Turkey, npkosar@gmail.com, npkosar@gantep.edu.tr

ABSTRACT

During the last forty years, many researchers impressed with the Korovkin theorems and they applied these theorems to the setting of different function spaces like Banach algebras, Banach spaces and so on.

The study for the statistical convergence plays very important role as an expanding of the basic convergence for the sequence spaces. The concept of statistical convergence was introduced by Fast [1] and Steinhaus [2], later reintroduced by Schoenberg [3].

The first work for the convergence of analytic functions of complex variables by the idea of k-positive linear operators was given by Gadjiev [4]. He offered “k-positive operators” and applied the Korovkin type theorem for these operators in the space of holomorphic functions on the different domains.

Let us

$$D_r^R := D(r; R) := \{z \in \mathbb{C}^n : r_j < |z_j| < R_j; j = \overline{1, n}\}$$

be polyradius in \mathbb{C}^n and $\mathcal{H}(D(r; R))$ be holomorphic functions space over $D(r; R)$ with the topologically compact convergence. In this study, the concept of convergence will have the same meaning as uniform convergence on compact subsets of the polyradius $D(r; R)$. We will take into account the semi-norms for $r < r' < R' < R$,

$$\|f\|_{\mathcal{H}(D(r; R)), r', R'} := \|f\|_{\mathcal{H}(D(r; R))} := \max_{r' < |z| < R'} |f(z)|$$

that convert $\mathcal{H}(D(r; R))$ into a Frechet-type space.

In this study we will be interested in the sequence of k-positive linear operators that stands for the Laurent coefficient. The aim of this research is to attain Korovkin-type theorems for k-positive linear operators on $\mathcal{H}(D(r; R))$.

Key Words: Korovkin type theorem, k -positive linear operators, statistical convergence, holomorphic functions in space of several complex variables.

REFERENCES

- [1] H. Fast, Sur la convergence statistique, Colloq. Math. 2 (1951), 241-244.
- [2] H. Steinhaus, Sur la convergence ordinaire et la convergence asymptotique, Colloq. Math. 2 (1951) 73-74.
- [3] I.J. Schoenberg, The integrability of certain functions and related summability methods, Amer. Math. Monthly 66 (1959), 361-375.
- [4] AD. Gadjiev, Linear k -positive operators in a space of regular functions and theorems of P.P. Korovkin type, Akad Nauk Azerb SSR Ser Fiz Tekh Mat Nauk, 5 (1974), 49-53 (in Russian).

On stability of the bacterial infection model through incommensurate fractional-order system

Bahatdin DAŞBAŞI¹

¹ Kayseri University, Faculty of Engineering, Architecture and Design, Department of Engineering Basic Sciences, 38280, Kayseri/Turkey

ABSTRACT

In this study, the time-dependent changes in antibiotic concentrations and the bacterial population causing the infection in a host receiving multiple antibiotic therapy in case of a bacterial infection were investigated by mathematically modeling with incommensurate fractional-order system (IFOS).

Let us consider that B represent the bacterial concentration in an individual at time t , and C_i for $i = 1, 2, \dots, n$ represent the i -th antibiotic concentration at time t in an individual receiving multiple antibiotic therapy against the infectious bacteria. The model for $t > t_0$ is

$$\begin{aligned} D^{\alpha_1} B(t) &= \beta B \left(1 - \frac{B}{\delta} \right) - B \left[\sum_{i=1}^n q_i C_i \right] - \eta B \\ D^{\alpha_2} C_1(t) &= \Lambda_1 - \mu_1 C_1 \\ &\vdots \\ D^{\alpha_{n+1}} C_n(t) &= \Lambda_n - \mu_n C_n \end{aligned} \quad (1)$$

where initial conditions are $B(0) = B_0$, $C_1(0) = C_{10}, \dots, C_n(0) = C_{n0}$. In addition, for the numbers $\alpha_1, \alpha_2, \dots, \alpha_{n+1}$, which are derivative orders in the model, it is satisfied the followings:

$$0 < \alpha_1, \alpha_2, \dots, \alpha_{n+1} \leq 1 \quad (2)$$

such that

$$k_1, k_2, \dots, k_n, m_1, m_2, \dots, m_n \in \mathbb{Z}^+ \quad (3)$$

and

$$\alpha_1 = \frac{k_1}{m_1}, \alpha_2 = \frac{k_2}{m_2}, \dots, \alpha_n = \frac{k_n}{m_n} \quad (4)$$

where the number m is the smallest of the common multiples of m_1, m_2, \dots, m_n . The parameters used in the model can be summarized as follows. The bacteria causing the infection increase according to the logistic rules, where β is the growth rate of the bacteria and δ is the carrying capacity of the bacteria. In addition, η indicates the natural death rate of the bacterium and q_i for $i = 1, 2, \dots, n$ indicates the death rate of the bacteria due to the i -th antibiotic. For $i = 1, 2, \dots, n$, the i -th antibiotic is taken in a fixed amount of Λ_i and is excreted from the body at a rate of μ_i . Also, we have the following inequalities.

$$\beta, \delta, \eta, q_i, \Lambda_i, \mu_i > 0 \quad (5)$$

In addition, the results, which are compatible with real situations, have obtained in numerical studies.

Key Words: Incommensurate Fractional-Order Differential Equation; Bacterial Infection Modeling; Stability Analysis.

REFERENCES

- [1] B. Daşbaşı, "Stability Analysis of Mathematical Model including Pathogen-Specific Immune System Response with Fractional-Order Differential Equations," Computational and Mathematical Methods in Medicine, no. 7930603, (2018), pp. 1-10.
- [2] B. Daşbaşı, "Stability analysis of the hiv model through incommensurate fractional-order nonlinear system," Chaos, Solitons & Fractals, vol. 137, no. 109870, (2020), pp. 1-12.
- [3] B. Daşbaşı and İ. Öztürk, "Mathematical modelling of bacterial resistance to multiple antibiotics and immune system response," SpringerPlus, vol. 5, no. 408 (2016), pp. 1-17.
- [4] E. I. Mondragón et al., "Mathematical modeling on bacterial resistance to multiple antibiotics caused by spontaneous mutations," BioSystems, vol. 117, (2014), pp. 60–67.
- [5] S.G. Gal, Shape-Preserving Approximation by Real and Complex Polynomials, Birkhäuser Boston, Inc., Boston, MA, 2008.

Parametric Identification in ODE Models as a Tool for Solving Real-world Problems

Galina Lyutskanova-Zhekova^{1,2}

1 Faculty of Mathematics and Informatics, University of Sofia, 5 J. Bourchier Ave., 1164 Sofia, Bulgaria, g.zhekova@fmi.uni-sofia.bg

2 Institute of Mathematics and Informatics, Bulgarian Academy of Science, Acad. G. Bonchev Str., Block 8, 1113 Sofia, Bulgaria, g.zhekova@math.bas.bg

ABSTRACT

One of the best ways to understand the deeper connections between the different aspects of mathematics to start working on interesting real-world problem, which has a great number of applications. Engaging students in such a way helps them to understand the material on a deeper level and apply it to solving real-world problems. In the current talk, I shall present two real-world problems which could be solved using parametric identification (a method to estimate the unknown parameters of a model). The problems are suitable for students at the University level.

Compute the transmission rate of an infectious disease. There is a given data set, which consists of data for the number of susceptible and infectious individuals to a certain disease. The aim is to find transmission rate of the disease. Firstly, students shall get familiar with the SIR model, which splits the population to three groups – susceptible, infectious, and removed [1]. The model consists of three ordinary differential equations for the three different groups and appropriate initial conditions are applied. It has two unknown parameters – the recovery rate and the transmission rate. Secondly, the problem shall be solved with fixed parameters. Then the unknown parameters are found in such a way that the model fits the data best.

Axisymmetric drop shape analysis. The problem can be formulated as follows. Using an image of a pendant drop of unknown liquid, find the surface tension of a liquid. This problem has a great number of applications – in food industry, oil industry, etc. Firstly, the students shall get familiar with the Young-Laplace equation of capillarity, which gives a relationship between the surface tension and the mean curvature of the drop and the pressure, applied to it.

Computing the curvature of the drop and using some geometrical considerations, they shall arrive at mathematical model, consisting of three ODEs with two unknown parameters [2]. In order to find the unknown parameters, the profile of the drop shall be extracted and the unknown parameters shall be computed, using parametric identification. Detailed description of all experiments from student point of view could be found in [3].

Key Words: SIR model, Young-Laplace equation of capillarity, mathematics education.

REFERENCES

- [1] W.O. Kermack and A.G. McKendrick, A contribution to the mathematical theory of epidemics, Proc. R. Soc. London 115 (1927), 700
- [2] Y. Rotenberg, L. Boruvka and A. W. Neumann, Determination of surface tension and contact angle from the shapes of axisymmetric fluid interfaces, J Colloid Interface Sci 93 (1983), 169-183.
- [3] G. Lyutskanova, K. Mihaylov, V. Kolev, Axisymmetric Drop Shape Analysis in Proceedings of Preparatory Modelling Week (Sofia, Bulgaria, 2015)
http://pmw2015.fmi.uni-sofia.bg/Documents/Problem_2_Report.pdf.

Several Weighted Hardy type Inequalities with Robin Boundary Conditions

Abdullah Yener

Department of Mathematics, Istanbul Commerce University, ayener@ticaret.edu.tr

ABSTRACT

The classical Hardy inequality on \mathbb{R}^n says, for $1 < p < \infty$,

$$\int_{\mathbb{R}^n} |\nabla \phi(x)|^p dx \geq \int_{\mathbb{R}^n} \frac{|\phi(x)|^p}{|x|^p} dx;$$

the constant is best possible and there are no nonzero extremals. Even when $p = n$; the inequality is interesting because of the optimality of the constant. For $p < n$ one assumes $\phi \in C_0^\infty(\mathbb{R}^n)$, while $p > n$ one assumes $\phi \in C_0^\infty(\mathbb{R}^n \setminus \{0\})$. Here the subscript zero signifies compact support, ∇ is the usual Euclidean gradient, and $|x| = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2}$.

Hardy-type inequalities have evoked the interest of many mathematicians because of their applications to elliptic and parabolic PDE's motivated by physics and geometry, and the literature concerning different versions of Hardy inequalities and their applications is extensive on the Euclidean setting.

On the other hand, very little work has been done on Hardy type inequalities with

Robin boundary conditions. However, there has been some initiation in this area of interest. In an interesting paper, H. Kovarik and A. Laptev [3] proved, among other results, a Hardy inequality for Laplace operators with Robin boundary conditions. For convex domains they also showed how the corresponding Hardy weight depends on the coefficient of the Robin boundary conditions. Later, T. Ekholm, H. Kovarik and A. Laptev [2] studied the best constant in a Hardy inequality for the p-Laplace operator on convex domains with Robin boundary conditions.

The main objective of this article is to study the general weighted Hardy type inequalities for the p-Laplace operators with Robin boundary conditions. We should emphasize that our unifying method is quite practical and constructive to obtain several weighted Hardy, Maz'ya and Heisenberg-Pauli-Weyl type inequalities with boundary terms.

Key Words: Hardy inequality, Boundary term, Robin boundary condition.

REFERENCES

- [1] E.B. Davies, A review of Hardy inequalities, in: The Maz'ya Anniversary Collection, vol. 2, in: Oper. Theory Adv. Appl., vol. 110, Birkhauser, Basel, 1999.

- [2]** T. Ekholm, H. Kovarik and A. Laptev, Hardy inequalities for p -Laplacians with Robin boundary conditions, *Nonlinear Anal.* 128 (2015), 365-379.
- [3]** H. Kovarik and A. Laptev, Hardy inequalities for Robin Laplacians, *J. Funct. Anal.* 262 (2012), 4972-4985.
- [4]** V.G. Maz'ya, *Sobolev Spaces*, Springer, Berlin, 1985.

Some Further Properties of BLUPs in SUR Models

Melek Eriş Büyükkaya¹, Nesrin Güler² and Melike Yiğit³

1 Department of Statistics and Computer Sciences, Karadeniz Technical University, Trabzon, Turkey, melekeris@ktu.edu.tr

2 Department of Econometrics, Sakarya University, Sakarya, Turkey, nesring@sakarya.edu.tr

3 Department of Mathematics, Sakarya University, Sakarya, Turkey, melikeyigitt@gmail.com

ABSTRACT

Seemingly unrelated regression (SUR) models, proposed by [1], are a class of multiple regression models. These models may seem unrelated but each of them can be linked to their correlated error terms across the models. In general, instead of considering the group of linear regression models individually, merging them into a combined model by making use of block matrices is a common procedure to gain more efficiency in the prediction of the unknown vectors.

Characterization of algebraic and statistical properties of predictors of the unknown vectors in regression models is one of the main subjects in statistical analysis. Furthermore, establishing the results on the connection among predictors in different models is one of the primary problems encountered in the theory of regression analysis. Therefore, it is natural to consider certain links among predictors under SUR models and their combined models. In this study, the best linear unbiased predictors (BLUPs) are considered in the context of SUR models. Some properties of BLUPs in SUR models are investigated. We present analytical expressions of the BLUPs and establish some properties of BLUPs by using block matrices' rank and inertia formulas. Our main purpose is to give a variety of equalities and inequalities of BLUPs and their covariance matrices for comparison of these predictors in considered models.

There are extensive works on SUR models, we can refer to [2,3] among others. SUR models are a special class of a system of linear regression models. We may refer to [4,5] for a system of linear regression models. Some results related to inertias, ranks, and Löwner partial ordering of symmetric matrices can be found in, e.g., [6]. Furthermore, we may refer to, e.g., [7,8] for rank and inertia formulas for covariance matrices of predictors/estimators.

Key Words: BLUP, inertia, rank, seemingly unrelated regression model.

REFERENCES

- [1] A. Zellner, An efficient method of estimating seemingly unrelated regressions and tests for aggregation bias, J. Am. Stat. Assoc., 57 (1962), 348-368.
- [2] J. Hou and Y. Zhao, Some remarks on a pair of seemingly unrelated regression models, Open Math., 17 (2019), 979-989.
- [3] L. Gong, Establishing equalities of OLSEs and BLUEs under seemingly unrelated regression models, J. Stat. Theory Pract., 13 (5) (2019).
- [4] L. G. Telser, Iterative estimation of a set of linear regression equations. J. Am. Stat. Assoc., 59 (1964), 845-862.
- [5] R. W. Parks, Efficient estimation of a system of regression equations when disturbances are both serially and contemporaneously correlated. J. Am. Stat. Assoc., 62 (1967), 500-509.
- [6] Y. Tian, Equalities and inequalities for inertias of Hermitian matrices with applications, Linear Algebra Appl., 433 (2010), 263-296.
- [7] Y. Tian, Some equalities and inequalities for covariance matrices of estimators under linear model, Stat. Pap., 58 (2) (2017), 467-484.
- [8] Y. Tian, Matrix rank and inertia formulas in the analysis of general linear models, Open Math., 15 (1) (2017), 126-150.

Some Remarks on Predictors in Reduced Linear Models

Melek Eriş Büyükkaya¹, Nesrin Güler² and Melike Yiğit³

1 Department of Statistics and Computer Sciences, Karadeniz Technical University, Trabzon, Turkey, melekeris@ktu.edu.tr

2 Department of Econometrics, Sakarya University, Sakarya, Turkey, nesring@sakarya.edu.tr

3 Department of Mathematics, Sakarya University, Sakarya, Turkey, melikeyigitt@gmail.com

ABSTRACT

Establishing a relationship between two or more different linear models in the theory of regression analysis is one of the classical research problems. One of these problems is to examine the relations between original linear model and its transformed models. Sometimes, linear models need to be transformed to meet the requirements of the analysis in statistical inferences of linear regression models. Using several linear transformations of the observable random vector, we can face different transformed models of the original model. One of these transformed models corresponds the reduced linear models of the original model.

In this study, we consider the best linear unbiased predictors (BLUPs) in the context of reduced linear models. Some properties of the BLUPs and their analytical expressions are investigated in reduced linear models. The results are established by making use of formulas of inertias and ranks of block matrices which are effective algebraic tools in matrix theory. Covariance matrices of BLUPs are usually used as comparison criteria to determine optimal predictors among other types of unbiased predictors because of their minimum covariance requirement in the Löwner partial ordering. The main purpose of the study is to derive some results on the comparison of covariance matrices of BLUPs under the original linear model and its reduced linear models. The subjects related to the results obtained in this study can also be found in [1-8].

Key Words: BLUP, covariance matrix, inertia, rank, reduced linear model.

REFERENCES

- [1] J. Groß and S. Puntanen, Estimation under a general partitioned linear model, *Linear Algebra Appl.*, 321 (2000), 131–144.
- [2] M. Liu, Y. Tian and R. Yuan, Statistical inference of a partitioned linear random-effects model, *Commun. Stat. - Theory Methods*, (2021), DOI: 10.1080/03610926.2021.1926509.
- [3] N. Güler and M. E. Büyükkaya, Rank and inertia formulas for covariance matrices of BLUPs in general linear mixed models, *Commun. Stat. - Theory Methods*, (2019), DOI: 10.1080/03610926.2019.1599950.
- [4] P. Bhimasankaram, and R. Saharay, On a partitioned linear model and some associated reduced models, *Linear Algebra Appl.*, 264 (1997) 329-39, DOI: 10.1016/S0024-3795(97)00011-6.
- [5] X. Ren, The equalities of estimations under a general partitioned linear model and its stochastically restricted model, *Commun. Stat. - Theory Methods*, 45 (22) (2016), 6495-6509, DOI: 10.1080/03610926.2014.960587.
- [6] Y. Tian, Some Decompositions of OLSEs and BLUEs Under a Partitioned Linear Model, *Int. Stat. Rev.*, 75 (2) (2007), 224–248.
- [7] Y. Tian and B. Jiang, Equalities for estimators of partial parameters under linear model with restrictions, *J. Multivar. Anal.*, 143 (2016), 299-313.
- [8] Y. Tian and X. Zhang, On connections among OLSEs and BLUEs of whole and partial parameters under a general linear model, *Stat. Probab. Lett.*, 112 (2016), 105-112.

Sub Pico-Second Optical Solitons in Birefringent Fibers with Kaup-Newell Equation by Two Strategic Integration Technologies

Yakup Yıldırım

Department of Mathematics, Faculty of Arts and Sciences, Near East University, 99138 Nicosia, Cyprus, yakup.yildirim@neu.edu.tr

ABSTRACT

This paper studies sub pico-second optical solitons in birefringent fibers. The term optical soliton is used to refer to any optical field that does not change during propagation because of a delicate balance between group velocity dispersion and linear effects in the medium. The coupled vector form of Kaup-Newell equation without four-wave mixing (4WM) is considered. The governing model is a case of derivative nonlinear Schrödinger's equation and describes sub pico-second optical solitons in birefringent fibers. These optical solitons are electromagnetic pulses which are commonly referred to as ultrafast events. Optical bright, dark, singular and combo singular solitons are recovered by the Riccati function and sine-Gordon function methods. These solitons depend on certain parameter restrictions which are given in this paper. The Riccati function method gives rise to dark and singular solitons while the sine-Gordon function method causes to bright, dark, singular and combo singular solitons. The steps of the Riccati function and sine-Gordon function methods are introduced. These methods are based on the principle of homogeneous balance between the highest order derivatives and the nonlinear terms of nonlinear evolution equations in abundant scientific fields such as in fluid dynamics, nonlinear optical fibers, nuclear physics, electromagnetism, medicine, finance, mathematical biosciences, etc.

Key Words: Kaup--Newell equation, Riccati function method, sine-Gordon function method, optical solitons.

REFERENCES

- [1] A. Biswas, Y. Yıldırım, E. Yaşar, Q. Zhou, S. P. Moshokoa and M. Belic, Sub pico-second pulses in mono-mode optical fibers with Kaup-Newell equation by a couple of integration schemes, Optik 167 (2018), 121-128.
- [2] Y. Yıldırım, A. Biswas, Q. Zhou, A. S. Alshomrani and M. R. Belic, Sub pico-second optical pulses in birefringent fibers for Kaup-Newell equation with cutting--edge integration technologies, Results Phys. 102660 (2019).

THE ASYMPTOTIC BEHAVIOR FOR GENERALIZED λ -BERNSTEIN-STANCU TYPE OPERATORS

Qing-Bo Cai¹, Gülten Torun² and Ülkü Dinlemez Kantar³

¹ Fujian Provincial Key Laboratory of Data-Intensive Computing, Key Laboratory of Intelligent Computing and Information Processing, School of Mathematics and Computer Science, Quanzhou Normal University, Quanzhou 362000, CHINA, qbcail@126.com

² Kastamonu University, Faculty of Education, Mathematics and Science Education, Kastamonu, TURKEY, gtorun@kastamonu.edu.tr

³ Gazi University, Faculty of Science, Department of Mathematics, Ankara TURKEY, ulku@gazi.edu.tr

ABSTRACT

The Bernstein operators, which are positive linear operators, are of great importance for the theory of approximation. In [1], Bernstein operators were introduced by Bernstein to prove the Weierstrass approximation theorem. In [2], Cai et al. introduced the Bernstein operators with shape parameter $\lambda \in [-1, 1]$ as follows:

$$B_m^\lambda(g, x) = \sum_{j=0}^m g\left(\frac{j}{m}\right) \tilde{p}_{m,j}(\lambda, x), \quad m \in \mathbb{N}, x \in [0, 1], \quad (1)$$

where $\tilde{p}_{m,j}(\lambda, x)$ are Bézier basis functions with shape parameter λ defined by

$$\tilde{p}_{m,0}(\lambda, x) := p_{m,0}(x) - \frac{\lambda}{m+1} p_{m+1,1}(x)$$

$$\tilde{p}_{m,j}(\lambda, x) := p_{m,j}(x) - \lambda \left[\frac{m-2j+1}{m^2-1} p_{m+1,j}(x) - \frac{m-2j-1}{m^2-1} p_{m+1,j+1}(x) \right]; j = 1, 2, \dots, m-1 \quad (2)$$

$$\tilde{p}_{m,m}(\lambda, x) := p_{m,m}(x) - \frac{\lambda}{m+1} p_{m+1,m}(x),$$

where the Bernstein basis functions $p_{m,j}(x)$ are defined as

$$p_{m,j}(x) := \binom{m}{j} x^j (1-x)^{m-j}; \quad j = 0, 1, \dots, m.$$

In [3], Srivastava et al. constructed λ -Bernstein-Stancu operators defined by

$$B_{m,\alpha,\beta}^\lambda(g, x) = \sum_{j=0}^m g\left(\frac{j+\alpha}{m+\beta}\right) \tilde{p}_{m,j}(\lambda, x); \quad m \in \mathbb{N}, x \in [0, 1], \quad (3)$$

where α and β are positive real numbers satisfying $0 \leq \alpha \leq \beta$, and Bézier basis functions $\tilde{p}_{m,j}(\lambda, x)$ with shape parameter λ are defined in (2).

In the present work, we introduce the following generalized λ -Bernstein-Stancu operators with shifted knots for $g \in C[0,1]$,

$$G_{m,\lambda}^{\alpha,\beta}(g, x) = \left(\frac{m+\beta_2}{m} \right) \sum_{j=0}^m g \left(\frac{j+\alpha_1}{m+\beta_1} \right) \tilde{p}_{m,j}(\lambda, x); \quad m \in \mathbb{N}, \quad (4)$$

where α_i and β_i are positive real numbers satisfying $0 \leq \alpha_i \leq \beta_i$, for $i = 1, 2$ and $x \in \left[\frac{\alpha_2}{m+\beta_2}, \frac{m+\alpha_2}{m+\beta_2} \right]$, and Bézier basis functions $\tilde{p}_{m,j}(\lambda, x)$ with shape parameter $\lambda \in [-1, 1]$ are defined in (2) such that

$$p_{m,j}(x) := \binom{m}{j} \left(x - \frac{\alpha_2}{m+\beta_2} \right)^j \left(\frac{m+\alpha_2}{m+\beta_2} - x \right)^{m-j}; \quad j = 0, 1, \dots, m.$$

We give a Korovkin type approximation theorem and obtain the rate of convergence of these type operators for Lipschitz type functions. Then, we give a Voronovskaja type theorem for the asymptotic behavior for these operators. Finally, we give numerical examples and their graphs to demonstrate the convergence of $G_{m,\lambda}^{\alpha,\beta}(f, x)$ to $f(x)$ with respect to m values.

Key Words: λ -Bernstein-Stancu type operators, Bézier bases functions, Voronovskaja type theorem.

REFERENCES

- [1] S.N. Bernstein, Démonstration du théorème de Weierstrass fondée sur la calcul des probabilités, Commun. Soc. Math. Charkow Sér. 13(2) (1912), 1-2.
- [2] Q.B. Cai, B.Y. Lian and G. Zhou, Approximation properties of λ -Bernstein operators, J. Inequal. Appl. 2018 (2018), 61.
- [3] H.M. Srivastava, F. Özger and S.A. Mohiuddine, Construction of Stancu-Type Bernstein Operators Based on Bézier Bases with Shape Parameter λ , Symmetry 11(2019), 316.

The Marshall–Olkin transmuted Rayleigh distribution. Statistical properties and different methods of estimation

Faton Merovci ¹

¹ University of Mitrovica “Isa Boletini” Kosovo, faton.merovci@umib.net

ABSTRACT

In this paper we have introduced a new four parameter Rayleigh distribution called the Marshall Olkin transmuted Rayleigh distribution obtained by new family of continuous distributions called the Marshall Olkin transmuted G family of univariate distribution given by Afify et al [1]. Some of its mathematical properties including the quantile function, limit behavior, hazard rate function, ordinary and incomplete moments, characteristic function, mean residual lifetime, mean past lifetime, mean deviation about mean and median, order statistics, probability weighted moments, Shannon and Rényi entropies, mean deviations, Bonferroni and Lorenz curves and stress-strength parameter are derived.

The estimation of the model parameters is performed by different methods like the maximum likelihood method, moments estimators, least squares estimators, weighted least squares, maximum product spacing estimates, methods of minimum distances, method of Cramer- von-Misses, methods of Anderson-Darling and right-tail Anderson-Darling and compare them using extensive simulations studies. We investigate the mean estimates, biases, and root mean square errors using Monte Carlo simulations and we showed that the methods of Cramer- von-Misses has the smallest mean square errors and mean relative estimates, for parameters, proving to be the most efficient method compared to other methods.

Two real data set are using to illustrate its applicability, and it is shown that Marshall Olkin transmuted Rayleigh distribution fits much better than some other distributions.

Key Words: Marshal-Olkin transmuted-G; maximum likelihood estimation; Order Statistics.

REFERENCES

- [1] A.Z. Afify, H.M. Yousof, M. Alizadeh, I. Ghosh, S. Ray, and G. Ozel, G, The Marshall–Olkin Transmuted-G Family of Distributions. *Stochastics and Quality Control*, 35(2) (2020), 79-96.
- [2] F. Merovci, Transmuted Rayleigh distribution. *Austrian Journal of statistics*, 42(1) (2013), 21-31.

The Poisson Topp Leone generalized exponential distribution

Faton Merovci ¹

1 University of Mitrovica "Isa Boletini" Kosovo, faton.merovci@umib.net

ABSTRACT

Recently many authors constructed several classes of distributions by extending common families of continuous distributions. In 2020 Merovci, Yousouf and Hamedani proposed a new family of distributions called Poisson Topp Leone-G family of distributions. By using this generator, in this paper we have proposed a new extension of the generalized exponential distribution called Poisson Topp Leone generalized exponential distribution. We study the main properties of this new distribution including the quantile function, limit behavior, hazard rate function, ordinary and incomplete moments, characteristic function, order statistics, Shannon and Rényi entropies, mean deviations and Bonferroni and Lorenz curves.

Several methods are used for the estimation of the unknown parameters: the maximum likelihood method, moments estimators, least squares estimators, weighted least squares, maximum product spacing estimates, methods of minimum distances, method of Cramer- von-Misses and methods of Anderson-Darling. We investigate the mean estimates, biases, and root mean square errors using Monte Carlo simulations and we showed that the methods of Anderson-Darling has the smallest mean square errors and mean relative estimates, for parameters, proving to be the most efficient method compared to other methods.

We present two applications with real data set to illustrate its applicability, and it is shown that Poisson Topp Leone generalized exponential distribution fits much better than its parent distribution.

Key Words: exponential distribution; maximum likelihood method; Shannon and Rényi entropies.

REFERENCES

- [1] R.D. Gupta and D. Kundu, Theory & methods: Generalized exponential distributions. *Australian & New Zealand Journal of Statistics*, 41(2) (1999), 173-188.
- [2] F. Merovci, H. Yousof, and G.G. Hamedani, The Poisson Topp Leone generator of distributions for lifetime data: theory, characterizations and applications. *Pakistan Journal of Statistics and Operation Research*, 343-355, 2020.

GEOMETRY

An Extension of Smarandache Ruled Surfaces According to Frenet Frame in E^3

Davut CANLI¹, Süleyman ŞENYURT² and Kebire Hilal AYVACI³

¹ Ordu University, Faculty of Arts and Sciences, Depart. of Maths., davutcanli@odu.edu.tr

² Ordu University, Faculty of Arts and Sciences, Depart. of Maths., ssenyurt@odu.edu.tr

³ Ordu University, Faculty of Arts and Sciences, Depart. of Maths., kehirehilalayvaci@odu.edu.tr

ABSTRACT

In this paper, we address the new notion namely Smarandache ruled surfaces according to Frenet frame and as an extension we introduce new ruled surfaces where all possible linear combinations of the vector elements of Frenet frame are taken into account as a base curve and the generator. Next, by referring the coefficients of first and second fundamental forms, we calculate the mean and Gaussian curvatures so that we could provide some characteristics for these ruled surfaces. In accordance with the calculations, we give the necessary conditions for any surface defined to be developable and minimal, as well. Finally, we present examples for each created ruled surfaces and picture those by using Maple 17 program.

Key Words: Smarandache ruled surfaces, first and second fundamental forms, mean and Gaussian curvatures, developable surfaces, minimal surfaces.

REFERENCES

- [1] A. Sarioğlu and A. Tutar, On ruled surfaces in Euclidean space, Int. J. Contemp. Math. Sci., 2(1) (2007), 1-11.
- [2] A. Yılmaz and B. Şahin, On geodesics of the tangent and normal surfaces defined by TN-Smarandache curve according to Frenet frame. 16th International Geometry Symposium (pp.1-10). Manisa, Turkey, 2018.
- [3] A. Yılmaz and B. Şahin, On Ruled Surfaces Defined by Smarandache Curve. 2nd International Students Science Congress, İzmir, Turkey, 2018.
- [4] A. Yılmaz and B. Şahin, On Geodesics of the Binormal Surface Defined by Smarandache Curve. 3rd International Students Science Congress, İzmir, Turkey, 2019.
- [5] S. Ouarab, Smarandache Ruled Surfaces according to Frenet-Serret Frame of a Regular Curve in E^3 , Abstract and Applied Analysis, vol. (2021).

- [6] S. Ouarab, NC-Smarandache Ruled Surface and NW-Smarandache Ruled Surface according to Alternative Moving Frame in E^3 , Journal of Mathematics, vol. (2021).
- [7] S. Ouarab, Smarandache Ruled Surfaces according to Darboux Frame in E^3 . Journal of Mathematics, vol. (2021).

Bertrand Curves and B-Lift Curves in Lorentzian 3-Space

Anıl Altinkaya¹, Mustafa Çalışkan²

1 Gazi University, Mathematics Department, Ankara, Turkey,
anilaltinkaya@gazi.edu.tr

2 Gazi University, Mathematics Department, Ankara, Turkey,
mustafacaliskan@gazi.edu.tr

ABSTRACT

The theory of curves has an important place in differential geometry. One of the known curves is the natural lift curve. The definition of the natural lift was introduced in Thorpe's "Elementary Topics in Differential Geometry" book. According to the definition, the natural lift curve was formed by combining the end points of the tangent vector of the main curve. In this study, based on Thorpe's definition, we define a new curve in Minkowski 3-space which called B-lift curve and we examine the Frenet operators of the B-lift curve. Besides, we investigate the correspondence of the Frenet vectors between the Bertrand and the B-Lift curve. Finally, we give some examples on these results.

Key Words: B-Lift curve, Bertrand curve, Frenet vectors.

REFERENCES

- [1] J. A. Thorpe, Elementary Topics in Differential Geometry, Springer-Verlag, New York, Heidelberg-Berlin, 1979.
- [2] M. Do. Carmo, Differential Geometry of Curves and Surfaces, Prentice-Hall. Inc. Englewood Cliffs, New Jersey, 1976.

Generating Circle-Foliated Transition Surfaces in Lorentz Space

Çağla RAMİS¹ and Yusuf YAYLI²

1 Nevşehir Hacı Bektaş Veli University Faculty of Arts and Sciences Department of
Mathematics, caglaramis@gmail.com

2 Ankara University Faculty of Sciences Department of Mathematics,
yayli@science.ankara.edu.tr

ABSTRACT

Cyclic surfaces can be defined as a one-parametric family of circles. Canal surfaces, Dupin cyclides, quadrics, the surface of revolution and transition surfaces are considered as well-known examples of cyclic surfaces. This study deals with transition surfaces foliated by the system of generating Lorentzian circles. The system of generating Lorentzian circles are constructed by a set of centers and radii functions with two parameters respect to the carrier Lorentzian planes. We give the structural properties of these surfaces by the fundamental forms and curvatures. Finally, we obtain the curvature lines and the singularity conditions so make some classifications for them.

Key Words: Cyclic surface, transition surface, foliation.

REFERENCES

- [1] L.P. Eisenhart, A Treatise on the Differential Geometry of Curves and Surfaces, Ginn., 1909.
- [2] R. Lopez, Cyclic surfaces of constant Gauss curvature, Houston J. Math. 27 (2001), 799-805.
- [3] W. Boehm, On cyclids in geometric modeling, Comput. Aided GEom. Design 7 (1990), 243-255.

On Some Closed Sets and Circles of the Added Sierpinski Triangle

Aslıhan İklim Şen¹ and Mustafa Saltan²

1 Department of Information Technologies, Kırklareli University, Kırklareli, Turkey
asen@klu.edu.tr

2 Department of Mathematics, Faculty of Science, Eskişehir Technical University, Eskişehir,
Turkey, mustafasaltan@eskisehir.edu.tr

ABSTRACT

Fractals are interesting and fascinating shapes. One of the most important features of the fractals is self-similarity. There are basic fractal models that are solutions to many mathematical problems such as Cantor set, Sierpinski triangle and carpet, Vicsek fractal, Koch curve and snowflake, Menger sponge, Mandelbrot set and Julia sets. Many properties of these self-similar sets have been investigated from every aspect for years. Especially the Sierpinski triangle, S , has been considered as a fundamental model in various studies. In recent years, studies in which the intrinsic metrics are formulated on this fractal come to the fore. To define the intrinsic metrics by using the code representations of the points on classical fractals make possible to investigate different geometrical and topological properties of these sets. In different studies, the intrinsic metrics are formulated on the classical Sierpinski triangles, added and mod-3 Sierpinski triangle and also Vicsek fractal by this method.

In this study, we consider the added Sierpinski triangle and its intrinsic metric formula and then we investigate some circles and closed sets of this fractal. Finally, we express them as the code sets with the help of the code representations of the points by using the intrinsic metric and give figures of some of them.

Key Words: Sierpinski gasket, fractal, intrinsic metric.

Acknowledgement: This study was supported by TÜBİTAK-1002 under contract 118F356.

REFERENCES

- [1] M.F.Barnsley, Fractals Everywhere. San Diego, CA, USA: Academic Press, 1988.

- [2]** M. Saltan, Intrinsic metrics on Sierpinski-like triangles and their geometric properties, *Symmetry* 10 (2018), 204, DOI:10.3390/sym10060204.
- [3]** M. Saltan, Some Interesting code sets of the Sierpinski triangle equipped with the intrinsic metric, *IJAMAS* 57 (2018), 152-160.
- [4]** M. Saltan, Y. Özdemir and B. Demir, An explicit formula of the intrinsic metric on the Sierpinski gasket via code representation, *Turk. J. Math.* 52 (2018), 716-725.
- [5]** A.İ. Şen and M. Saltan, The formulization of the intrinsic metric on the added Sierpinski triangle by using the code representations, *Turk. J. Math.* 42 (2018), 716-725.

Some Notes on $N(k)$ -contact Metric Manifolds

Halil İbrahim Yoldaş

Department of Mathematics, Faculty of Science and Arts, Mersin University,
hibrahimyoldas@mail.com

ABSTRACT

Differential geometry, which has a wide application area in many branches of science, is one of the most active fields of study in modern mathematics. The theory of manifolds is one of the most comprehensive and important fields of study of differential geometry since manifolds explain spaces in terms of simpler and easily understandable structures. There are many special and different classes of manifolds in the literature. One of them is $N(k)$ -contact metric manifolds that are almost contact metric manifolds with nullity distribution. The first study on such manifolds were given by Tanno. Since then, such manifolds have been studied and investigated widely by several mathematicians.

On the other hand, in differential geometry, mathematicians have used many different methods to characterize the geometric properties and class of a given manifold. One of them is special vector fields. The existence of the vector fields defined on the manifolds plays a significant role in determining and characterizing the most geometric properties of the related object of the manifolds. This situation make vector fields useful tools in the study of differential geometry of manifolds. As a result of this, Therefore, in recent years, the manifolds which admit special vector fields have been investigated comprehensively by many geometers in many context.

Motivated by the above circumstances, in this work we deal with some special vector fields on $N(k)$ -contact metric manifolds, which prove to be rich in geometrical structures. Also, we obtain some important characterizations as regards both such vector fields and such manifolds.

Key Words: $N(k)$ -contact metric manifold, Sasakian Manifold, Torse-forming vector field.

REFERENCES

- [1] D. E. Blair, Contact Manifolds in Riemannian Geometry, Lecture Notes in Mathematics, Springer-Verlag, Berlin, 1976.
- [2] D. E. Blair, T. Koufogiorgos, B. J. Papantoniou, Contact Metric Manifold Satisfying a Nullity Condition, Israel J. Math., 91 (1995), 189-214.
- [3] B.-Y. Chen, Classification of Torqued Vector Fields and Its Applications to Ricci Solitons, Kragujevac J. Math., 41(2) (2017), 239-250.
- [4] U. C. De, A. Yildiz, S. Ghosh, On a Class of $N(k)$ -Contact Metric Manifolds, Math. Reports, 16 (2014), 207-217.
- [5] S. Tanno, Ricci Curvatures of Contact Riemannian Manifolds, Tohoku Math. J., 40 (1988), 441-448.
- [6] H. İ. Yoldaş, Ş. E. Meriç, E. Yaşar, On Generic Submanifold of Sasakian Manifold with Concurrent Vector Field, Commun. Fac. Sci. Univ. Ank. Ser. A1 Math. Stat., 68(2) (2019), 1983-1994.

Some Special Smarandache Ruled Surfaces According to Flc Frame in E^3

Kebire Hilal AYVACI¹, Süleyman ŞENYURT² and Davut CANLI³

¹ Ordu University, Faculty of Arts and Sciences, Depart. of Maths. kebrehilalatvace@odu.edu.tr

² Ordu University, Faculty of Arts and Sciences, Depart. of Maths. ssenyurt@odu.edu.tr

³ Ordu University, Faculty of Arts and Sciences, Depart. of Maths. davutcanli@odu.edu.tr
(use Arial 10 typeface for author's Affiliation)

ABSTRACT

In this study, we introduce some special ruled surfaces according to the Flc frame of a given polynomial curve. We name these ruled surfaces as T-D₂, T-D₁ ve D₁-D₂ Smarandache ruled surfaces and provide their characteristics such as Gauss and mean curvatures in order to specify their developability and minimality conditions. Moreover, we examine the conditions if the parametric curves of the surfaces are asymptotic, geodesic or curvature line. Such conditions are also argued in terms of the developability and minimality conditions. Finally, we give an example and picture the corresponding graphs of ruled surfaces by using Maple 17.

Key Words: Smarandache ruled surfaces, Mean Curvature, Gaussian Curvature, Flc Frame, Polynomial curves.

REFERENCES

- [1] A. Karger and J. Novak, Space Kinematics and Lie Groups. STNL Publishers of Technical Lit. Prague, Czechoslovakia. 1978.
- [2] S. Izumiya and N. Takeuchi, Special curves and ruled surfaces, Cont. to Alg. And Geo. 44 (2003), 200–212.
- [3] Y. Yu, H. Liu, S.D. Jung.. Structure and Characterization of Ruled Surfaces in Euclidean 3-space , Applied Math. and Comp 233 (2014), 252-259.
- [4] P. Alegre, K. Arslan, A. Carriazo, C. Murathan and G. Öztürk, "Some special types of developable ruled surface," Hacettepe Journal of Mathematics and Statistics, 39(3) (2010), pp. 319–325.
- [5] G. Hu, H. Cao, J. Wu, and G. Wei. "Construction of developable surfaces using generalized C-Bezier bases with shape parameters," Computational and Applied Mathematics, 39(3) (2020).

- [6] F. Özsoy and İ. Güven Özel Eğrilerden Üretilen Regle Yüzeyler, Yüksek Lisans Tezi, Matematik, Temmuz, 44 Syf, Gaziantep, 2019.
- [7] C. Ashbacher, Smarandache geometries, Smarandache Notions Journal, 8,1-3 (1997), 212-215.
- [8] M. Turgut and S.Yilmaz, Smarandache Curves in Minkowski Space-time, Int. J. Math. Comb., 3 (2008), 51-55.
- [9] A.T. Ali, Special Smarandache curves in the Euclidean space, International Journal of Mathematical Combinatorics, 2 (2010), 30-36.
- [10] S. Senyurt and A. Çalışkan N^*C^* Smarandache Curves of Mannheim Curve Couple According to Frenet Frame. International Journal of Mathematical Combinatorics, 1 (2015), 1-13.
- [11] S. Senyurt, A. Çalışkan and Ü. Çelik, N^*C^* Smarandache curves of Bertrand curves pair according to Frenet frame. International Journal of Mathematical Combinatorics, 1 (2016), 1-7.
- [12] M. Çetin, Y. Tuncer ve M.K. Karacan, Smarandache Curves According to Bishop Frame in Euclidean 3-Space, arXiv:1106. 3202 v1 [math. DG], (2011).
- [13] Ö. Bektaş ve S. Yüce, Special Smarandache Curves According to Darboux Frame in E^3 Rom. J. Math. Comput. Sci., 3 (2013), 48-59.
- [14] K. Taşköprü and M. Tosun, Smarandache curves according to Sabban frame on S^3 , Boletim da Sociedade Paranaense de Matemtica, 32 (2014), 51-59.
- [15] N. Demircan Smarandache Eğrilerine Dair, Sinop Üniversitesi Fen Bilimleri Enstitüsü, Yüksek Lisans Tezi, Sinop, (2015).
- [16] M.B. Mandal and C. Ekici q-çatılı Timelike Eğrilerin Smarandache Eğrileri Üzerine, Yüksek Lisans Tezi, Kasım, Eskişehir (2019).
- [17] S. Ouarab, -Smarandache Ruled Surface and-Smarandache Ruled Surface according to Alternative Moving Frame in. Journal of Mathematics, (2021).
- [18] S. Ouarab, Smarandache Ruled Surfaces according to Darboux Frame in. Journal of Mathematics, (2021).

The properties of projective, concircular and conharmonic curvature tensor fields on the complex Sasakian manifold

Aysel TURGUT VANLI¹

¹ Gazi University Faculty of Science Department of Mathematics, avanli@gazi.edu.tr

ABSTRACT

Real Sasakian manifolds are well known in the literature, and there are many articles on this subject. But the concept of complex Sasakian manifold is a new field of study. Studies on complex Sasakian manifolds are scarce in the literature. Properties of many tensors and curvatures on real Sasakian manifolds have been studied, and there are many articles in the literature on this subject. Complex Sasakian manifolds are very difficult to work with, the calculations are complex and long. For this reason, obtaining the properties of tensor fields and curvature tensor fields on the complex Sasakian manifold requires a very laborious work. The best known of the curvature tensor fields are the projective, concircular and conharmonic curvature tensor fields. Therefore, in this article, the properties of projective, concircular and conharmonic curvature tensor fields on the complex Sasakian manifold are investigated.

Key Words: Projective, concircular, conharmonic, complex Sasakian manifolds

REFERENCES

- [1] D. E. Blair, Riemannian Geometry of Contact and Symplectic Manifolds, 2nd edn. Birkhauser, Boston 2010.
- [2] D. Fetcu, Harmonic maps between complex Sasakian manifolds. Rendiconti del Seminario Matematico. Universita e Politecnico Torino, 64, (2006). 319-329.
- [3] B. Foreman, Complex contact manifolds and hyperkahler geometry, Kodai Mathematical Journal, 23 (2000) 12–26.
- [4] A. Turgut Vanlı, İ. Ünal and K. Avcu, On Complex Sasakian manifolds, Afrika Matematika, v.32 (2021), 507-516.

MATHEMATICS EDUCATION

A Disease Modeling Approach to Simulate the Spread of Math Anxiety Between Students

Dilek Soysal¹, Majid Bani-Yaghoub²

¹Department of Mathematics and Statistics, University of Missouri-Kansas City
Kansas City, Missouri 64110-2499, USA
dswbz@mail.umkc.edu

²Department of Mathematics and Statistics, University of Missouri-Kansas City
Kansas City, Missouri 64110-2499, USA
baniyaghooubm@umkc.edu

ABSTRACT

Tobias and Weissbrod (1980) define math anxiety as “the panic, helplessness, paralysis, and mental disorganization that arises among some people when they are required to solve a mathematical problem” that affects a large percentage of the population [1]. There is mounting evidence that math anxiety is contagious [2]. In the present study, we will simulate the spread of math anxiety between students using a disease modeling approach. We use a next generation matrix approach to calculate the basic reproduction number, R_0 , associated with math anxiety. We prove that students remain anxiety free when $R_0 < 1$ whereas math anxiety becomes endemic when $R_0 > 1$. In other words, when one anxious student transmits their math anxiety to more than one classmate, math anxiety will spread throughout the sample population. Otherwise, the class of students will eventually become anxiety-free. Using numerical simulations of our model, we show that the number of students with math anxiety can periodically increase or decrease during the semester. The merit of the present work is the innovative use of disease modeling to understand the dynamics of math anxiety in a population of students enrolled in a math course.

Key Words: Math Anxiety, Mathematical Modelling, Compartment Model, Numerical Simulation, Mathematical Epidemiology, Stability Analysis.

ACKNOWLEDGEMENT:

The authors would like to acknowledge the work and comments of Mr. Arash Arjmand and Deepak Sireeshan.

REFERENCES

- [1] S. Tobias and C. Weissbrod, (1980). Anxiety and mathematics: An update. *Harvard Educational Review*, 50(1) (1980), 63–70.
- [2] A. Gurin, G. Jeanneret, M. Pearson, M. Pulley, A. Salinas and C. W. Castillo-Garsow, The dynamics of math anxiety as it is transferred through peer and teacher interactions. Tech. Rep. MTBI-14-05M, Arizona State University, 2017.

An Application On Comparison of Student Achievements in Face-to-Face and Online Education Models in University Mathematics Education

Göksu İnan¹, Zeynep Hande Toyganözü² and Cüneyt Toyganözü³

1 Suleyman Demirel University, Institute of Social Sciences, Isparta, Turkey,
yl2030227008@ogr.sdu.edu.tr

2 Suleyman Demirel University, Faculty of Arts and Sciences, Department of Mathematics,
Isparta, Turkey, handetoyganozu@sdu.edu.tr

3 Suleyman Demirel University, Faculty of Economics and Administrative Sciences, Department
of Econometrics, Isparta Turkey, cuneyttoyganozu@sdu.edu.tr
(use Arial 10 typeface for author's Affiliation)

ABSTRACT

Today, online education is an education model that is used not only in the education sector, but also in different sectors, regardless of time and place. With the development of technology this education model has become one of the basic education models that provide lifelong learning opportunities.

Due to the Covid-19 coronavirus, which emerged in the last months of 2019 and affected the whole world as a global epidemic, universities in our country had to take a break from traditional face-to-face education in line with the decision taken by YÖK. In this period, which is called the coronavirus pandemic, universities continued their education via online, depending on their digital infrastructure. In this way, it is aimed to reduce the negative effect of the pandemic by participating in the classes in the virtual environment from where they are, without coming to the university campuses.

Mathematics education is fundamental in our country as in the world. The existence of a general prejudice about the difficulty of learning mathematics can affect the educational process. For this reason, the effectiveness of face-to-face and online education methods in mathematics education can be discussed.

In this study, students' achievements in face-to-face and online education methods of a basic theoretical course given in freshmen year in the Department of Mathematics were compared. For this, the grades of the students obtained by face-to-face education method in the 2019-2020 Fall semester and the grades obtained by the online education method in the 2020-2021 Fall semester were used. Whether

these two methods differ in terms of average success or their superiority to each other was examined using the t-statistic.

Key Words: Mathematics education, face-to-face education, online education, t-statistic.

REFERENCES

- [1] M. Başaran, E. Doğan, E. Karaoğlu, and E. Şahin, Koronavirüs (Covid-19) pandemi sürecinin getirisi olan uzaktan eğitimin etkililiği üzerine bir çalışma, *Academia Eğitim Araştırmaları Dergisi* 5(2) (2020), 368–397.
- [2] A. Bozkurt, Koronavirüs (Covid-19) pandemi süreci ve pandemi sonrası dünyada eğitime yönelik değerlendirmeler: Yeni normal ve yeni eğitim paradigması. *AUAd.* 6(3) (2020) 112-142.
- [3] W. D. Muirhead, Online education in schools, *International Journal of Educational Management*, 14(7) (2000), 315-324.
- [4] K.C. Carr and C.L. Farley, Redesigning courses for the World Wide Web, *Journal of Midwifery & Women's Health*, 48(6) (2003), 407-417.

Behavioural Analysis of a Predator-Prey Model

Seval İŞİK¹, Figen KANGALGİL²

1 Mathematics and Science Education, Sivas Cumhuriyet University, Turkey,
skaracan@cumhuriyet.edu.tr

2 Bergama Vocational High School, Dokuz Eylül University, Turkey,
figen.kangalgil@deu.edu.tr,

ABSTRACT

Multiple mathematical models have been proposed in the studies on populations, since Lotka [1] and Volterra [2] suggested to model population dynamics of predator- prey model. Population dynamics is an important subject in the mathematical biology. Among the models mentioned above, predator-prey models have an important place in the biological field. In mathematical biology, the dynamics of the predator-prey models, which can show relationships between two predator-prey species, have been a great interest by mathematician [3-8] and the reference therein. Qualitative analyses of these studies were found many rich dynamics.

The recent articles have been shown that discrete-time models described by difference equations are appropriate than the continuous-time models governed by differential equations. So, in this study we considered a discrete-time model. We showed that the existence and topological classification of the fixed points of the considered model. Moreover, we discussed bifurcation analysis of a predator-prey model via bifurcation theory. Furthermore, our theoretical prediction is proved by numerical results that all calculations described with the aid of Mathematica. Finally, the analytical results obtained are interpreted biologically in support of numerical studies.

Key Words: Predator- Prey Model, Stability, Bifurcation.

REFERENCES

- [1] A.J. Lotka, Elements of Physical Biology, Waverly Press, Williams&Wilkins Company, Baltimore, MD, USA 1925.

- [2] V. Volterra, Variations and Fluctuations of the Number of Individuals in Animal Species Living Together, ICES J. Mar. Sci. 3 (1928), 3-51.
- [3] J. Smith, Mathematical ideas in biology, Cambridge Press, Cambridge 1968.
- [4] L. Xiaoli and X. Dongmei, Complex dynamics behaviors of a discrete-time predator-prey system, Chaos Solitons and Fractals, 32, (2007), 80-94.
- [5] S. N. Elaydi, An Introduction to Difference Equations, Springer-Verlag, New York, NY, USA, 1996.
- [6] Kuznetsov Y. A., "Elements of Applied Bifurcation Theory", Springer-Verlag, New York, NY, USA, 2nd edition, 1998.
- [7] S. Wiggins, Introduction to Applied Nonlinear Dynamical System and Chaos, vol. 2, Springer-Verlag, New York, NY, USA, 2003.
- [8] Khan, A. Q., "Neimark-Sacker bifurcation of a two-dimensional discrete-time predator-prey model", SpringerPlus, 5 (2016), 126-136.

Development of Preservice Mathematics Teachers' TPACK through Digital Storytelling

Tugba UYGUN¹,

¹ Alanya Alaaddin Keykubat University, Dr., tugba.uygun@alanya.edu.tr

ABSTRACT

Mathematics education, that students are forced to learn, can become more enjoyable when taught by a method that attracts students. In this respect, the use of digital stories with visual and auditory elements that both narrators and listeners enjoy may be effective in mathematics education. Digital storytelling can be used as a beneficial tool enhancing learning and interaction among students (Hofer & Swan, 2005). In this respect, it may be important to provide preservice mathematics teachers opportunities to acquire the necessary knowledge and skills about creating and using digital stories effectively in classrooms. Hence, the study focused on the preservice mathematics teachers' technological pedagogical content knowledge (TPACK) (Mishra & Koehler, 2006) in the context of digital stories. The participants of the current study designed with one group pretest-posttest experimental design as a kind of experimental design was composed of 50 preservice mathematics teachers. Initially, they were asked to design lessons by using any technological tool. Then, they were instructed based on designed instructional sequence related to how to use the Powtoon, and to write stories for creating digital stories. Then, the participants were asked to design lessons using digital stories prepared by them. These lessons were analysed by a rubric prepared by the author. The findings of the study showed that most of the lessons designed and created using digital stories could show preservice teachers' technological pedagogical content knowledge effectively and help preservice mathematics teachers improve their TPACK thorough the instructional process enacted based on digital storytelling.

Key Words: Digital storytelling, mathematics, technological pedagogical content knowledge, preservice mathematics teachers.

REFERENCES

- [1] M. Hofer and K. Swan, Digital storytelling: moving from promise to practice. In C. Crawford et al. (Eds.), Proceedings of Society for Information Technology and Teacher Education International Conference, March 19 (pp. 679-684), Chesapeake, VA: AACE, (2006).
- [2] P. Mishra and M.J. Koehler, Technological pedagogical content knowledge: A framework for teacher knowledge. Teachers College Record, 108(6) (2006), 1017–1054.

EXAMINING THE EFFECTS OF PRE-SCHOOL EDUCATION ON MATHEMATICS ACHIEVEMENT AT DIFFERENT SOCIOECONOMIC LEVELS: THE CASE OF TIMSS 2019 TURKEY

Burcu Parlak

Ministry National Education, not a member, enbaparlak@gmail.com

ABSTRACT

Research shows that the skills acquired during the pre-school education process in early childhood are important for children to take the necessary foundation for their success in primary school and beyond. Because the experiences that children have from pre-school to the end of primary school form the basis of their future life skills (OECD, 2011a).

Studies show that children who receive pre-school education are in a better position in terms of social, emotional and physical development, language and self-care skills development, academic success in the first years of primary education compared to children who do not (Tantekin, Erden & Altun, 2014). Pre-school education prepares children for primary school in terms of all developmental areas and reduces the differences arising from socioeconomic level (Yazar, 2013). A qualified pre-school education strengthens social equality by compensating for the developmental differences of children.

Studies reveal that socioeconomic differences are also effective on students' academic success (Yayan & Berberoğlu, 2004; Sandoval-Hernandez & Bialowolski, 2016). Studies show that children from families with low and middle socioeconomic status are more disadvantaged than others. The fact that children in this disadvantaged group are included in the formal education process at an early age and receive a good education further increases the importance of pre-school education.

When we look at the results of the international exam practices (TIMSS, PISA), which Turkey also participates in, two points related to pre-school education point to important findings on the academic success of students. First of all, whether students receive pre-school education or not is an important factor that causes differences between students' achievement scores. However, the results show that there is a strong relationship between the duration of preschool education and achievement scores (OECD, 2014; Mullis, I.V.S.vd, 2020).

Purpose of the study

The aim of the study is to determine whether the pre-school education status and duration of education, and depending on these variables, the literacy and numerical skills and socioeconomic levels of the students at the time they started school predict the mathematics achievement of the students.

Research Group

The research was conducted on the data of 3675 fourth grade students who participated in the TIMSS 2019 application from Turkey. Mathematics scores and questionnaire data of fourth grade students were used in the study.

Analysis of Data and Results

Multivariate linear regression analysis was used to determine the variables that predicted students' success. Considering the findings, it was determined that 1094 students, approximately one third of the group, did not receive pre-school education. Also, it was determined that as the duration of pre-school education increased, mathematics achievement also increased. When the results of the regression analysis are examined, it is seen that *socioeconomic level*, *early numeracy activities beginning school*, *early literacy activities beginning school*, are significant predictors of students' mathematics achievement. This variable gives a moderate and significant relationship with students' math scores ($R=0.584$, $R^2=0.341$, $p<0.05$).

Key Words: Mathematics achievement, pre-school education, socioeconomic level, TIMSS.

REFERENCES

- [1] I.V.S. Mullis, M.O. Martin, P. Foy, L.D. Kelly, and B. Fishbein *TIMSS 2019 international results in mathematics and science*. Boston College, TIMSS & PIRLS International Study Center, (2020).
- [2] OECD. (2011a). PISA in Focus: Does participation in pre-primary education translate into better learning outcomes at school?
- [3] OECD. (2014). PISA 2012 results in focus: What 15-year-olds know and what they can do with what they know?
- [4] A. Sandoval-Hernandez and P. Bialowolski, Factors and conditions promoting academic resilience: a TIMSS based analysis of five Asian education system. *Asia Pacific Educ. Rev.* 17 (2016), 511-520.
- [5] F. Tantekin Erden and D. Altun, Sınıf öğretmenlerinin okul öncesi eğitim ve okul öncesinde ilköğretime geçiş süreci hakkındaki düşüncelerinin incelenmesi. *İlköğretim Online*, 13(2) (2014), 481-502.
- [6] B. Yayan and G. Berberoğlu, A re-analysis of the TIMSS 1999 mathematics assessment data of the Turkish students. *Studies in Educational Evaluation*, 30 (2004), 87–104.
- [7] A. Yazar, Okul öncesi eğitim programının ilkokula hazırlık açısından etkililiğinin incelenmesi. Yayımlanmamış doktora tezi, Atatürk Üniversitesi Eğitim Bilimleri Enstitüsü, Erzurum (2013).

On Stability Analysis and Flip Bifurcation of a Discrete-Time Predator-Prey System with Allee Effect and Prey Immigration

Feda İlhan¹, Figen Kangalgil²

¹ Bolu Abant İzzet Baysal University, Bolu, Turkey, gumusboga_f@ibu.edu.tr

² Dokuz Eylül University, İzmir, Turkey, figen.kangalgil@deu.edu.tr

ABSTRACT

In this study the dynamics of a discrete-time predator prey model including both Allee effect and immigration factor on prey population is studied. Predator-prey model is an important subject in mathematical biology. In 1930's, famous ecologist Warder Clyde Allee observed that individual fitness declines at lower critical density often occurs in many natural populations. By Allee effect, a positive correlation is described between any measure of species fitness and population numbers. Biologically, the Allee effect term can be explained as the factors that genetic inbreeding and loss of heterozygosity lead to decreased fitness and that there will be shortage of cooperation interactions among conspecific at low densities.

Allee effect and the immigration parameter have an important role in increasing the realism of the population models, besides they help to gain a more accurate description of the model. We have concentrated on the existence and stability analysis of the equilibrium points of the model. Also, the topological classification of the equilibrium points has been made. Furthermore, dealing with the bifurcation analyses of the model the existence of Flip bifurcation is shown. The dynamical analysis of Flip bifurcation is determined by using the center manifold theorem. Some numerical simulations including stability and bifurcation diagrams are presented.

Key Words: Predator-prey model, stability analysis, Flip bifurcation.

REFERENCES

- [1] F. Kangalgil and N. Topsakal, Stability Analysis and Flip Bifurcation of a Discrete-Time Predator-Prey Model with Predator Immigration, Asian J. of Math. And Comput. Sci., 27(3) (2020), 1-10.

- [2]** G. Zuhu and J. Wei, Global Stability and Bifurcation Analysis of a Delayed Predator-Prey System with Prey Immigration, *Electronic Journal of Qualitative Theory of Diff. Eqns.*, 13 (2016), 1-20.
- [3]** Q. Din, T. Donchev and D. Kolev, Stability, Bifurcation Analysis and Chaos Control in Chlorine Dioxide-Iodine-Malonic Acid Reaction, *Match Commun. Math. Comput. Chem.*, 79 (2018), 577-606.
- [4]** F. Kangalgil, Flip Bifurcation and Stability in a Discrete_Time Prey-Predator Model with Allee Effect, *Cumhuriyet Science Journal*, 40-1 (2019), 141-149.
- [5]** Allen and J. S. Linda, *An Introduction to Mathematical Biology*, Texas Tech. University, 2007.

ON THE DEVELOPMENT OF PLANNING ABILITY IN PROBLEM SOLVING FROM GRADES 5 THROUGH 7

B.Damla BABER ELBISTAN¹ , Kolja PUSTELNIK²

¹ Social Science Faculty of Goettingen University, birgueldamla.baber@stud.uni-goettingen.de

² Mathematics Institute of Goettingen University,
kolja.pustelnik@mathematik.uni-goettingen.de

ABSTRACT

Planning ability is considered an important feature for successful problem-solving in mathematics which should develop over the years at school. It involves many different processes and cannot be measured with imprecise or over-simplified operationalizations (Boghi et al, 2006). Most studies have focused on problem-solving longitudinally within planning ability (Siegler, 1983; Gauvain and Rogoff, 1989). In this study, with the help of the Tower of London (TOL) test, which has been developed to measure planning ability, the effectiveness of executive functions for problem-solving in different degrees of complexity is investigated with N=223 students at secondary school in grades five, six and seven. In the five and six move problem sections the planning skills increased; in fact, a significant difference depending on the youngest and the oldest age is showing ($p < .05$). However, no statistically significant difference is found between the planning with respect to gender ($p > .05$).

Key Words: Tower of London Test, planning ability, problem-solving.

REFERENCES

- [1] A. Boghi, R. Rasetti, F. Avidano, C. Manzone, L. Orsi, F. D'Agata, ... and P. Mortara, The effect of gender on planning: An fMRI study using the Tower of London task. *Neuroimage*, 33(3) (2006), 999-1010.
- [2] M. Gauvain and B. Rogoff, Collaborative problem solving and children's planning skills. *Developmental psychology*, 25 (1) (1989), 139.
- [3] R.S. Siegler, Five generalizations about cognitive development. *American Psychologist*, 38(3) (1983), 263.

Pre-service and In-service Teachers' Proof Schemes and Their Opinions on Mathematical Proof

Prof. Dr. Sare Şengül¹, Arş. Gör. Dilara Yılmaz²

¹ Marmara Üniversitesi Atatürk Eğitim Fakültesi Matematik Öğretmenliği A.B.D.

² Kocaeli Üniversitesi Eğitim Fakültesi Sınıf Öğretmenliği A.B.D.

ABSTRACT

Making sense of mathematical proof is equivalent to making sense of mathematics as a discipline (Pair & Calva, 2020). At this point, questions such as how to teach mathematics, how to learn it and how to understand how it is learned arise (Harel, 2008). How and in what ways to teach mathematics; The thought patterns that are desired to be gained in the student and many more are the focus of proof-based teaching, and educators have important tasks at this point. In this study, the views and proof schemes of primary school teachers and prospective teachers who will be teachers of the future on mathematical proof were examined. When the literature is examined, the skills in probability and statistics at the secondary school level do not develop at the expected level (Bakırcı, 2014; Dereli, 2009; Güven & Özmen, 2014; Memnun, Altun & Yılmaz, 2010; Mills & Holloway, 2013), and the majority of secondary school students (Aydoğdu İskenderoğlu, 2003; Ören, 2007) cannot think analytically during the proof process. These results show that these subjects and areas should be supported for lower level students such as primary school and even pre-school. In this study, perspectives of primary school teachers and pre-service primary teachers about mathematical proof and proof schemes were examined. Case study design and maximum diversity sampling were used. Participants' views on mathematical proof were collected with a questionnaire consisting of open-ended questions. In order to examine the proof schemes, the problems created by the researchers that require graphing about data processing were used. The analysis of the obtained data was analyzed using the content analysis method, which is the most commonly used method among qualitative data analysis methods. According to the findings obtained in the research, the classroom teachers and candidates emphasized the meanings of proving, proving the truth and revealing with concrete evidence; It has been determined that they associate the

purpose of proving with increasing confidence in information and confirming meaningfulness. It has been observed that being understandable, consistent, objective, and free from contradictions are the most frequently stated expressions among the features of the proof. According to the findings regarding proof schemes, it was seen that teachers and pre-service teachers mostly had experimental proof schemes. In the light of these results, researchers are recommended to conduct research involving pupils, primary teachers and teacher candidates for the development of primary school level mathematics education.

Key Words: Pre-service and in-service teachers, proof based research, problem solving, data process

REFERENCES

- [1] Aydoğdu İskenderoğlu, T. (2003). Farklı sınıf düzeylerindeki öğrencilerin matematik problemlerini kanıtlama süreçleri, (Yayınlanmamış Yüksek Lisans Tezi), Abant İzzet Baysal Üniversitesi, Sosyal Bilimler Enstitüsü, Bolu.
- [2] Bakırcı, S. (2014). İlköğretim 7. sınıf öğrencilerinin olasılıkla ilgili problem çözme süreçlerinin incelenmesi üzerine nitel bir çalışma. (Yayınlanmamış Yüksek Lisans Tezi), Necmettin Erbakan Üniversitesi Eğitim Bilimleri Enstitüsü, Konya.
- [3] Dereli, A. (2009). Sekizinci sınıf öğrencilerinin olasılık konusundaki hataları ve kavram yanılgıları. (Yayınlanmamış Yüksek Lisans Tezi), Eskişehir Osmangazi Üniversitesi Fen Bilimleri Enstitüsü, Eskişehir.
- [4] Güven, B. ve Özmen, Z.M. (2014). Ortaokul 8. sınıf öğrencilerinin grafik okuryazarlığı düzeylerinin belirlenmesi. 11. Ulusal Fen Bilimleri ve Matematik Eğitimi Kongresi, Adana, 264-268.
- [5] Harel, G. (2008). DNR perspective on mathematics curriculum and instruction Part II: With reference to teacher's knowledge base Zentralblatt'ur Didaktik der Mathematik.
- [6] Memnun, D.S., Altun, M. ve Yılmaz A. (2010). İlköğretim sekizinci sınıf öğrencilerinin olasılıkla ilgili temel kavramları anlama düzeyleri. Eğitim Fakültesi Dergisi 23 (1), 11-29.
- [7] Mills, J.D. & Holloway C.E. (2013). The development of statistical literacy skills in the eighth grade: exploring the TIMSS data to evaluate student achievement and teacher characteristics in the United States. Educational Research and Evaluation, 19(4), 323-345. <http://dx.doi.org/10.1080/13803611.2013.771110>.
- [8] Ören, D. (2007). An Investigation of 10th Grade Students' Proof Schemes in Geometry with Respect to Their Cognitive Styles and Gender. (Yayınlanmamış Yüksek Lisans Tezi), Orta Doğu Teknik Üniversitesi, Ankara.
- [9] Pair, J. & Calva, G. (2020). Understanding the roles of proof through exploration of unsolved conjectures. Mathematics Education Across Cultures: Proceedings of

the 42nd Meeting of the North American Chapter of the International Group for the
Psychology of Mathematics Education, Mexico. Cinvestav / AMIUTEM / PME-NA
Doi: 10.51272/pmena.42.2020-183.

Preservice High School Mathematics Teachers' Experiences on their Fieldwork During the COVID-19 Pandemic

Ebru Ersari

Balıkesir University, ebru.ersari@balikesir.edu.tr

ABSTRACT

The COVID-19 Pandemic impacted many fields including the field of education (Bakker & Wagner, 2020; Erduran, 2020; Onyema, Eucheria, Obafemic, Sen, Atonye, Sharma, 2020). Most schools had to shift from face-to-face learning to distance learning in a short period of time (Ng, P. T., 2021). Students, teachers, and preservice teachers had different learning and teaching experiences during the pandemic (Baber, 2020). The purpose of this study was to explore three high school preservice teachers' field experience during the COVID-19 pandemic. Their performances on the Teaching Practice course that they took in their eighth semester were analysed. The researcher of this study was the instructor of the course. One preservice teacher observed one high school teacher's classroom, and two preservice teachers observed another high school teacher's classroom online. They shared their experiences on the virtual fieldwork in the Teaching Practice course. All course sessions were recorded. In addition to attending course meetings online, as part of the course, the preservice teachers reported their experience over the semester. The reports consist of 12 week online observations of the high school classroom. The data consists of the preservice teachers' videotaped sessions as well as their written work. The findings and conclusions will be shared.

Key Words: Pandemic, preservice teacher education, field experience.

REFERENCES

- [1] A. Bakker and D. Wagner, Pandemic: Lessons for today and tomorrow? Educational Studies in Mathematics, 104, (2020), 1-4.

- [2] H. Baber, H, Determinants of students' perceived learning outcome and satisfaction in online learning during the pandemic of COVID-19. *Journal of Education and e-Learning Research*, 7(3), (2020), 285-292.
- [3] S. Erduran, Science education in the era of a pandemic: How can history, philosophy and sociology of science contribute to education for understanding and solving the COVID-19 crisis? *Science & Education*, 29, (2020), 233–235.
- [4] P.T. Ng, Timely change and timeless constants: COVID-19 and educational change in Singapore. *Educational Research for Policy and Practice*, 20(1) (2021), 19-27.
- [5] E.M. Onyem, N.C. Eucheria, F.A. Obafemi, S. Sen, F.G. Atonye, A. Sharma and A.O. Alsayed, (2020). Impact of Coronavirus pandemic on education. *Journal of Education and Practice*, 11(13) (2020), 108-121.

THE EFFECTS OF ACTIVITIES CARRIED OUT WITH WEB 2.0 TOOLS ON SOME FIELD COMPETENCIES OF MATHEMATICS TEACHER CANDIDATES

Aybige ARABACI¹, Keziban ORBAY²

1 Suluova Cumhuriyet Secondary School, Amasya, aybige9045@gmail.com

2 Amasya University, Faculty of Education, Amasya, keziban.orbay@amasya.edu.tr

ABSTRACT

The fact that educational environments are affected by rapid changes in information and communication technologies necessitate the use of technology in these environments (Akkoyunlu, 1995). This situation has revealed the importance of developing the competencies of teachers who will integrate technology to education (Dağ, 2016). Web 2.0 tools, one of the best technologies that can provide integration into education nowadays, offer innovative web-based alternative digital applications that can provide professional development of teachers without time and place restrictions (Dalgarno ve Lee, 2010). Therefore, the aim of this study was to examine the effects of design experiences of mathematics teaching course activities organized with Web 2.0 tools on teacher candidates' technological pedagogical content knowledge, their attitude towards instructional technologies, and their self-efficacy for practical content development with Web 2.0 tools.

The study was structured in a single-group pretest-posttest design, one of the quantitative methods. The universe of the study consists of 33 teachers candidates studying in a primary school mathematics teaching program in a state university in the 2020-2021 academic year. "Technological Pedagogical Content Knowledge", "Web 2.0 Practical Content Development Self-Efficacy Belief" and "Attitudes Towards Instructional Technologies" scales were used as data collection tools. The study was carried out within the scope of the "Teaching Numbers" course. Twelve different Web 2.0 tools, and how to create examples of activities that can be used in mathematics teaching was presented at the end of the course through the videos prepared by the researcher for 10 weeks. The content of the presented activity examples was consist

of the topics covered in the course. At the end of the process, teacher candidates were asked to prepare and present a lesson plan integrated with Web 2.0 tools.

According to the results obtained from the study, it is concluded that, design experiences of mathematics teaching course activities of the teacher candidates had significant effects on “Web 2.0 Practical Content Development Self-Efficacy Belief” level and the “belief regarding usage of instructional technology in lesson”, “appreciation to usage of instructional technology in lesson” and “unappreciated using instructional technology” sub-dimensions of the “Attitude Towards Instructional Technologies” scale; however there were no significant effect on the “Technological Pedagogical Content Knowledge” scale and it’s sub-dimensions.

Key Words: Web 2.0 tools, Instructional technology, Teacher candidates

REFERENCES

- [1] Akkoyunlu, B. (1995). Bilgi teknolojilerinin okullarda kullanımı ve öğretmenlerin rolü. *Hacettepe Üniversitesi Eğitim Fakültesi Dergisi*, 11(11).
- [2] Dağ, F. (2016). Examination of the professional development studies for the development of technological competence of teachers in Turkey in the context of lifelong learning. *Journal of Human Sciences*, 13(1), 90-111.
- [3] Dalgarno, B. & Lee, M. J. (2010). What are the learning affordances of 3-D virtual environments? *British Journal of Educational Technology*, 41(1), 10-32.

The Effectiveness of the Subject Area Textbook Review Course Amid the COVID-19 Pandemic

Ebru Ersari

Balıkesir University, ebru.ersari@balikesir.edu.tr

ABSTRACT

Even though courses had to be switched to online during the COVID-19, the effectiveness of this shift needs to be investigated (Daniel, 2020; Tartavuela, Albu, Albu, Dieaconescu, & Petre, 2020). Institutions including schools had to take precautions to survive or continue to function (Azhari & Fajri, 2021, Engelbrecht, Borba, Llinares, & Kaiser, 2020). Compared to the past, research on textbook analysis has also been increased (citation). Textbooks could have the potential to improve preservice teachers' knowledge. Thus, the purpose of this study is to explore the effectiveness of the Subject Area Textbook Review Course during the COVID-19 pandemic. The participants were 57 preservice mathematics teachers, and they took the course over the summer 2019 during the pandemic. The course was a required course that preservice mathematics teachers needed to take in their eighth semester in their program. They met 4,5 hours per week consisting of 3 days of 90 minutes per day. Mathematics textbooks from the website of the Educational Informatics Network (EBA) were reviewed over the course. Preservice teachers needed to present their review of the books as part of the course. The course sessions over the course were videotaped. The results of the analysis of the course materials will be discussed.

Key Words: COVID-19, pandemic, preservice teacher education, textbook analysis

REFERENCES

- [1] Daniel, S. J. (2020). Education and the COVID-19 pandemic. *Prospects*, 49(1), 91–96.
- [2] Azhari, B., & Fairi, I. (2021) Distance learning during the COVID-19 pandemic: School closure in Indonesia. *International Journal of Mathematical Education in Science and Technology*, <https://doi.org/10.1080/0020739X.2021.1875072>

- [3] Davis, J. D. (2009). Understanding the influence of two mathematics textbooks on prospective secondary teachers' knowledge. *Journal of Mathematics Teacher Education*, 12(5), 365–389.
- [4] Engelbrecht, J. Borba, M.C., Llinares, S. & Kaiser, G., (2020) Will 2020 be remembered as the year in which education was changed? *ZDM Mathematics Education*, 52, 821–824.
<https://doi.org/10.1007/s11858-020-01185-3>
- [5] Tartavulea, C. V., Albu, C. N., Albu, N., Dieaconescu, R. I., & Petre, S. (2020). Online teaching practices and the effectiveness of the educational process in the wake of the COVID-19 pandemic, *Amfiteatru Economic*, 22(55), 920–936.

THE FIELD EXPERIENCES ON THE ONLINE APPLICATION OF THE CASE STUDY METHOD IN MATHEMATICS LESSON

Sare Sengül¹ and Gülşah Gerez Cantimer²

¹ Marmara University, zsengul@marmara.edu.tr

² Ministry of Education, Sakarya/Serdivan IHO, gulsahgerez@subu.edu.tr

ABSTRACT

The case study method is one of the effective methods that can be used to provide learning [1], [2] and has areas of use in different disciplines of education [3]. The case study method which brings students together with real life problems [4], enables students to develop their ability to use the information they learn in lessons in practice and to develop their thinking and decision-making skills together with their friends [5]. Although the case study method is not a new teaching method, there are limited studies in this field in mathematics education [6], [7], [8]. For this reason, the applicability of the case study method in the mathematics lesson was discussed.

With this study it is aimed to examine the field experiences obtained in terms of teachers and students within the scope of the online application of the case study method in the mathematics lesson. The study was carried out with 7th grade students from two different branches who were studying at a public school in Sakarya during the fall semester of the 2020-2021 academic year and took distance education courses. The participants of the study consisted of 23 students who constantly followed the online courses during the pandemic process. The study was designed in a qualitative research design and the data collection tools of the study; The opinions of the researchers about the preparation of the case study lesson plans before the application, the teacher's in-class observations during the application of the lesson plans and the written opinions of the students about the application. The experiences gained during this process are presented separately from the perspective of the teacher and the student.

According to the findings; there have been difficulties experienced in the design of the case study, in which daily life problems can be questioned in an empathic, intellectual and moral dimension. It has been determined that the case studies designed according to the experiences gained from the teacher's point of view are

structured based on the pre-cognitive knowledge of the students, the difficulties experienced in the efforts to communicate the case studies within the scope of the mathematical context, the difficulties experienced by the students in the course follow-up due to the online learning environment. In terms of students, even if they are not at a sufficient level of mathematical knowledge, the ease of expressing their opinions within the scope of the case studies prepared, the ease of intellectual flexibility brought by the transfer of mathematical structures to a social context, the gains in awareness of their own personal rights, the recognition of the responsibilities required by living in a social environment, the respect for different views. As a result of this process, suggestions will be presented to researchers about how to design case studies on different subjects and the issues to be considered during the implementation.

Key Words: Mathematics education, case study method, online mathematics lesson, field experiences.

REFERENCES

- [1] D. Hung, D. T. Chen and S. C. Tan, A social- constructivist adaptation of casebased reasoning: Integrating goalbased scenarios with computer- supported collaborative learning. *Educational Technology*, 43(2) (2003), 30-35.
- [2] K. Ü. Açıkgöz, *Aktif öğrenme*. İzmir, Biliş Yayıncılık, 2009.
- [3] S. Şahin, B. Atasoy and S. Somyürek, Öğretmen eğitiminde örnek olay yöntemi. *Gaziantep University Journal of Social Sciences*, 9(2), 2010.
- [4] V. Sönmez, *Program geliştirmede öğretmen el kitabı*. Ankara, Anı Yayıncılık, 2008.
- [5] D. Gözütok, *Öğretim ilke ve yöntemleri*. Ankara, Ekinoks, 2007.
- [6] C. Barnett, Building a case-based curriculum to enhance the pedagogical content knowledge of mathematics teachers. *Journal of Teacher Education*, 42(4) (1991), 263-272.
- [7] D. H. Jonassen and J. Hernandez-Serrano, Case-based reasoning and instructional design: Using stories to support problem solving. *Educational Technology Research & Development*, 50(2) (2002), 65-77.
- [8] C. Öksüz and S. Uça, Matematik dersinde probleme dayalı öğrenme üzerine bir örnek olay. *Adnan Menderes Üniversitesi Eğitim Fakültesi Eğitim Bilimleri Dergisi*, 2(2) (2011), 20-29.

The Investigation of Studies on Case Study Method in Education

Gülşah Gerez Cantimer¹ and Sare Şengül²

¹ Ministry of Education, Sakarya/Serdivan IHO, gulsahgerez@subu.edu.tr

² Marmara University, zsengul@marmara.edu.tr

ABSTRACT

The case study method is a method that helps fill the gap between theory and practice in the teaching environment by bringing students face to face with real life experiences [1]. The case study method has application in different disciplines and students can learn problem solving, critical thinking, reflective thinking, decision making, multicultural awareness, etc. It is very effective in gaining skills [2].

The study aimed to examine the researches on the case study method in education through systematic review. Thus, a general framework from the descriptive features of the relevant researches from their aims to their results was drawn and their tendencies were evaluated. In the study, a total of 48 researches that consisted of 21 articles, 20 master's thesis and 7 doctoral dissertations published between 2010-2020 were included. The researches were examined within the scope of the education field that the case study method was used, purpose, method, sample, data collection tools and results by content analysis. The obtained data were presented on tables with frequencies.

As a result of the study there were determined that as the education area where the case study method was used, that the most researches were made in the fields of teacher education, chemistry education and Turkish education; that quantitative research methods were mostly preferred; that mostly working with students and preservice teachers; that it was determined that scale and achievement tests were used. When the researches examined were evaluated within the scope of their purposes, the effects of the use of the case study method on academic achievement were related to critical thinking, creative thinking, social problem solving, decision making, etc. in the field of education. It was observed that the effect on gaining skills was tried to be determined. Within the scope of the results obtained, the findings regarding the effects of the use of the case study method on knowledge, skills and affective areas were determined. In this direction, different studies can be designed and implemented with the participation of interdisciplinary studies on the use of the

case study method in the field of education and with the participation of student groups at all levels, starting from younger age groups, instructors, teachers and preservice teachers who are the implementers of the method and students' parents who have an important role in the education of students. The effect size can be increased by sharing the results.

Key Words: Education, case study method, systematic review.

REFERENCES

- [1] C. Stensmo, Case methodology in teacher education compared to traditional academic teaching: A field experiment. Paper presented at the 8th European Conference for Research on learning and Instruction (EARLI), August 24-28, Sweden, 1999.
- [2] S. Şahin, B. Atasoy and S. Somyürek, Öğretmen eğitiminde örnek olay yöntemi. Gaziantep University Journal of Social Sciences, 9(2) (2010), 253-277.

The Opinions of Secondary School Students' Parents about Mathematics Course in Distance Education

Zeynep ÇAKMAK GÜREL

Erzincan Binali Yildirim University, Education Faculty, zcakmak@erzincan.edu.tr

ABSTRACT

This study aims to determine the opinions of secondary school students' parents related to the mathematics course in distant education. A case study method was employed in the research. The sample of the research consisted of 405 parents whose children were studying at the state secondary schools in the city centre of a medium-scale province in the Eastern Anatolia Region. An online form was prepared to determine the opinions of the parents and sent to them. The parents' opinions were asked with the questions related to distance education, mathematics course in distance education, their level of responsibility and their ability to help students. In addition, the demographic information of the students was obtained. The SPSS package program was applied in the data analysis and the chi-square test was done. According to the findings, it was determined that the opinions of the majority of parents on distance education were negative (56%); therewithal, it was not appropriate to the mathematics course (55%). Moreover, it was noticed that the parents, who claimed positive opinion for the distance education, stated, at the same time, that mathematics was appropriate to distance education; on the other hand, those, who expressed a negative opinion, thought that distance education was not appropriate to mathematics course. It was determined in the study that, there was not a significant correlation between the opinions of the parents related to mathematics in distance teaching and students' gender, number of siblings, grade reports variables; there was a significant relationship with class-level variables. Considering this, it can be claimed that the parents of 8th class students expressed more negative opinions compared with the parents of other classes. In addition, it was determined that the parents, who expressed positive opinions related to mathematics in distance education, generally expressed positive opinions; those, who expressed negative opinions related to mathematics in distance education, generally expressed a negative opinion. Besides, a significant correlation was found between the opinions of parents related to mathematics in distance education and the responsibility towards the student. Most of the parents, who claimed negative opinions in distance education, stated that they had more responsibility. It was determined that half of the parents thought that the students needed extra help in a mathematics course in distance education, and some of the parents stated that they helped their children. Consequently, it can be claimed that as the class level increases the opinions of students and parents on distance education are negative because of the anxiety related to the exam. In addition, it was noticed that the responsibility of parents increased in this process, the students needed help in a mathematics course. For this reason, it can be claimed that they stated negative

opinions about mathematics courses in distance education. The findings of the study can be supported by the findings of qualitative studies by Kuzu (2020) and Aslan, Arı and Kanat (2021). These results can be taken into consideration in the education policies.

Key Words: Distance education, mathematics, parent

REFERENCES

- [1] Ç.i. Kuzu, Ç. İ. (2020). The views of the parents on primary school distance education program (eba tv) implemented during the covid-19 pandemic, Millî Eğitim, 49(1) (2020), 505-527.
- [2] K. Aslan, A.G. Arı and M.H. Kanat, Views of the parents on the distance education process given in the covid-19 pandemic. Ulakbilge Sosyal Bilimler dergisi, 9(57) (2021), 192-205.

Translation of the Digital Game Critter Corral into Turkish and Its Evaluation in Terms of Mathematics Education Standards and Appropriateness for Children

H. Elif DAĞLIOĞLU ¹ and Hilal GENÇ ÇOPUR ²

¹ Prof. Dr., Gazi University, Gazi Faculty of Education, Department of Preschool Education,
edaglioglu1@gmail.com

² Res. Assist. Dr., Kırşehir Ahi Evran University, Faculty of Education, Department of Preschool Education, hilal.genc40@gmail.com

ABSTRACT

Tablet computers are significantly effective in helping preschool children gain mathematical skills (Papadakis, Kalogiannakis & Zaranis, 2016; Papadakis, Kalogiannakis & Zaranis 2018). The number of digital games on tablet computers is increasing day by day (Blair, Pfaffman, Cutumisu, Hallinen & Schwartz, 2015). Therefore, it is needed to have research-based educational digital games. The aim of this research is to translate the game Critter Corral (CC), which was developed for pre-school children, into Turkish and to evaluate it in terms of mathematics education standards, its appropriateness for pre-school children. CC which is developed at Stanford University (Blair, 2013), is a research-based educational digital game for pre-school children aiming to support basic math skills related to counting such as one-to-one matching, comparison, counting, addition and subtraction. In the translation process of this game, permission was obtained from the game developer (K. P. Blair). Blair sent the English audio recordings of CC, the written version of these recordings, and the written documents containing the English expressions on the game screen, all of which were translated into Turkish by an expert in the Department of English Language Education, together with the researchers. Then, expert opinion was taken to evaluate these translations in terms of their appropriateness for the age and development of children, and the translations were completed by making necessary corrections in line with expert opinions. The researcher voiced the Turkish translations of the audio recordings and sent them to the game developer. Blair prepared the Turkish version of the game by integrating the Turkish translation of CC into the game software and requested the Turkish version to be tested by the researchers before uploading it to the application market.

Researchers played the Turkish version of the game from beginning to end and checked whether there were any deficiencies in the Turkish translations. The researchers noticed several omissions and technical errors in the audio recordings and translations and reported them to Blair. After the necessary corrections, the game was uploaded to the application market. Thus, CC was translated into Turkish and took its place in the Apple Store under the name Critter Corral Turkish. Besides, in this study the "Educational Digital Game Evaluation Form for Preschool Mathematics Education" prepared by Genç Çopur, Dağlıoğlu and Dağlı (2020) was used for the evaluation of the game. In conclusion, it is observed that CC Turkish game includes mathematical content and process standards together, supports the cognitive and psychomotor development areas of the child, has an interesting story in the game, has visual and verbal instructions, is suitable for the child's independent use, progresses gradually from simple to difficult, does not have any negative content, and includes hints and directions.

Key Words: Preschool, Math education, Digital game, Number Concept.

REFERENCES

- [1] K.P. Blair, Learning in Critter Corral: Evaluating three kinds of feedback in a preschool math game. In. Sawhney, N., Reardon, E., & Hourcade, J. P. (Eds.), Proceedings of the Interaction Design and Children 2013 Conference (pp. 372-375). New York: ACM, (2013).
- [2] K.P. Blair, J. Pfaffman, M. Cutumisu, N. Hallinen and D. Schwartz, "Testing the effectiveness of iPad math game: Lessons learned from running a multi-classroom study," Conference on Human Factors in Computing Systems - Proceedings, 18 (2015), pp. 727-734.
- [3] Papadakis, Kalogiannakis, & Zaranis, Comparing tablets and pcs in teaching mathematics: an attempt to improve mathematics competence in early childhood education. Preschool and Primary Education, 4 (2016), 241-253.
- [4] S. Papadakis, M. Kalogiannakis and N. Zaranis, Educational apps from the Android Google Play for Greek preschoolers: A systematic review. Computers & Education, 116, (2018), 139-160.
- [5] H. Genç Çopur, H., H.E. Dağlıoğlu and H. Dağlı, Google Play Store ve Apple Store'daki dijital oyunların matematik eğitimi standartları ve çocuğa uygunluğu açısından incelenmesi 75. TURAN: Stratejik Arastirmalar Merkezi, 12(47) (2020), 163-176.

LISTS OF PARTICIPANTS OF ICMME-2021

Name	Institution Name	Country
A. Nazlı Uresin	Ahep University	Turkey
Abdulhamit Kucukaslan	Pamukkale University	Turkey
Abdullah Akkurt,	Kahramanmaraş Sütçü İmam University	Turkey
Abdullah Yener	Istanbul Commerce University	Turkey
Allaberen Ashyralyev	Near East University	Turkey
Alparslan Tunaalp	Ankara University	Algeria
Alperen Öztürk	Istanbul University	Turkey
Anıl Altinkaya	Gazi University	Turkey
Aslıhan İklim Şen	Kırklareli University	Turkey
Asma Hanif	Government College University	Pakistan
Aybike Arabacı	Suluova Cumhuriyet Secondary School	Turkey
Ayhan Serbetci	Ankara University	Turkey
Aynur Keskin Kaymakçı	Selcuk University	Turkey
Aysel Turgut Vanlı	Gazi University	Turkey
Bahatdin Daşbaşı	Kayseri University	Turkey
Bilge Zehra Sergi	Gazi University	Turkey
Birgül Damla Baber Elbistan	Göttingen University	Turkey
Burak Aydos	Ankara University	Turkey
Burak Oğul	Istanbul Aydın University	Turkey
Burcu Balca	Hacettepe University	Turkey
Burcu Parlak	Ministry National Education	Turkey
Çağla Ramis	Nevşehir Hacı Bayram Veli University	Turkey
Cansu Keskin	Dumlupınar University	Turkey
Çetin Cemal Özek	Gazi University	Turkey
Davut Canlı	Ordu University	Turkey
Derya Arslan	Bitlis Eren University	Turkey
Derya Sekman	Ahi Evran University	Turkey
Dilara Yılmaz	Kocaeli University	Turkey
Dilek Soysal	University of Missouri-Kansas City	USA
Ebenezer Bonyah	University of Education Winneba	Ghana
Ebru Ersarı	Balıkesir University	Turkey
Elif Ercan	Gazi University	Turkey
Elif Ece Demir	Gazi University	Turkey

Emre Sevgi	Gazi University	Turkey
Enes Ata	University of Ahi Evran	Turkey
Ercan Çelik	Atatürk University	Turkey
Erkan Taşdemir	Kırklareli University	Turkey
Ersan Erdem	Necmettin Erbakan University	Turkey
Fadime Gökçe	University of Pamukkale	Turkey
Fatih Korkmaz	Kahramanmaraş Sütçü İmam University	Turkey
Fatih Rıza Çelik	Iğdır University	Turkey
Fatma Özceran	Necmettin Erbakan University	Turkey
Faton Merovci	University of Mitrovica	Albania
Feda İlhan	Bolu Abant İzzet Baysal University	Turkey
Fercan Filiz	Ege University	Turkey
Ferhan Şola Erduran	Gazi University	Turkey
Filiz Çıtak	Tokat Gaziosmanpaşa University	Turkey
Galina Lyutskanova-Zhekova	University of Sofia	Bulgaria
Gencay Oğuz	Ankara University	Turkey
Gökçe Efeoğlu	Gazi University	Turkey
Gökhan Mutlu	Gazi University	Turkey
Göksu İnan	Suleyman Demirel University	Turkey
Gonca Çelikten	Kafkas University	Turkey
Güler Başak Öznur	Gazi University	Turkey
Gülnur Çelik Kızılkın	Necmettin Erbakan University	Turkey
Gülnur Haçat	Gazi University	Turkey
Gülşah Başaran,	Gazi University	Turkey
Gülşah Gerez Cantimer	Ministry of Education	Turkey
Gülten Torun	Kastamonu University	Turkey
Günseli Gümüşel	Atılım University	Turkey
Gürhan Soylu	Gazi University	Turkey
Hadi Kutlay	Tokat Gaziosmanpaşa University	Turkey
Halil İbrahim Yoldaş	Mersin University	Turkey
Harun Baldemir	Cankiri Karatekin University	Turkey
Hasan Es	Gazi University	Turkey
Hasan Hüseyin Sayan	Gazi University	Turkey
Hazal Yüksekaya	Dicle University	Turkey
Hilal Genç Çopur	Kırşehir Ahi Evran University	Turkey

Hülya Acar	Duzce University	Turkey
Ilhame Amiral	Duzce University	Turkey
İsmail Ekincioğlu	Dumlupinar University	Turkey
Jan Zemlicka	Charles University	Czech Republic
Kazım Özcan	Gazi University	Turkey
Kebire Hilal Ayvaci	Ordu University	Turkey
Kemal Aydın	Selcuk University	Turkey
M. Tamer Koşan	Gazi University	Turkey
Maja Nedović	Novi Sad University	Serbia
Melek Eriş Büyükkaya	Karadeniz Technical University	Turkey
Merve İnce	Gazi University	Turkey
Meryem Kaya	Gazi University	Turkey
Mesrur Ümit Bingöl	Ankara University	Turkey
Mihaela Ribicic Penava	Osijek University	Croatia
Mine Uysal	Erzincan Binali Yıldırım University	Turkey
Mithat Özçırkıcı	Necmettin Erbakan University	Turkey
Mithat Raşit Özçırkıcı	Necmettin Erbakan University	Turkey
Muhammet Yazıcı	Karadeniz Technical University	Turkey
Mustafa Özkan	Gazi University	Turkey
Neslihan Biricik	Gazi University	Turkey
Neslihan Yıldırım	Gazi University	Turkey
Nezaket Javanshir	Hacettepe University	Turkey
Nida Palamut Koşar	Gaziantep University	Turkey
Nigar İrem Aras	Ankara University	Turkey
Nilay Akgönüllü Pirim	MATDER	Turkey
Nizamettin Ufuk Bostan	Ankara University	Turkey
Nurettin Doğan	Selçuk University	Turkey
Nurhayat İspir	Gazi University	Turkey
Oğuz Alkış	Gazi University	Turkey
Ökkeş Erdem Tekerek		Turkey
Ömer Akın	TOBB Economy and Technology University	Turkey
Özge Kazar	Hitit University	Turkey
Özgür Taşdemir	Trakya University	Turkey
Rahime Dere Paçin	Alanya Alaaddin Keykubat University	Turkey

Rıdvan Yaprak	Karadeniz Technical University	Turkey
Rümeysa Sacide Altınkaya	Gazi University	Turkey
Saif Ullah	GC University	Pakistan
Samet Sarioğlu	Hacettepe University	Turkey
Sanem Akdeniz Şafak	Aksaray University	Turkey
Sare Şengül	Marmara University	Turkey
Selami Ercan	Gazi University	Turkey
Semih Çavuşoğlu	Tokat Gaziosmanpasa University	Turkey
Serdal Yazıcı	Gazi University	Turkey
Serkan Ayan	Bursa Technical University	Turkey
Seval Işık	Sivas Cumhuriyet University	Turkey
Şeyda Solmaz	Ankara University	Turkey
Şeyma Firdevs Korkmaz	Gazi University	Turkey
Sezer Erdem	Battalgazi Farabi Anatolian Imam Hatip High School	Turkey
Sezin Çit	Gazi University	Turkey
Shamyrat Ataev		Turkey
Solomon Anokye	Kwame Nkrumah University	South Africa
Soukaina Hadiri	Moulay Ismail University	Morocco
Şule Yüksel Güngör	Gazi University	Turkey
Tahir Feyzioğlu	MATDER	Turkey
Takwa Hamzaoui		Tunisia
Tuğba Uygun ,	Alanya Alaaddin Keykubat University	Turkey
Tuğçe Pekacar Çalıcı	Ankara University	Turkey
Tülay Yağmur	Aksaray University	Turkey
Volkan Mutlu	Recep Tayyip Erdoğan University	Turkey
Yakup Yıldırım	Near East University	Cyprus
Zeliha Özdemir	Selcuk University	Turkey
Zeynep Çakmak Gürel	Erzincan Binali Yıldırım University	Turkey
Züleyha Açelya Üçer	Dumlupınar University	Turkey

SPONSORS



MATDER



GAZI UNIVERSITY