

INTERVAL VALUED PENTAPARTITIONED NEUTROSOPHIC GRAPHS WITH AN APPLICATION TO MCDM

Said Broumi ^{1,2*}, D. Ajay ³, P. Chellamani ³, Lathamaheswari Malayalan ⁴,
Mohamed Talea ¹, Assia Bakali ⁵, Philippe Schweizer ⁶, Saeid Jafari ⁷

¹ Laboratory of Information Processing, Faculty of Science Ben M'Sik, University of Hassan II, Casablanca, Morocco

² Regional Center for the Professions of Education and Training (C.R.M.E.F), Casablanca-Settat, Morocco.

³ Department of Mathematics, Sacred Heart College (Autonomous), Tamilnadu, India

⁴ Department of Mathematics, Hindustan Institute of Technology & Science, Chennai, India

⁵ Ecole Royale Navale-Boulevard Sour Jdid, Morocco

⁶ Independent researcher, Switzerland

⁷ College of Vestsjaelland South Herrestarede 11, Denmark

Received: 20 June 2022

Accepted: 27 August 2022

First online: 03 October 2022

Original scientific paper

Abstract: The concept of interval valued pentapartitioned neutrosophic set is the extension of interval-valued neutrosophic set, quadripartitioned neutrosophic set, interval valued quadripartitioned neutrosophic set and pentapartitioned neutrosophic set. The powerful mathematical tool known as the interval valued pentapartitioned neutrosophic set divides indeterminacy into three separate components: unknown, contradiction, and ignorance. There are several applications for graph theory in everyday life, and it is a rapidly growing topic. The concept of an interval valued pentapartitioned neutrosophic set is used in graph theory. A decision-making method (multicriteria, MCDM) is proposed by using the developed Interval valued Pentapartitioned Neutrosophic set with a numerical illustration. In this paper, as an extension of interval valued neutrosophic graph theory, we introduce the notions of Interval-Valued Pentapartitioned Neutrosophic Graph (IVPPN-graph) with degree, size, and order of an IVPPN-graph.

Key words: Neutrosophic Set, Interval valued Pentapartitioned Neutrosophic sets, Neutrosophic Graph, IVPPN-Graph.

* Corresponding author

broumisaid78@gmail.com, S.broumi@flbenmsik.ma (S.B)

dajaypravin@gmail.com (A. Pravin), joshmani238@gmail.com (P. Chellamani),

lathamax@gmail.com (L. Karthik), taleamohamed@yahoo.fr (M. Talea), assiabakali@yahoo.fr

(A. Bakali), flippe2@gmail.com (P. Schweizer), jafaripersia@gmail.com (S. Jafari)

1. Introduction

Linguistic variables with uncertainties such as vagueness, imprecision and ambiguity are handled with a special type of mathematical tool known as fuzzy set theory (Zadeh, 1965). The contribution of research articles in the field of fuzzy sets in mathematics and wide range of applications are growing exponentially. The advantage of introducing Zadeh's fuzzy sets in the place of classical sets has been its accuracy and precision in theory and compatibility and efficiency in case of applications.

Fuzzy set has been extended to Intuitionistic fuzzy set (Atanasov, 1986), in which the membership and non-membership degree of the element of that set ranges between 0 and 1. Intuitionistic set was further extended to Interval valued fuzzy sets, Neutrosophic sets, Pythagorean sets and so on. (Smarandache, 1999 & 2020) presented a new type of set as an extension of intuitionistic fuzzy set which is called Neutrosophic set. A neutrosophic set is characterized by a truth membership degree (T), an indeterminacy membership degree (I) and a falsity membership degree (F) independently, which are within the real unit interval $]0-,1+[$ satisfying the condition that the total of the membership grades is within the range of 0 to 3. In (Wang, 2010) single valued neutrosophic set (SVNS) with set-theoretic operators was investigated by Wang et al. SVNSs have been developed into many new concepts and are applied in different disciplines (Broumi et al., 2016 & 2016a & 2016b & 2016c; Chatterjee et al., 2016 & 2016b). Pythagorean set (Yager, 2013). is an extension of the intuitionistic set in which the condition differs from the intuitionistic set that the sum of the squares the membership is less than 1.

By combining the idea of fuzzy graphs and Pythagorean neutrosophic set Ajay and Chellamani et al. presented Pythagorean neutrosophic graphs (Ajay & Chellamani, 2020) is a combination of the Pythagorean and Neutrosophic set in which the the sum of the squares of the membership, non-membership and indeterminacy membership lies in $[0,2]$ and further theoretical concepts were developed in (Ajay & Chellamani, , 2021; Chellamani & Ajay, 2021; Ajay et al., 2021 , 2022 ,2020a); Chellamani et al., 2021). The partition of indeterminacy function of the neutrosophic set into contradiction part and ignorance part is defined as the quadripartitioned single-valued neutrosophic set (Chatterjee et al., 2016, 2020)

(Quek et al, 2022) introduced the notion of pentapartitioned neutrosophic graphs (PPNGs), as an extended version of Single valued neutrosophic graphs. (Mallick & Pramanik, 2020) proposed the concept of interval-valued pentapartitioned neutrosophic sets (IVPPNSs) as a generalization of pentapartitioned neutrosophic sets in the spirit of interval-valued neutrosophic sets (Broumi, 2016) The concept of Interval valued pentapartitioned neutrosophic sets (IVPPNSs) allows a decision-making expert to represented the membership degree contradiction membership function $C_A(x)$, an unknown membership function $U_A(x)$ and a falsity membership degree of a set of options in terms of the interval; hence, the range of uncertain information they can describe is widened. Interval-valued pentapartitioned neutrosophic numbers (IVPPNNs) are the generalized version of pentapartitioned neutrosophic numbers, quadripartitioned neutrosophic numbers, interval-valued quadripartitioned neutrosophic numbers. IVPPNNs will help model problems with incomplete, uncertain and indeterminate information. And further theoretical concepts were developed in (Das et al, 2022 & 2022a; Pramanik, 2022). (Hussain et al, 2022) introduced the notion of quadripartitioned single-valued neutrosophic graphs

and developed some operations on it. The Cartesian product, cross product, lexicographic product, strong product and composition of quadripartitioned single-valued neutrosophic graph have been investigated. The proposed concepts are illustrated with an application in the climatic analysis of apple cultivation. (Kaufmann, 1973), based on fuzzy relation (Zadeh, 2020). developed the idea of fuzzy graphs (FGs). Later, (Rosenfeld, 1975) defined the basic properties of fuzzy relations which are generalized with fuzzy set as a base set and fuzzy analogues of graphic theoretical concepts like bridges and trees were established with their properties and further fuzzy graph concepts were developed and applied in the real life situations in (Mohamed et al., 2020; Mordeson & Chang-Shyh, 1994; Naz et al, 2017 & 2018; Quek, et al., 2018 ; Smarandache, 2013 ; Tan et al., 2021 ; Das et al., 2021 ; Majumder et al, 2021; Saha et al., 2022). Recent developments of the fuzzy extensions are developed as in (Kumar et al., 2019 ; Radha et al., 2021 ; Vellapandi & Gunasekaran, 2020 ; Das & Edalatpanah, 2020 ; Polymenis, 2021; Mao et al, 2020 ; Voskoglou & Broumi, 2022 ; Zhang et al., 2022 ; Al-Hamido, 2022). To the best of authors knowledge, a very less work is being done on the interval-valued pentapartitioned neutrosophic set (IVPPNS), so the present study define the operations on the IVPPNS and later on extend it to graphs environment. The major contributions in this work are explained as follows:

1. The notions of interval valued pentapartitioned neutrosophic Graphs (IVPPNGs) are introduced. This manuscript makes the first attempt in the literature about the concept in neutrosophic graphs.
2. In addition, the complete and strong IVPPNG are defined. The operations like a Cartesian product, cross product, lexicographic product, strong product and the composition of IVPPNGs with their properties are discussed.
3. In addition, the complete, strong and complement of IVPPNGs are defined.
4. We extend some of the basic properties for this PNFG along with few examples. We have developed a decision-making model using the introduced Interval valued Pentapartitioned Neutrosophic graphs and applied it for a numerical illustration. A list of contribution (Table 1) of authors is presented below.

Table 1. Contribution of authors to extension of neutrosophic graphs

Authors	Year	Contributions
Akram and Siddique	2017	Introduced neutrosophic competition graph
Broumi et al.	2018	Introduced Generalized neutrosophic graph
Das et al	2020	Introduced generalized neutrosophic competition graph
Ajay and Chellamani	2020	Pythagorean Neutrosophic Fuzzy Graphs
S. Satham Hussain	2022	Quadripartitioned Neutrosophic Soft Graphs
S. Satham Hussain	2022	Quadripartitioned Bipolar Single Valued Neutrosophic Graph
S. Satham Hussain	2022	Quadripartitioned Single-Valued Neutrosophic Graph
Quek et al.	2022	Pentapartitioned Single-Valued Neutrosophic Graph
Our approach	2022	Interval valued Pentapartitioned Single-Valued Neutrosophic Graph

2. Preliminaries

This section summarizes some basic concepts from the theory of IVNSs and the concept of IVPPNSs, which is the foundation for the concept of IVPPNGs. Further details on the NS, SVN and IVNS theories can be found in [3-6] Smarandache (1999) and (Wang et al. 2010), respectively.

Definition 2.1: [5] Let A and B be two SVNNSs over a universe Y.

(i) A is contained in B, if $T_A(y) \leq T_B(y)$, $I_A(y) \geq I_B(y)$, and $F_A(y) \geq F_B(y)$, for all $y \in Y$. This relationship is denoted as $A \subseteq B$ (1)

(ii) A and B are said to be equal if $A \subseteq B$ and $B \subseteq A$ (2)

(iii) $A^c = (y, (F_A(y), 1 - I_A(y), T_A(y)))$, for all $y \in Y$ (3)

(iv) $A \cup B = (y, (\max(T_A, T_B), \min(I_A, I_B), \min(F_A, F_B)))$, for all $y \in Y$. (4)

(v) $A \cap B = (y, (\min(T_A, T_B), \max(I_A, I_B), \max(F_A, F_B)))$, for all $y \in Y$. (5)

Definition 2.2: (Wang et al, 2010) An Interval valued neutrosophic Sets (IVNS) A in X is denoted by an interval truth-membership function $T_A(x)$, an interval indeterminacy-membership function $I_A(x)$, and an interval falsity membership function $F_A(x)$ for each point x in X, there are

$$\tilde{T}_A(x) = [T_A^L(x), T_A^U(x)] \subseteq [0,1], \quad (6)$$

$$\tilde{I}_A(x) = [I_A^L(x), I_A^U(x)] \subseteq [0,1], \quad (7)$$

$$\tilde{F}_A(x) = [F_A^L(x), F_A^U(x)] \subseteq [0,1]. \quad (8)$$

Therefore an IVNS A can be denoted as, $A = \{ \langle x, \tilde{T}_A(x), \tilde{I}_A(x), \tilde{F}_A(x) \rangle \mid x \in X \}$

Then the sum of $\tilde{T}_A(x)$, $\tilde{I}_A(x)$, $\tilde{F}_A(x)$ satisfies the condition

$$0 \leq T_A^U(x) + I_A^U(x) + F_A^U(x) \leq 3 \quad (9)$$

If the upper and lower ends of the interval values of $T_A(x)$, $I_A(x)$ and $F_A(x)$ in an IVNS are equal then IVNS reduces to the SVN.

Definition 2.3: (Wang et al, 2010) Let A and B be two IVNSs over a universe Y.

(i) A is contained in B, if $T_A^L(y) \leq T_B^L(y)$, $T_A^U(y) \leq T_B^U(y)$, $I_A^L(y) \geq I_B^L(y)$, $I_A^U(y) \geq I_B^U(y)$, and $F_A^L(y) \geq F_B^L(y)$, $F_A^U(y) \geq F_B^U(y)$, for all $y \in Y$. This relationship is denoted as

$$A \subseteq B. \quad (10)$$

(ii) A and B are said to be equal if $A \subseteq B$ and $B \subseteq A$. (11)

(iii) $A^c = (y, (F_A(y), 1 - I_A(y), T_A(y)))$, for all $y \in Y$. (12)

(iv) $A \cup B = (y, (\max(T_A, T_B), \min(I_A, I_B), \min(F_A, F_B)))$, for all $y \in Y$. (13)

(v) $A \cap B = (y, (\min(T_A, T_B), \max(I_A, I_B), \max(F_A, F_B)))$, for all $y \in Y$ (14)

Definition 2.4: (Chatterjee et al., 2016) **Quadripartitioned neutrosophic sets**

Let X be a universe. A Quadripartitioned neutrosophic set A with independent neutrosophic components on X is an object of the form

$$A = \{ \langle x, T_A(x), C_A(x), U_A(x), F_A(x) \rangle : x \in X \} \quad (15)$$

$$\text{and } 0 \leq T_A(x) + C_A(x) + U_A(x) + F_A(x) \leq 4 \quad (16)$$

Here, $T_A(x)$ is the truth membership, $C_A(x)$ is contradiction membership, $U_A(x)$ is ignorance membership and $F_A(x)$ is the false membership.

Definition 2.5: (Pramanik, 2022). **Interval quadripartitioned neutrosophic Sets**

An Interval quadripartitioned neutrosophic Sets (IQNS) A in X is denoted by truth-membership function $T_A(x)$, a contradiction membership function $C_A(x)$, an unknown membership function $U_A(x)$ and a falsity membership function $F_A(x)$, for each point x in X , there are

$$T_A(x) = [T_A^L(x), T_A^U(x)] \subseteq [0,1], C_A(x) = [C_A^L(x), C_A^U(x)] \subseteq [0,1],$$

$$U_A(x) = [U_A^L(x), U_A^U(x)] \subseteq [0,1], F_A(x) = [F_A^L(x), F_A^U(x)] \subseteq [0,1],$$

Therefore an IQNS A can be denoted as,

$$A = \{ \langle x, T_A(x), C_A(x), U_A(x), F_A(x) \rangle \mid x \in X \} \quad (17)$$

$$A = \left\{ \left\langle x, [T_A^L(x), T_A^U(x)], [C_A^L(x), C_A^U(x)], [U_A^L(x), U_A^U(x)], [F_A^L(x), F_A^U(x)] \right\rangle \mid x \in X \right\} \quad (18)$$

Then the sum of $T_A(x), C_A(x), U_A(x), F_A(x)$ satisfies the condition

$$0 \leq T_A^U(x) + C_A^U(x) + U_A^U(x) + F_A^U(x) \leq 4 \quad (19)$$

If the upper and lower bounds of the interval values of $T_A(x), C_A(x), U_A(x)$ and $F_A(x)$ in an IQNS are equal then IQNS reduces to the QSVNS.

Definition 2.6: (Mallick & Pramanik, 2020). **Pentapartitioned neutrosophic sets**

Let X be a universe. A Pentapartitioned neutrosophic set A with independent neutrosophic components on X is an object of the form

$$A = \{ \langle x, T_A(x), C_A(x), U_A(x), G_A(x), F_A(x) \rangle : x \in X \} \quad (20)$$

$$\text{and } 0 \leq T_A(x) + C_A(x) + U_A(x) + G_A(x) + F_A(x) \leq 5 \quad (21)$$

Here, $T_A(x)$ is the truth membership, $C_A(x)$ is contradiction membership, $U_A(x)$ is ignorance membership, $G_A(x)$ is ignorance membership, and $F_A(x)$ is the false membership.

Definition 2.7: (Das et al., 2022) **Interval Pentapartitioned neutrosophic sets**

An Interval Pentapartitioned neutrosophic Sets (IPNS) A in X is denoted by truth-membership function $T_A(x)$, a contradiction membership function $C_A(x)$, an unknown membership function $U_A(x)$, $G_A(x)$ is ignorance membership and a falsity membership function $F_A(x)$ for each point x in X , there are

$$T_A(x) = [T_A^L(x), T_A^U(x)] \subseteq [0,1], C_A(x) = [C_A^L(x), C_A^U(x)] \subseteq [0,1],$$

$$U_A(x) = [U_A^L(x), U_A^U(x)] \subseteq [0,1], G_A(x) = [G_A^L(x), G_A^U(x)] \subseteq [0,1],$$

$$F_A(x) = [F_A^L(x), F_A^U(x)] \subseteq [0,1],$$

Therefore an IPNS A can be denoted as,

$$A = \{ \langle x, T_A(x), C_A(x), U_A(x), G_A(x), F_A(x) \rangle \mid x \in X \} \quad (22)$$

$$A = \left\{ \left\langle x, [T_A^L(x), T_A^U(x)], [C_A^L(x), C_A^U(x)], [U_A^L(x), U_A^U(x)], [G_A^L(x), G_A^U(x)], [F_A^L(x), F_A^U(x)] \right\rangle \mid x \in X \right\}$$

Then the sum of $T_A(x)$, $C_A(x)$, $U_A(x)$, $G_A(x)$, $F_A(x)$ satisfies the condition

$$0 \leq T_A^U(x) + C_A^U(x) + U_A^U(x) + G_A^U(x) + F_A^U(x) \leq 5 \quad (23)$$

If the upper and lower bounds of the interval values of $T_A(x)$, $C_A(x)$, $U_A(x)$, $G_A(x)$ and $F_A(x)$ in an IPNS are equal then IPNS reduces to the PSVNS.

Definition 2.8: (Broumi et al., 2016) Let V be a set. Let $E \subseteq \{\{\mu, v\} : \mu, v \in V \text{ with } \mu \neq v\}$.

Let A be an IVNS on V and B be an IVNS on E with

$$T_B^l(\mu, v) \leq \min\{T_A^l(\mu), T_A^l(v)\}, T_B^u(\mu, v) \leq \min\{T_A^u(\mu), T_A^u(v)\} \quad (24)$$

$$I_B^l(\mu, v) \geq \max\{I_A^l(\mu), I_A^l(v)\}, I_B^u(\mu, v) \geq \max\{I_A^u(\mu), I_A^u(v)\} \quad (25)$$

$$F_B^l(\mu, v) \geq \min\{F_A^l(\mu), F_A^l(v)\}, F_B^u(\mu, v) \geq \min\{F_A^u(\mu), F_A^u(v)\} \text{ for all } \{\mu, v\} \in E. \text{ Then } G = (A, B, V, E) \text{ is said to be an Interval-valued neutrosophic graph.} \quad (26)$$

Definition 2.9: [22] Let V be a set. Let $E \subseteq \{\{v, \omega\} : v, \omega \in V' \text{ with } \mu \neq v\}$. Let A be a PPNS on V , and B be a PPNS on E , with $t_B(\mu, v) \leq \min\{t_A(\mu), t_A(v)\}$, $c_B(\mu, v) \geq \max\{c_A(\mu), c_A(v)\}$, $g_B(\mu, v) \geq \max\{g_A(\mu), g_A(v)\}$, $u_B(\mu, v) \geq \max\{u_A(\mu), u_A(v)\}$ and $f_B(\mu, v) \geq \max\{f_A(\mu), f_A(v)\}$ for all $\{\mu, v\} \in E$. Then we have the following:

(i) $G = (A, B, V, E)$ is said to be a pentapartitioned neutrosophic graph (PPNG).

(ii) Each $v \in V$ is said to be a vertex of G .

(iii) Each $\{\mu, v\} \in E$ is said to be an edge of G .

3. Interval Pentapartitioned neutrosophic graphs

In this section, first the concept of interval pentapartitioned neutrosophic graphs (IVPPNGs) is introduced. Then based on the definitions of IVNS, interval-valued neutrosophic graphs [24] and IVPPNS given in definition 2.6 and definition 2.7, respectively will be used to put forward the novel concept of IVPPNGs. The related properties pertaining to this concept will be subsequently investigated.

Definition 3.1. Let V be a set. Let $E \subseteq \{\{\mu, v\} : v, \mu \in V \text{ with } \mu \neq v\}$. Let A be an IVPPNS on V , and B be an IVPPNS on E , with

$$t_B^l(\mu, v) \leq \min\{t_A^l(\mu), t_A^l(v)\}, t_B^u(\mu, v) \leq \min\{t_A^u(\mu), t_A^u(v)\}, \quad (27)$$

$$c_B^l(\mu, v) \geq \max\{c_A^l(\mu), c_A^l(v)\}, c_B^u(\mu, v) \geq \max\{c_A^u(\mu), c_A^u(v)\}, \quad (28)$$

$$g_B^l(\mu, v) \geq \max\{g_A^l(\mu), g_A^l(v)\}, g_B^u(\mu, v) \geq \max\{g_A^u(\mu), g_A^u(v)\}, \quad (29)$$

$$u_B^l(\mu, v) \geq \max\{u_A^l(\mu), u_A^l(v)\}, u_B^u(\mu, v) \geq \max\{u_A^u(\mu), u_A^u(v)\}, \quad (30)$$

$$\text{and } f_B^l(\mu, v) \geq \max\{f_A^l(\mu), f_A^l(v)\}, f_B^u(\mu, v) \geq \max\{f_A^u(\mu), f_A^u(v)\}, \quad (31)$$

for all $\{\mu, v\} \in E$. Then we have the following:

(i) $G=(A, B, V, E)$ is said to be an interval pentapartitioned neutrosophic graph (IVPPNG). (32)

(ii) Each $v \in V$ is said to be a vertex of G . (33)

(iii) Each $\{\mu, v\} \in E$ is said to be an edge of G . (34)

Fig. 1 A graphical representation of the PPNG G .

Notation 3.1.1 Let $G = (A, B, V, E)$ be a IVPPNG. Denote $m_A : V \rightarrow [0, 1]^5$, where $m_A(v) = (\tilde{t}_A(v), \tilde{c}_A(v), \tilde{g}_A(v), \tilde{u}_A(v), \tilde{f}_A(v))$ for all $v \in V$. Denote $m_B : E \rightarrow \text{Int}([0, 1]^5)$, where $m_B(\mu, v) = (\tilde{t}_B(\mu, v), \tilde{c}_B(\mu, v), \tilde{g}_B(\mu, v), \tilde{u}_B(\mu, v), \tilde{f}_B(\mu, v))$ for all $\{\mu, v\} \in E$.

Example 3.2 Let $G = (A, B, V, E)$ be an IVPPNG with $V = \{v_1, v_2, v_3, v_4, v_5\}$ and $E = \{\{v_1, v_2\}, \{v_1, v_3\}, \{v_1, v_4\}, \{v_2, v_5\}, \{v_4, v_5\}\}$. Then we have the following (Fig. 1):




Figure 1. Interval Valued Pentapartitioned Neutrosophic Graph

$$\begin{aligned}
 m_A(v_1) &= ([.7, .9], [.6, .7], [.5, .7], [.4, .6], [.3, .5]), \\
 m_A(v_2) &= ([.6, .7], [.5, .8], [.3, .6], [.4, .9], [.1, .5]), \\
 m_A(v_3) &= ([.4, .8], [.5, .7], [.6, .9], [.7, .9], [.2, .5]), \\
 m_A(v_4) &= ([.3, .6], [.4, .7], [.5, .9], [.7, .9], [.2, .5]), \\
 m_B(v_5) &= ([.7, .9], [.6, .8], [.5, .7], [.3, .6], [.2, .5]), \\
 m_B(v_1 v_2) &= ([.6, .7], [.7, .9], [.5, .7], [.5, .9], [.4, .6]), \\
 m_B(v_1 v_3) &= ([.4, .8], [.6, .8], [.6, .9], [.4, .6], [.3, .6]), \\
 m_B(v_1 v_4) &= ([.3, .6], [.6, .8], [.5, .9], [.7, .9], [.3, .5]), \\
 m_B(v_2 v_5) &= ([.6, .7], [.6, .8], [.5, .7], [.4, .9], [.2, .6]), \\
 m_B(v_4 v_5) &= ([.3, .6], [.6, .8], [.5, .9], [.7, .9], [.3, .7]).
 \end{aligned}$$

Definition 3.3. Let $G = (A, B, V, E)$ and $H = (A', B', V', E')$ be two IVPPNGs that satisfies the following conditions:

(i) $V' \subseteq V$ (35)

$$(ii) E' \subseteq \{ \{v, \mu\} : v, \mu \in V' \text{ with } \mu \neq v \} \quad (36)$$

$$(iii) m_{A'}(v) = m_A(v), \text{ for all } v \in V' \subseteq V \quad (37)$$

$$(iv) m_{B'}(v, \mu) = m_B(v, \mu), \text{ for all } \{v, \mu\} \in E' \subseteq E. \quad (38)$$

Then, H is said to be a partial *interval valued pentapartitioned neutrosophic graph* (partial-IVPPNG) of G.

Definition 3.4. Let $G = (A, B, V, E)$ and $H = (A', B', V', E')$ be two IVPPNGs that satisfies the following conditions:

$$(i) V' \subseteq V \quad (39)$$

$$(ii) E' \subseteq \{ \{v, \omega\} : v, \omega \in V' \text{ with } \omega \neq v \} \quad (40)$$

$$(iii) t_{A'}^l(v) \leq t_A^l(v), t_{A'}^u(v) \leq t_A^u(v), c_{A'}^l(v) \geq c_A^l(v), c_{A'}^u(v) \geq c_A^u(v), g_{A'}^l(v) \geq g_A^l(v), g_{A'}^u(v) \geq g_A^u(v), u_{A'}^l(v) \geq u_A^l(v), u_{A'}^u(v) \geq u_A^u(v), f_{A'}^l(v) \geq f_A^l(v), f_{A'}^u(v) \geq f_A^u(v) \text{ for all } v \in V' \subseteq V \quad (41)$$

$$(iv) t_{B'}^l(v, \omega) \leq t_B^l(v, \omega), t_{B'}^u(v, \omega) \leq t_B^u(v, \omega), c_{B'}^l(v, \omega) \geq c_B^l(v, \omega), c_{B'}^u(v, \omega) \geq c_B^u(v, \omega), g_{B'}^l(v, \omega) \geq g_B^l(v, \omega), g_{B'}^u(v, \omega) \geq g_B^u(v, \omega), u_{B'}^l(v, \omega) \geq u_B^l(v, \omega), u_{B'}^u(v, \omega) \geq u_B^u(v, \omega), f_{B'}^l(v, \omega) \geq f_B^l(v, \omega), f_{B'}^u(v, \omega) \geq f_B^u(v, \omega), \text{ for all } \{v, \omega\} \in E' \subseteq E. \quad (42)$$

Then, H is said to be an *interval valued pentapartitioned neutrosophic subgraph* (IVPPNSG) of G.

Example 3.5. Let G_1 be an IVPPNG and H_1, H_2 be the partial-IVPPNG and IVPPNSG of G_1 , respectively. The graphical representation of G_1, H_1 and H_2 are shown in Figs. 2, 3 and 4, respectively. Here H_2 is a IVPPNSG of G_1 but not a partial-IVPPNG of G_1 .

$$m_A(v_1) = ([.7, .9], [.6, .7], [.5, .7], [.4, .6], [.3, .5]),$$

$$m_A(v_2) = ([.6, .7], [.5, .8], [.3, .6], [.4, .9], [.1, .5]),$$

$$m_A(v_3) = ([.4, .8], [.5, .7], [.6, .9], [.7, .9], [.2, .5]),$$

$$m_A(v_4) = ([.3, .6], [.4, .7], [.5, .9], [.7, .9], [.2, .5]),$$

$$m_B(v_5) = ([.7, .9], [.6, .8], [.5, .7], [.3, .6], [.2, .5]).$$




Figure 2. Interval Valued Pentapartitioned Neutrosophic Graph




Figure 3. Partial Interval Valued Pentapartitioned Neutrosophic Graph




Figure 4. Interval Valued Pentapartitioned Neutrosophic Subgraph

Definition 3.6. Let $G = (A, B, V, E)$ be an IVPPNG. Let $v_0, v_1, \dots, v_n \in V$, with $\{v_i, v_{i+1}\} \in E$ for all $0 < i \leq n-1$, and with $v_i \neq v_j$ for all $0 \leq i < j \leq n-1$. Then we have the following:

$P = (v_0, v_1, \dots, v_n)$ is said to be a an interval valued pentapartitioned neutrosophic path (IVPPNP) in G .

For each i , $\{v_i, v_{i+1}\}$ is said to be an *edge* of P .

n is said to be the *length* of P .

Definition 3.7. Let $G = (A, B, V, E)$ be an IVPPNG. Then G is said to be *connected* if there exists at least one $\{v, \omega\} \in E$ for all $v \in V$.

Definition 3.8. Let $G = (A, B, V, E)$ be an IVPPNG. Then $v \in V$ is said to be *isolated* if $\{v, \omega\} \notin E$ for all $\omega \in V \setminus \{v\}$. the IVPPNG has v_1 as the isolated vertex.

Definition 3.9. Let $G = (A, B, V, E)$ be an IVPPNG and let $v \in V$. The degree of v , written as $d(v)$, is defined as $d(v) = \sum_{\mu \in V \text{ with } \{v, \omega\} \in E} m_B(v, \omega)$.

Remark 3.9.1. It follows that $d(v) \in [0, 1]^5$.

Example 3.10. In Fig. 3 under Eg. 3.5, the degree of the vertices are as follows.

$d(v_1) = (0.5, 1.7, 1.5, 1.1, 1.2)$, $d(v_3) = (0.6, 1.7, 1.7, 1.2, 1.2)$, $d(v_4) = (0.5, 1.8, 1.6, 1.3, 1.6)$.

Definition 3.11. Let $G = (A, B, V, E)$ be an IVPPNG and let $P = (v_0, v_1, \dots, v_n)$ be an IVPPNP in G . The strength of P , denoted as $s(P)$, is defined as :

$$s(P) = \left(\left[s_{t_A^l}(P), s_{t_A^u}(P) \right], \left[s_{c_A^l}(P), s_{c_A^u}(P) \right], \left[s_{g_A^l}(P), s_{g_A^u}(P) \right], \left[s_{u_A^l}(P), s_{u_A^u}(P) \right], \left[s_{f_A^l}(P), s_{f_A^u}(P) \right] \right) \\ = (s_{\bar{t}}(P), s_{\bar{c}}(P), s_{\bar{g}}(P), s_{\bar{u}}(P), s_{\bar{f}}(P)) \quad (43)$$

where

$$s_{t_A^l}(P) = \min\{t_B^l(v_i, v_{i+1}) : 0 \leq i \leq n-1\}, s_{t_A^u}(P) = \min\{t_B^u(v_i, v_{i+1}) : 0 \leq i \leq n-1\}, \\ s_{c_A^l}(P) = \max\{c_B^l(v_i, v_{i+1}) : 0 \leq i \leq n-1\}, s_{c_A^u}(P) = \max\{c_B^u(v_i, v_{i+1}) : 0 \leq i \leq n-1\}, \\ s_{g_A^l}(P) = \max\{g_B^l(v_i, v_{i+1}) : 0 \leq i \leq n-1\}, s_{g_A^u}(P) = \max\{g_B^u(v_i, v_{i+1}) : 0 \leq i \leq n-1\}, \\ s_{u_A^l}(P) = \max\{u_B^l(v_i, v_{i+1}) : 0 \leq i \leq n-1\}, s_{u_A^u}(P) = \max\{u_B^u(v_i, v_{i+1}) : 0 \leq i \leq n-1\}, \text{ and} \\ s_{f_A^l}(P) = \max\{f_B^l(v_i, v_{i+1}) : 0 \leq i \leq n-1\}, s_{f_A^u}(P) = \max\{f_B^u(v_i, v_{i+1}) : 0 \leq i \leq n-1\}, \text{ Moreover, the strength of connectedness among the vertices } a, b \in V \text{ in } G, \text{ denoted as } r_G(a, b), \text{ is defined as:}$$

$$r_G(a, b) = (r_{\bar{t},G}(a, b), r_{\bar{c},G}(a, b), r_{\bar{g},G}(a, b), r_{\bar{u},G}(a, b), r_{\bar{f},G}(a, b)) \quad \text{With} \quad (44)$$

$$r_{\bar{t},G}(a, b) = \left[r_{t_{A,G}^l}(a, b), r_{t_{A,G}^u}(a, b) \right], \quad r_{\bar{c},G}(a, b) = \left[r_{c_{A,G}^l}(a, b), r_{c_{A,G}^u}(a, b) \right] \\ r_{\bar{g},G}(a, b) = \left[r_{g_{A,G}^l}(a, b), r_{g_{A,G}^u}(a, b) \right], \quad r_{\bar{u},G}(a, b) = \left[r_{u_{A,G}^l}(a, b), r_{u_{A,G}^u}(a, b) \right] \text{ and} \\ r_{\bar{f},G}(a, b) = \left[r_{f_{A,G}^l}(a, b), r_{f_{A,G}^u}(a, b) \right]$$

Where

$$r_{t_{A,G}^l}(a, b) = \max\{s_{t_A^l}(P) : P = (v_0, v_1, \dots, v_{n_P}) \text{ in } G \text{ with } v_0 = a \text{ and } v_{n_P} = b\}, \\ r_{t_{A,G}^u}(a, b) = \max\{s_{t_A^u}(P) : P = (v_0, v_1, \dots, v_{n_P}) \text{ in } G \text{ with } v_0 = a \text{ and } v_{n_P} = b\}, \\ r_{c_{A,G}^l}(a, b) = \min\{s_{c_A^l}(P) : P = (v_0, v_1, \dots, v_{n_P}) \text{ in } G \text{ with } v_0 = a \text{ and } v_{n_P} = b\},$$

$$\begin{aligned}
r_{c_A^u, G}(a, b) &= \min\{s_{c_A^u}(P) : P = (v_0, v_1, \dots, v_{nP}) \text{ in } G \text{ with } v_0 = a \text{ and } v_{nP} = b\}, \\
r_{g_A^l, G}(a, b) &= \min\{s_{g_A^l}(P) : P = (v_0, v_1, \dots, v_{nP}) \text{ in } G \text{ with } v_0 = a \text{ and } v_{nP} = b\}, \\
r_{g_A^u, G}(a, b) &= \min\{s_{g_A^u}(P) : P = (v_0, v_1, \dots, v_{nP}) \text{ in } G \text{ with } v_0 = a \text{ and } v_{nP} = b\}, \\
r_{u_A^l, G}(a, b) &= \min\{s_{u_A^l}(P) : P = (v_0, v_1, \dots, v_{nP}) \text{ in } G \text{ with } v_0 = a \text{ and } v_{nP} = b\}, \\
r_{u_A^u, G}(a, b) &= \min\{s_{u_A^u}(P) : P = (v_0, v_1, \dots, v_{nP}) \text{ in } G \text{ with } v_0 = a \text{ and } v_{nP} = b\}, \\
r_{f_A^l, G}(a, b) &= \min\{s_{f_A^l}(P) : P = (v_0, v_1, \dots, v_{nP}) \text{ in } G \text{ with } v_0 = a \text{ and } v_{nP} = b\}, \\
r_{f_A^u, G}(a, b) &= \min\{s_{f_A^u}(P) : P = (v_0, v_1, \dots, v_{nP}) \text{ in } G \text{ with } v_0 = a \text{ and } v_{nP} = b\}.
\end{aligned}$$

Definition 3.12. Let $G = (A, B, V, E)$ be an IVPPNG and let $\{v, \omega\}$ be an edge in G . Denote $G'_{\{v, \omega\}}$ as the partial-IVPPNG of G , in which $G'_{\{v, \omega\}} = (A', B', V', E')$ with $V = V'$ and $E' = \{\{v, \omega\}\}$. Then, $\{v, \omega\}$ is said to be an *interval valued pentapartitioned neutrosophic bridge* (IVPPNB) in G if at least one of the following conditions holds for some $a, b \in V$.

$$(i) \ r_{\bar{t}, G'_{\{v, \omega\}}}(a, b) < r_{\bar{t}, G}(a, b) \quad (45)$$

$$(ii) \ r_{\bar{c}, G'_{\{v, \omega\}}}(a, b) > r_{\bar{c}, G}(a, b) \quad (46)$$

$$(iii) \ r_{\bar{g}, G'_{\{v, \omega\}}}(a, b) > r_{\bar{g}, G}(a, b) \quad (47)$$

$$(iv) \ r_{\bar{u}, G'_{\{v, \omega\}}}(a, b) > r_{\bar{u}, G}(a, b) \quad (48)$$

$$(v) \ r_{\bar{f}, G'_{\{v, \omega\}}}(a, b) > r_{\bar{f}, G}(a, b) \quad (49)$$

With $r_{t_A^l, G'_{\{v, \omega\}}}(a, b) < r_{t_A^l, G}(a, b)$, $r_{t_A^u, G'_{\{v, \omega\}}}(a, b) < r_{t_A^u, G}(a, b)$,
 $r_{c_A^l, G'_{\{v, \omega\}}}(a, b) > r_{c_A^l, G}(a, b)$, $r_{c_A^u, G'_{\{v, \omega\}}}(a, b) > r_{c_A^u, G}(a, b)$,
 $r_{g_A^l, G'_{\{v, \omega\}}}(a, b) > r_{g_A^l, G}(a, b)$, $r_{g_A^u, G'_{\{v, \omega\}}}(a, b) > r_{g_A^u, G}(a, b)$,
 $r_{u_A^l, G'_{\{v, \omega\}}}(a, b) > r_{u_A^l, G}(a, b)$, $r_{u_A^u, G'_{\{v, \omega\}}}(a, b) > r_{u_A^u, G}(a, b)$ and
 $r_{f_A^l, G'_{\{v, \omega\}}}(a, b) > r_{f_A^l, G}(a, b)$, $r_{f_A^u, G'_{\{v, \omega\}}}(a, b) > r_{f_A^u, G}(a, b)$.

In particular, if all of the conditions (i)–(v) are true for some $a, b \in V$, then $\{v, \omega\}$ is said to be a strong interval valued pentapartitioned neutrosophic bridge (strong-IVPPNB) in G .

Definition 3.13. Let $G = (A, B, V, E)$ be an IVPPNG and let v be a vertex in G . Denote G'_v as the partial-IVPPNG of G , in which $G'_v = (A', B', V', E')$ with $V' = V - \{v\}$ and $E' = E - \{\{a, v\} : a \in V - \{v\}\}$. Then, v is said to be an interval valued pentapartitioned neutrosophic cut vertex (IVPPNCV) in G if at least one of the following conditions holds for some $a, b \in V$

$$(i) \ r_{\bar{t}, G'_v}(a, b) < r_{\bar{t}, G}(a, b) \quad (50)$$

$$(ii) \ r_{\bar{c}, G'_v}(a, b) > r_{\bar{c}, G}(a, b) \quad (51)$$

$$(iii) \ r_{\bar{g}, G'_v}(a, b) > r_{\bar{g}, G}(a, b) \quad (52)$$

$$(iv) r_{\bar{u},G'_v}(a, b) > r_{\bar{u},G}(a, b) \quad (53)$$

$$(v) r_{\bar{f},G'_v}(a, b) > r_{\bar{f},G}(a, b) \quad (54)$$

With $r_{t_{A,G'_v}}^l(a, b) < r_{t_{A,G}}^l(a, b)$, $r_{t_{A,G'_v}}^u(a, b) < r_{t_{A,G}}^u(a, b)$,
 $r_{c_{A,G'_v}}^l(a, b) > r_{c_{A,G}}^l(a, b)$, $r_{c_{A,G'_v}}^u(a, b) > r_{c_{A,G}}^u(a, b)$,
 $r_{g_{A,G'_v}}^l(a, b) > r_{g_{A,G}}^l(a, b)$, $r_{g_{A,G'_v}}^u(a, b) > r_{g_{A,G}}^u(a, b)$,
 $r_{u_{A,G'_v}}^l(a, b) > r_{u_{A,G}}^l(a, b)$, $r_{u_{A,G'_v}}^u(a, b) > r_{u_{A,G}}^u(a, b)$ and
 $r_{f_{A,G'_v}}^l(a, b) > r_{f_{A,G}}^l(a, b)$, $r_{f_{A,G'_v}}^u(a, b) > r_{f_{A,G}}^u(a, b)$.

In particular, if all of the conditions (i)–(v) are true for some $a, b \in V$, then v is said to be a strong interval valued pentapartitioned neutrosophic cut vertex (strong-IVPPNCV) in G .

Definition 3.15. Suppose that $\hat{G} = (P, Q)$ is an SVPN-graph. Then, the size of $\hat{G} = (P, Q)$, denoted by $S(\hat{G})$ is defined by $S(\hat{G}) = (S_T(\hat{G}), S_C(\hat{G}), S_R(\hat{G}), S_U(\hat{G}), S_F(\hat{G}))$, where

$$S_T(\hat{G}) = \sum_{u \neq k} T_Q(u, k) \text{ denotes the T-size of } \hat{G} = (P, Q); \quad (55)$$

$$S_C(\hat{G}) = \sum_{u \neq k} C_Q(u, k) \text{ denotes the C-size of } \hat{G} = (P, Q); \quad (56)$$

$$S_R(\hat{G}) = \sum_{u \neq k} R_Q(u, k) \text{ denotes the R-size of } \hat{G} = (P, Q); \quad (57)$$

$$S_U(\hat{G}) = \sum_{u \neq k} U_Q(u, k) \text{ denotes the U-size of } \hat{G} = (P, Q); \quad (58)$$

$$S_F(\hat{G}) = \sum_{u \neq k} F_Q(u, k) \text{ denotes the F-size of } \hat{G} = (P, Q). \quad (59)$$

4. Application in Decision Making Problem

Rapid development of the application of the fuzzy decision making in the real-life application is remarkable. There are many applications of the implementation of the fuzzy graph model in decision making is evident in the recent research articles. Multi attribute decision-making is one of the decision-making methods in which we analyze the attributes according to the criteria for the problems chosen and using an algorithm the attributes are evaluated and the best attributes will be selected. More models are proposed according to each further developments and extensions of the fuzzy theory, we have proposed the decision-making method according to the developed concept of interval valued Pentapartitioned Neutrosophic graphs. The proposed algorithm using the interval valued Pentapartitioned Neutrosophic graphs is as follows:

4.1. Algorithm:

Step 1: Input the attributes $A = \{A_1, A_2, \dots, A_m\}$ and set of criteria $C = \{C_1, C_2, \dots, C_n\}$.

Step 2: Construct the interval valued Pentapartitioned relation $K^i = (k_{lp}^{(i)})_{mxm}$ where $i = 1, 2, \dots, n$ & $k_{lp}^i = ([T_{lp}^{(i)L}, T_{lp}^{(i)U}], [C_{lp}^{(i)L}, C_{lp}^{(i)U}], [U_{lp}^{(i)L}, U_{lp}^{(i)U}], [G_{lp}^{(i)L}, G_{lp}^{(i)U}], [F_{lp}^{(i)L}, F_{lp}^{(i)U}])$, $l, p = 1, 2, \dots, n$.

Step 3: Compute the resultant adjacency matrix (K) for the attributes under the criteria using the intersection of all the interval valued Pentapartitioned relation (K^i) as given, $K = \bigcap_i K^i$

Step 4: Calculate the score value of resultant adjacency matrix K by using score function S_{ij} .

$$S_{lp} = \frac{T_{lp}^L + T_{lp}^U + (1 - C_{lp}^L) + (1 - C_{lp}^U) + (1 - U_{lp}^L) + (1 - U_{lp}^U) + (1 - F_{lp}^L) + (1 - F_{lp}^U)}{5} \quad (60)$$

Step 5: Calculate the choice values $A_h = \sum_j S_{hj}$ $h, j=1,2,...m$ of each alternative.

Step 6: The final decision is the A_j with maximum choice value.

Step 7: If the j has more than one maximum value, then any one may be chosen.

4.2. An Illustrative Example:

In the fuzzy set and its extensions, the membership and the other membership degree function is sufficient but does not satisfy the needed vague categories. Thus, the introduction of the Pentapartitioned Neutrosophic set have paved the way for that advancement in the application area. It has five membership degree which mostly consider the vagueness of the real-life situations. The introduction of Interval valued Pentapartitioned Neutrosophic set is like a cream on top of the cake which adds more fuzziness in the conditions and criteria.

Now we consider the decision-making problems to choose the best international airline to take up the journey. The proposed decision-making method based on the interval valued Pentapartitioned Neutrosophic set and graph is used to solve this decision-making problem. The proposed method is more effective than the other proposed models due its membership degrees. Let the attributes be $A = \{A_1, A_2, ... A_6\}$ where A_i 's are the Airlines which we consider in our list. The selection of the best airline is based the criteria's $C = \{C_1, C_2, ... C_5\}$. where the criteria's are

C_1 = Safe air travel for passengers during COVID-19 Airline safety rating.

C_2 = High standards of Airport.

C_3 = Onboard product with excellent standards of staff service delivery

C_4 = Specialist Intelligence and Save time & Resources

C_5 = Trusted Experience and Efficient workflows

Using the proposed method, we input the attributes and the criteria's. According to each criteria Interval valued Pentapartitioned Neutrosophic relation is developed (K_i) as directed graphs. Then the intersection of all the criteria's are calculated and the resultant matrix is given in a graph. Then the score values are calculated for each alternative and framed into a graph structure. The choice values are found for each attribute and arranged according to find the optimal alternative. The method of solving the selected problem is as follows.:

Step 1: The attributes $A = \{A_1, A_2, ... A_6\}$ and criteria are $C = \{C_1, C_2, ... C_6\}$.

Step 2: The interval valued Pentapartitioned relation K^i ($i = 1, 2, 3, 4, 5$) is developed for each criteria as in the tables 2 to 6.

Table 2. Interval Valued Pentapartitioned Neutrosophic relation (K^1) for Criteria 1

K^1	A_1	A_2	A_3	A_4	A_5	A_6
A_1	$[[0.3,0.7],$ $[0.2,0.6],$ $[0.4,0.5],$ $[0.1,0.4],$ $[0.2,0.6]]$	$[[0.1,0.5],$ $[0.2,0.7],$ $[0.3,0.8],$ $[0.5,0.9],$ $[0.1,0.4]]$	$[[0.1,0.5],$ $[0.2,0.8],$ $[0.3,0.6],$ $[0.4,0.7],$ $[0.5,0.8]]$	$[[0.2,0.4],$ $[0.3,0.5],$ $[0.2,0.8],$ $[0.1,0.7],$ $[0.3,0.6]]$	$[[0.4,0.8],$ $[0.3,0.6],$ $[0.3,0.7],$ $[0.2,0.6],$ $[0.1,0.6]]$	$[[0.2,0.7],$ $[0.1,0.4],$ $[0.3,0.7],$ $[0.6,0.9],$ $[0.4,0.6]]$
A_2	$[[0.1,0.5],$ $[0.2,0.4],$ $[0.2,0.6],$ $[0.3,0.7],$ $[0.4,0.8]]$	$[[0.3,0.7],$ $[0.2,0.6],$ $[0.4,0.5],$ $[0.1,0.4],$ $[0.2,0.6]]$	$[[0.2,0.6],$ $[0.3,0.7],$ $[0.4,0.8],$ $[0.2,0.7],$ $[0.1,0.6]]$	$[[0.1,0.6],$ $[0.2,0.5],$ $[0.3,0.7],$ $[0.4,0.9],$ $[0.5,0.8]]$	$[[0.2,0.6],$ $[0.4,0.8],$ $[0.1,0.6],$ $[0.3,0.7],$ $[0.5,0.9]]$	$[[0.4,0.8],$ $[0.1,0.7],$ $[0.5,0.8],$ $[0.3,0.6],$ $[0.2,0.6]]$
A_3	$[[0.3,0.6],$ $[0.4,0.7],$ $[0.1,0.5],$ $[0.2,0.8],$ $[0.1,0.6]]$	$[[0.1,0.5],$ $[0.2,0.8],$ $[0.3,0.6],$ $[0.4,0.7],$ $[0.1,0.7]]$	$[[0.3,0.7],$ $[0.2,0.6],$ $[0.4,0.5],$ $[0.1,0.4],$ $[0.2,0.6]]$	$[[0.5,0.9],$ $[0.3,0.8],$ $[0.2,0.6],$ $[0.1,0.5],$ $[0.3,0.7]]$	$[[0.3,0.8],$ $[0.2,0.7],$ $[0.5,0.9],$ $[0.4,0.8],$ $[0.1,0.5]]$	$[[0.1,0.5],$ $[0.2,0.6],$ $[0.3,0.7],$ $[0.4,0.8],$ $[0.5,0.8]]$
A_4	$[[0.2,0.5],$ $[0.3,0.7],$ $[0.4,0.9],$ $[0.5,0.9],$ $[0.1,0.7]]$	$[[0.1,0.6],$ $[0.2,0.7],$ $[0.3,0.8],$ $[0.5,0.9],$ $[0.2,0.8]]$	$[[0.2,0.8],$ $[0.3,0.9],$ $[0.5,0.7],$ $[0.3,0.6],$ $[0.1,0.4]]$	$[[0.3,0.7],$ $[0.2,0.6],$ $[0.4,0.5],$ $[0.1,0.4],$ $[0.2,0.6]]$	$[[0.3,0.9],$ $[0.1,0.4],$ $[0.4,0.6],$ $[0.6,0.9],$ $[0.2,0.8]]$	$[[0.3,0.6],$ $[0.4,0.7],$ $[0.2,0.5],$ $[0.2,0.8],$ $[0.1,0.4]]$
A_5	$[[0.2,0.6],$ $[0.1,0.5],$ $[0.3,0.7],$ $[0.4,0.7],$ $[0.1,0.4]]$	$[[0.1,0.4],$ $[0.3,0.7],$ $[0.2,0.6],$ $[0.4,0.9],$ $[0.1,0.7]]$	$[[0.5,0.9],$ $[0.4,0.7],$ $[0.3,0.6],$ $[0.1,0.7],$ $[0.2,0.8]]$	$[[0.3,0.9],$ $[0.5,0.8],$ $[0.1,0.6],$ $[0.2,0.7],$ $[0.4,0.8]]$	$[[0.3,0.7],$ $[0.2,0.6],$ $[0.4,0.5],$ $[0.1,0.4],$ $[0.2,0.6]]$	$[[0.2,0.6],$ $[0.4,0.7],$ $[0.2,0.5],$ $[0.2,0.8],$ $[0.1,0.4]]$
A_6	$[[0.2,0.6],$ $[0.3,0.5],$ $[0.3,0.8],$ $[0.4,0.9],$ $[0.3,0.7]]$	$[[0.2,0.8],$ $[0.3,0.6],$ $[0.4,0.8],$ $[0.5,0.7],$ $[0.2,0.9]]$	$[[0.6,0.8],$ $[0.5,0.7],$ $[0.3,0.6],$ $[0.2,0.7],$ $[0.1,0.4]]$	$[[0.3,0.6],$ $[0.2,0.8],$ $[0.5,0.7],$ $[0.4,0.9],$ $[0.3,0.8]]$	$[[0.1,0.6],$ $[0.3,0.8],$ $[0.4,0.7],$ $[0.1,0.8],$ $[0.2,0.9]]$	$[[0.3,0.7],$ $[0.2,0.6],$ $[0.4,0.5],$ $[0.1,0.4],$ $[0.2,0.6]]$

Table 3. Interval Valued Pentapartitioned Neutrosophic relation (K^2) for Criteria 2

K^2	A_1	A_2	A_3	A_4	A_5	A_6
A_1	$[[0.2,0.5],$ $[0.3,0.7],$ $[0.4,0.9],$ $[0.5,0.9],$ $[0.1,0.7]]$	$[[0.1,0.5],$ $[0.2,0.7],$ $[0.3,0.8],$ $[0.5,0.9],$ $[0.1,0.4]]$	$[[0.1,0.5],$ $[0.2,0.8],$ $[0.3,0.6],$ $[0.4,0.7],$ $[0.1,0.7]]$	$[[0.1,0.6],$ $[0.2,0.7],$ $[0.3,0.8],$ $[0.5,0.9],$ $[0.2,0.8]]$	$[[0.1,0.4],$ $[0.3,0.7],$ $[0.2,0.6],$ $[0.4,0.9],$ $[0.1,0.7]]$	$[[0.2,0.8],$ $[0.3,0.6],$ $[0.4,0.8],$ $[0.5,0.7],$ $[0.2,0.9]]$
A_2	$[[0.1,0.6],$ $[0.2,0.7],$ $[0.3,0.8],$ $[0.5,0.9],$ $[0.2,0.8]]$	$[[0.2,0.5],$ $[0.3,0.7],$ $[0.4,0.9],$ $[0.5,0.9],$ $[0.1,0.7]]$	$[[0.2,0.6],$ $[0.3,0.7],$ $[0.4,0.8],$ $[0.2,0.7],$ $[0.1,0.5]]$	$[[0.2,0.7],$ $[0.3,0.5],$ $[0.4,0.8],$ $[0.5,0.9],$ $[0.4,0.6]]$	$[[0.2,0.6],$ $[0.4,0.8],$ $[0.1,0.6],$ $[0.3,0.7],$ $[0.5,0.9]]$	$[[0.4,0.8],$ $[0.1,0.7],$ $[0.5,0.8],$ $[0.3,0.6],$ $[0.2,0.6]]$
A_3	$[[0.2,0.8],$ $[0.3,0.9],$ $[0.5,0.7],$ $[0.2,0.6],$	$[[0.3,0.6],$ $[0.4,0.7],$ $[0.1,0.5],$ $[0.2,0.8],$	$[[0.2,0.5],$ $[0.3,0.7],$ $[0.4,0.9],$ $[0.5,0.9],$	$[[0.5,0.9],$ $[0.3,0.8],$ $[0.3,0.7],$ $[0.2,0.8],$	$[[0.4,0.7],$ $[0.5,0.8],$ $[0.3,0.6],$ $[0.1,0.4],$	$[[0.4,0.9],$ $[0.5,0.8],$ $[0.6,0.9],$ $[0.3,0.7],$

Interval Valued Pentapartitioned Neutrosophic Graphs with an Application to MCDM

	$[0.1,0.4]$	$[0.1,0.6]$	$[0.1,0.7]$	$[0.4,0.9]$	$[0.2,0.5]$	$[0.2,0.5]$
A ₄	$([0.3,0.7],$	$([0.1,0.5],$	$([0.5,0.9],$	$([0.2,0.5],$	$([0.3,0.8],$	$([0.2,0.6],$
	$[0.2,0.6],$	$[0.2,0.8],$	$[0.3,0.8],$	$[0.3,0.7],$	$[0.2,0.7],$	$[0.1,0.5],$
	$[0.4,0.5],$	$[0.3,0.6],$	$[0.2,0.6],$	$[0.4,0.9],$	$[0.5,0.9],$	$[0.3,0.7],$
	$[0.1,0.4],$	$[0.4,0.7],$	$[0.1,0.5],$	$[0.5,0.9],$	$[0.4,0.8],$	$[0.5,0.8],$
	$[0.2,0.6])]$	$[0.1,0.7])]$	$[0.3,0.7])]$	$[0.1,0.7])]$	$[0.2,0.5])]$	$[0.6,0.9])]$
A ₅	$([0.3,0.9],$	$([0.4,0.7],$	$([0.6,0.8],$	$([0.7,0.9],$	$([0.2,0.5],$	$([0.5,0.7],$
	$[0.1,0.4],$	$[0.2,0.6],$	$[0.6,0.9],$	$[0.5,0.8],$	$[0.3,0.7],$	$[0.4,0.6],$
	$[0.4,0.6],$	$[0.1,0.4],$	$[0.5,0.7],$	$[0.4,0.6],$	$[0.4,0.9],$	$[0.3,0.6],$
	$[0.6,0.9],$	$[0.3,0.8],$	$[0.4,0.6],$	$[0.3,0.5],$	$[0.5,0.9],$	$[0.2,0.5],$
	$[0.2,0.8])]$	$[0.5,0.8])]$	$[0.1,0.3])]$	$[0.2,0.7])]$	$[0.1,0.7])]$	$[0.1,0.6])]$
A ₆	$([0.3,0.6],$	$([0.4,0.7],$	$([0.5,0.8],$	$([0.3,0.5],$	$([0.4,0.6],$	$([0.2,0.5],$
	$[0.4,0.7],$	$[0.3,0.5],$	$[0.3,0.4],$	$[0.2,0.6],$	$[0.3,0.5],$	$[0.3,0.7],$
	$[0.2,0.5],$	$[0.2,0.6],$	$[0.4,0.7],$	$[0.5,0.8],$	$[0.2,0.8],$	$[0.4,0.9],$
	$[0.2,0.8],$	$[0.3,0.8],$	$[0.3,0.6],$	$[0.4,0.9],$	$[0.1,0.5],$	$[0.5,0.9],$
	$[0.1,0.4])]$	$[0.5,0.9])]$	$[0.2,0.6])]$	$[0.6,0.9])]$	$[0.6,0.9])]$	$[0.1,0.7])]$

Table 4. Interval Valued Pentapartitioned Neutrosophic relation (K^3) for Criteria 3

K^3	A ₁	A ₂	A ₃	A ₄	A ₅	A ₆
A ₁	$([0.3,0.6],$	$([0.5,0.9],$	$([0.1,0.5],$	$([0.3,0.7],$	$([0.4,0.6],$	$([0.5,0.7],$
	$[0.2,0.8],$	$[0.3,0.8],$	$[0.2,0.8],$	$[0.2,0.6],$	$[0.3,0.5],$	$[0.4,0.6],$
	$[0.5,0.7],$	$[0.2,0.6],$	$[0.3,0.6],$	$[0.4,0.5],$	$[0.2,0.8],$	$[0.3,0.6],$
	$[0.4,0.9],$	$[0.1,0.5],$	$[0.4,0.7],$	$[0.1,0.4],$	$[0.1,0.5],$	$[0.2,0.5],$
	$[0.3,0.8])]$	$[0.3,0.7])]$	$[0.1,0.7])]$	$[0.2,0.8])]$	$[0.6,0.9])]$	$[0.1,0.6])]$
A ₂	$([0.5,0.9],$	$([0.3,0.6],$	$([0.6,0.8],$	$([0.1,0.6],$	$([0.3,0.9],$	$([0.4,0.8],$
	$[0.3,0.8],$	$[0.2,0.8],$	$[0.6,0.9],$	$[0.2,0.7],$	$[0.1,0.4],$	$[0.1,0.7],$
	$[0.3,0.7],$	$[0.5,0.7],$	$[0.5,0.7],$	$[0.3,0.8],$	$[0.4,0.6],$	$[0.5,0.8],$
	$[0.2,0.8],$	$[0.4,0.9],$	$[0.4,0.6],$	$[0.5,0.9],$	$[0.6,0.9],$	$[0.3,0.6],$
	$[0.4,0.9])]$	$[0.3,0.8])]$	$[0.1,0.3])]$	$[0.2,0.5])]$	$[0.2,0.8])]$	$[0.2,0.6])]$
A ₃	$([0.3,0.9],$	$([0.4,0.7],$	$([0.3,0.6],$	$([0.5,0.8],$	$([0.1,0.4],$	$([0.3,0.6],$
	$[0.1,0.4],$	$[0.5,0.8],$	$[0.2,0.8],$	$[0.3,0.4],$	$[0.3,0.7],$	$[0.4,0.7],$
	$[0.4,0.6],$	$[0.3,0.6],$	$[0.5,0.7],$	$[0.4,0.7],$	$[0.2,0.6],$	$[0.2,0.5],$
	$[0.6,0.9],$	$[0.1,0.4],$	$[0.4,0.9],$	$[0.3,0.6],$	$[0.4,0.9],$	$[0.2,0.8],$
	$[0.2,0.8])]$	$[0.2,0.5])]$	$[0.3,0.8])]$	$[0.2,0.5])]$	$[0.1,0.7])]$	$[0.1,0.4])]$
A ₄	$([0.4,0.7],$	$([0.3,0.6],$	$([0.4,0.9],$	$([0.3,0.6],$	$([0.5,0.9],$	$([0.2,0.8],$
	$[0.2,0.6],$	$[0.4,0.7],$	$[0.5,0.8],$	$[0.2,0.8],$	$[0.3,0.8],$	$[0.3,0.6],$
	$[0.1,0.4],$	$[0.1,0.5],$	$[0.6,0.9],$	$[0.5,0.7],$	$[0.3,0.7],$	$[0.4,0.8],$
	$[0.3,0.8],$	$[0.2,0.8],$	$[0.5,0.7],$	$[0.4,0.9],$	$[0.2,0.8],$	$[0.5,0.7],$
	$[0.5,0.8])]$	$[0.1,0.6])]$	$[0.2,0.5])]$	$[0.3,0.8])]$	$[0.4,0.9])]$	$[0.2,0.9])]$
A ₅	$([0.6,0.8],$	$([0.3,0.7],$	$([0.1,0.5],$	$([0.2,0.5],$	$([0.3,0.6],$	$([0.7,0.9],$
	$[0.6,0.9],$	$[0.2,0.6],$	$[0.2,0.8],$	$[0.3,0.7],$	$[0.2,0.8],$	$[0.5,0.8],$
	$[0.5,0.7],$	$[0.4,0.5],$	$[0.3,0.6],$	$[0.4,0.9],$	$[0.5,0.7],$	$[0.4,0.6],$
	$[0.4,0.6],$	$[0.1,0.4],$	$[0.4,0.7],$	$[0.5,0.8],$	$[0.4,0.9],$	$[0.3,0.5],$
	$[0.1,0.3])]$	$[0.2,0.8])]$	$[0.1,0.7])]$	$[0.1,0.7])]$	$[0.3,0.8])]$	$[0.2,0.7])]$
A ₆	$([0.7,0.9],$	$([0.3,0.9],$	$([0.3,0.6],$	$([0.4,0.7],$	$([0.1,0.5],$	$([0.3,0.6],$
	$[0.5,0.8],$	$[0.1,0.4],$	$[0.4,0.7],$	$[0.2,0.6],$	$[0.2,0.7],$	$[0.2,0.8],$
	$[0.4,0.6],$	$[0.4,0.6],$	$[0.2,0.5],$	$[0.1,0.4],$	$[0.3,0.8],$	$[0.5,0.7],$
	$[0.3,0.5],$	$[0.6,0.9],$	$[0.2,0.8],$	$[0.3,0.8],$	$[0.5,0.9],$	$[0.4,0.9],$
	$[0.2,0.7])]$	$[0.2,0.8])]$	$[0.1,0.4])]$	$[0.5,0.8])]$	$[0.1,0.4])]$	$[0.3,0.8])]$

Table 5. Interval Valued Pentapartitioned Neutrosophic relation (K^4) for Criteria 4

K^4	A_1	A_2	A_3	A_4	A_5	A_6
A_1	[[0.3,0.5], [0.2,0.7], [0.5,0.9], [0.4,0.8], [0.3,0.8]]	[[0.5,0.9], [0.3,0.8], [0.3,0.7], [0.2,0.8], [0.4,0.9]]	[[0.3,0.9], [0.1,0.4], [0.4,0.6], [0.6,0.9], [0.2,0.8]]	[[0.4,0.7], [0.2,0.6], [0.1,0.4], [0.3,0.8], [0.5,0.8]]	[[0.6,0.8], [0.6,0.9], [0.5,0.7], [0.4,0.6], [0.1,0.3]]	[[0.7,0.9], [0.5,0.8], [0.4,0.6], [0.3,0.5], [0.2,0.7]]
A_2	[[0.5,0.9], [0.3,0.8], [0.2,0.6], [0.1,0.5], [0.3,0.7]]	[[0.3,0.5], [0.2,0.7], [0.5,0.9], [0.4,0.8], [0.3,0.8]]	[[0.4,0.7], [0.5,0.8], [0.3,0.6], [0.1,0.4], [0.2,0.5]]	[[0.3,0.6], [0.4,0.7], [0.1,0.5], [0.2,0.8], [0.1,0.6]]	[[0.3,0.7], [0.2,0.6], [0.4,0.5], [0.1,0.4], [0.2,0.8]]	[[0.3,0.9], [0.1,0.4], [0.4,0.6], [0.6,0.9], [0.2,0.8]]
A_3	[[0.1,0.5], [0.2,0.8], [0.3,0.6], [0.4,0.7], [0.1,0.7]]	[[0.6,0.8], [0.6,0.9], [0.5,0.7], [0.4,0.6], [0.1,0.3]]	[[0.3,0.5], [0.2,0.7], [0.5,0.9], [0.4,0.8], [0.3,0.8]]	[[0.4,0.9], [0.5,0.8], [0.6,0.9], [0.5,0.7], [0.2,0.5]]	[[0.1,0.3], [0.2,0.8], [0.3,0.7], [0.4,0.7], [0.1,0.5]]	[[0.3,0.6], [0.4,0.7], [0.2,0.5], [0.2,0.8], [0.1,0.4]]
A_4	[[0.3,0.7], [0.2,0.6], [0.4,0.5], [0.1,0.4], [0.2,0.8]]	[[0.1,0.6], [0.2,0.7], [0.3,0.8], [0.5,0.9], [0.2,0.5]]	[[0.5,0.8], [0.3,0.4], [0.4,0.7], [0.3,0.6], [0.2,0.5]]	[[0.3,0.5], [0.2,0.7], [0.5,0.9], [0.4,0.8], [0.3,0.8]]	[[0.2,0.5], [0.3,0.7], [0.4,0.9], [0.5,0.8], [0.1,0.6]]	[[0.4,0.7], [0.2,0.6], [0.1,0.4], [0.3,0.8], [0.5,0.8]]
A_5	[[0.4,0.6], [0.3,0.5], [0.2,0.8], [0.1,0.4], [0.6,0.9]]	[[0.3,0.9], [0.1,0.4], [0.4,0.6], [0.6,0.9], [0.2,0.8]]	[[0.1,0.4], [0.3,0.7], [0.2,0.6], [0.4,0.9], [0.1,0.7]]	[[0.5,0.9], [0.3,0.8], [0.3,0.7], [0.2,0.8], [0.4,0.6]]	[[0.3,0.5], [0.2,0.7], [0.5,0.9], [0.4,0.8], [0.3,0.8]]	[[0.2,0.7], [0.1,0.5], [0.3,0.8], [0.5,0.9], [0.1,0.4]]
A_6	[[0.5,0.7], [0.4,0.6], [0.3,0.6], [0.2,0.5], [0.1,0.6]]	[[0.4,0.8], [0.1,0.7], [0.5,0.8], [0.3,0.6], [0.2,0.5]]	[[0.3,0.6], [0.4,0.7], [0.2,0.5], [0.2,0.8], [0.1,0.4]]	[[0.2,0.8], [0.3,0.6], [0.4,0.8], [0.5,0.7], [0.2,0.9]]	[[0.7,0.9], [0.5,0.8], [0.4,0.6], [0.3,0.5], [0.2,0.7]]	[[0.3,0.5], [0.2,0.7], [0.5,0.9], [0.4,0.8], [0.3,0.8]]

Table 6. Interval Valued Pentapartitioned Neutrosophic relation (K^5) for Criteria 5

K^5	A_1	A_2	A_3	A_4	A_5	A_6
A_1	[[0.5,0.9], [0.3,0.8], [0.4,0.7], [0.2,0.6], [0.3,0.5]]	[[0.3,0.7], [0.2,0.5], [0.1,0.4], [0.4,0.8], [0.3,0.6]]	[[0.1,0.5], [0.2,0.8], [0.3,0.6], [0.4,0.7], [0.2,0.6]]	[[0.3,0.7], [0.2,0.6], [0.4,0.5], [0.1,0.4], [0.2,0.8]]	[[0.4,0.6], [0.3,0.5], [0.2,0.8], [0.1,0.4], [0.6,0.9]]	[[0.5,0.7], [0.4,0.6], [0.3,0.6], [0.2,0.5], [0.1,0.6]]
A_2	[[0.4,0.8], [0.3,0.5], [0.4,0.6], [0.2,0.5], [0.3,0.9]]	[[0.5,0.9], [0.3,0.8], [0.4,0.7], [0.2,0.6], [0.3,0.5]]	[[0.6,0.8], [0.6,0.9], [0.5,0.7], [0.4,0.6], [0.2,0.4]]	[[0.1,0.6], [0.2,0.7], [0.3,0.8], [0.5,0.9], [0.2,0.5]]	[[0.3,0.9], [0.1,0.4], [0.4,0.6], [0.6,0.9], [0.1,0.5]]	[[0.4,0.8], [0.1,0.7], [0.5,0.8], [0.3,0.6], [0.2,0.5]]
A_3	[[0.3,0.9], [0.1,0.4], [0.4,0.6], [0.6,0.9], [0.1,0.4]]	[[0.4,0.7], [0.5,0.8], [0.3,0.6], [0.4,0.7], [0.1,0.4]]	[[0.5,0.9], [0.3,0.8], [0.4,0.7], [0.2,0.6], [0.2,0.6]]	[[0.5,0.8], [0.3,0.4], [0.4,0.7], [0.3,0.6], [0.3,0.6]]	[[0.1,0.4], [0.3,0.7], [0.2,0.6], [0.4,0.9], [0.4,0.9]]	[[0.3,0.6], [0.4,0.7], [0.2,0.5], [0.2,0.8], [0.2,0.8]]

	$[0.2,0.8]$	$[0.2,0.5]$	$[0.3,0.5]$	$[0.2,0.5]$	$[0.1,0.5]$	$[0.1,0.4]$
A_4	$([0.4,0.7], [0.2,0.6], [0.1,0.4], [0.3,0.8], [0.5,0.8])$	$([0.3,0.6], [0.4,0.7], [0.1,0.5], [0.2,0.8], [0.1,0.6])$	$([0.4,0.9], [0.5,0.8], [0.6,0.9], [0.5,0.7], [0.2,0.5])$	$([0.5,0.9], [0.3,0.8], [0.4,0.7], [0.2,0.6], [0.3,0.5])$	$([0.5,0.9], [0.3,0.8], [0.3,0.7], [0.2,0.8], [0.4,0.6])$	$([0.2,0.8], [0.3,0.6], [0.4,0.8], [0.5,0.7], [0.2,0.9])$
A_5	$([0.6,0.8], [0.6,0.9], [0.5,0.7], [0.4,0.6], [0.1,0.3])$	$([0.3,0.7], [0.2,0.6], [0.4,0.5], [0.1,0.4], [0.2,0.8])$	$([0.1,0.3], [0.2,0.8], [0.3,0.7], [0.4,0.7], [0.6,0.9])$	$([0.2,0.5], [0.3,0.7], [0.4,0.9], [0.5,0.8], [0.1,0.6])$	$([0.5,0.9], [0.3,0.8], [0.4,0.7], [0.2,0.6], [0.3,0.5])$	$([0.7,0.9], [0.5,0.8], [0.4,0.6], [0.3,0.5], [0.2,0.7])$
A_6	$([0.7,0.9], [0.5,0.8], [0.4,0.6], [0.6,0.9], [0.2,0.8])$	$([0.3,0.9], [0.1,0.4], [0.4,0.6], [0.6,0.9], [0.2,0.8])$	$([0.3,0.6], [0.4,0.7], [0.2,0.5], [0.2,0.8], [0.1,0.4])$	$([0.4,0.7], [0.2,0.6], [0.1,0.4], [0.3,0.8], [0.5,0.9])$	$([0.2,0.7], [0.1,0.5], [0.3,0.8], [0.5,0.9], [0.1,0.4])$	$([0.5,0.9], [0.3,0.8], [0.4,0.7], [0.2,0.6], [0.3,0.5])$




Figure 5: Interval Valued Pentapartitioned Neutrosophic directed graph structure for $C_i (i = 1, 2, 3, 4, 5)$

Step 3: From the interval valued Pentapartitioned relation $K^i (i = 1, 2, 3, 4, 5)$, computing the intersection and find the resultant matrix as in Table 7.

Table 7. The resultant matrix

K	A ₁	A ₂	A ₃	A ₄	A ₅	A ₆
A ₁	[[0.2,0.5], [0.3,0.8], [0.5,0.9], [0.5,0.9], [0.3,0.8]]	[[0.1,0.5], [0.3,0.8], [0.3,0.8], [0.5,0.9], [0.4,0.9]]	[[0.1,0.5], [0.2,0.8], [0.4,0.6], [0.6,0.9], [0.5,0.8]]	[[0.1,0.4], [0.3,0.7], [0.4,0.8], [0.5,0.9], [0.5,0.8]]	[[0.1,0.4], [0.6,0.9], [0.5,0.8], [0.4,0.9], [0.6,0.9]]	[[0.2,0.7], [0.5,0.8], [0.4,0.8], [0.6,0.9], [0.4,0.9]]
A ₂	[[0.1,0.5], [0.3,0.8], [0.4,0.8], [0.5,0.9], [0.4,0.9]]	[[0.2,0.5], [0.3,0.8], [0.5,0.9], [0.5,0.9], [0.3,0.8]]	[[0.2,0.6], [0.6,0.9], [0.5,0.8], [0.4,0.7], [0.2,0.6]]	[[0.1,0.6], [0.3,0.8], [0.4,0.8], [0.5,0.9], [0.5,0.8]]	[[0.2,0.6], [0.4,0.8], [0.4,0.6], [0.6,0.9], [0.5,0.9]]	[[0.3,0.8], [0.1,0.7], [0.5,0.8], [0.6,0.9], [0.2,0.8]]
A ₃	[[0.1,0.5], [0.4,0.9], [0.5,0.7], [0.6,0.9], [0.2,0.8]]	[[0.1,0.5], [0.6,0.9], [0.5,0.7], [0.4,0.8], [0.2,0.7]]	[[0.2,0.5], [0.3,0.8], [0.5,0.9], [0.5,0.9], [0.3,0.8]]	[[0.4,0.8], [0.5,0.8], [0.6,0.9], [0.5,0.8], [0.4,0.9]]	[[0.1,0.3], [0.5,0.8], [0.5,0.9], [0.4,0.9], [0.2,0.7]]	[[0.1,0.5], [0.5,0.8], [0.6,0.9], [0.5,0.7], [0.5,0.8]]
A ₄	[[0.2,0.5], [0.3,0.7], [0.4,0.9], [0.5,0.9], [0.5,0.8]]	[[0.1,0.5], [0.4,0.8], [0.3,0.8], [0.5,0.9], [0.2,0.8]]	[[0.2,0.8], [0.5,0.9], [0.6,0.9], [0.5,0.7], [0.3,0.7]]	[[0.2,0.5], [0.3,0.8], [0.5,0.9], [0.5,0.9], [0.3,0.8]]	[[0.2,0.5], [0.3,0.8], [0.5,0.9], [0.6,0.9], [0.4,0.9]]	[[0.2,0.6], [0.4,0.7], [0.4,0.8], [0.5,0.8], [0.6,0.9]]
A ₅	[[0.2,0.6], [0.6,0.9], [0.5,0.8], [0.6,0.9], [0.6,0.9]]	[[0.1,0.4], [0.3,0.7], [0.4,0.6], [0.6,0.9], [0.5,0.8]]	[[0.1,0.3], [0.6,0.9], [0.5,0.7], [0.4,0.9], [0.6,0.9]]	[[0.2,0.5], [0.5,0.8], [0.4,0.9], [0.5,0.8], [0.4,0.8]]	[[0.2,0.5], [0.3,0.8], [0.5,0.9], [0.5,0.9], [0.3,0.8]]	[[0.2,0.6], [0.5,0.8], [0.4,0.8], [0.5,0.9], [0.2,0.7]]
A ₆	[[0.2,0.6], [0.5,0.8], [0.4,0.8], [0.6,0.9], [0.3,0.8]]	[[0.2,0.7], [0.3,0.7], [0.5,0.8], [0.6,0.9], [0.5,0.9]]	[[0.3,0.6], [0.5,0.7], [0.4,0.7], [0.3,0.8], [0.2,0.6]]	[[0.2,0.5], [0.3,0.8], [0.5,0.8], [0.5,0.9], [0.6,0.9]]	[[0.1,0.5], [0.5,0.8], [0.4,0.8], [0.5,0.9], [0.6,0.9]]	[[0.2,0.5], [0.3,0.8], [0.5,0.9], [0.5,0.9], [0.3,0.8]]

Step 4: By using the score function S_{ij} , in the resultant matrix of Table 7, find the score matrix as in Table 8.

Table 8. The score matrix of K and the choice values of the Attributes A_j

K	A ₁	A ₂	A ₃	A ₄	A ₅	A ₆	Choice Value
A ₁	0.74	0.74	0.76	0.72	0.58	0.72	4.24
A ₂	0.72	0.74	0.82	0.74	0.74	0.9	4.66
A ₃	0.72	0.76	0.74	0.68	0.7	0.66	4.26
A ₄	0.74	0.78	0.78	0.86	0.68	0.74	4.58
A ₅	0.6	0.74	0.58	0.72	0.74	0.76	4.14
A ₆	0.74	0.78	0.94	0.68	0.64	0.74	4.52




Figure 6. Interval Valued Pentapartitioned Neutrosophic directed graph with score values

Step 5: From the score matrix of Table 8, calculate the choice values for each attribute A_i ($i = 1$ to 6).

Step 6: Arrange the attributes according to their choice values in descending order, the final decision is A_2 .

$$4.66 > 4.58 > 5.52 > 4.26 > 4.24 > 4.14$$

$$A_2 > A_4 > A_6 > A_3 > A_1 > A_5$$

By using the proposed decision-making method, the optimal decision is decided. By using the method, the alternatives are arranged in descending order and the second airline is chosen as the best airline with the condition which satisfies all the criteria.

5. Discussion

Interval valued pentapartitioned neutrosophic sets provide a powerful tool to represent the information with imprecise and indeterminate data and have fruitful applications. Pentapartitioned neutrosophic sets is an extension of the Neutrosophic sets with the membership function classified into five categories. The interval valued pentapartitioned neutrosophic sets which is introduced is an advancement of the pentapartitioned set like each membership representation comes in an interval. In this

paper, the interval valued pentapartitioned neutrosophic sets have been implemented on the graph theoretical concepts and interval valued pentapartitioned neutrosophic graph have been developed. The significant operations of interval-valued pentapartitioned neutrosophic sets and its graphs are investigated. This helps the decision makers more sufficient for taking their input best suit to their domain of reference. The properties of interval valued pentapartitioned neutrosophic graph as cut vertex, bridge, degree are studied and examined with suitable examples. Hence, the proposed graphs and their basic properties have enough capabilities to address the related dependability on the indeterminate information. A decision-making method have been developed using the interval valued pentapartitioned neutrosophic graph with a numerical illustration.

6. Conclusion

Interval valued pentapartitioned neutrosophic graphs are generalization of pentapartitioned neutrosophic graphs and provide a sufficient space for complex decision-making situations. In this research paper some properties of interval valued pentapartitioned neutrosophic graph as cut vertex, bridge, degree are studied and examined with suitable examples. Using the proposed Interval valued Pentapartitioned Neutrosophic graphs, a decision-making method has been developed and applied in a real-life situation with numerical illustrations. In future, concepts can be developed in the interval valued pentapartitioned neutrosophic soft graphs, interval valued quadripartitioned neutrosophic graphs, Strong interval valued pentapartitioned neutrosophic graphs, etc. Also, one can extend the developed concepts into isomorphic properties and regularity properties in the proposed graph structures. The Interval valued pentapartitioned neutrosophic graph can be extended to regular and irregular interval valued pentapartitioned neutrosophic graph, interval-valued pentapartitioned neutrosophic intersection graphs, interval-valued pentapartitioned neutrosophic hypergraphs, and so on. The interval-valued pentapartitioned neutrosophic graph can be used in modeling the network, telecommunication, image processing, computer networks, expert systems...etc.

References:

- Ajay, D., & Chellamani, P. (2020). Pythagorean neutrosophic fuzzy graphs. *International Journal of Neutrosophic Science*, 11(2), 108-114. DOI: 10.5281/zenodo.4172124
- Ajay, D., & Chellamani, P. (2021). Pythagorean neutrosophic soft sets and their application to decision-making scenario. In *International Conference on Intelligent and Fuzzy Systems* (pp. 552-560). Springer, Cham. <https://doi.org/10.1007/978-3-030-85577-265>
- Ajay, D., Karthiga, S., & Chellamani, P. (2021a). A study on labelling of pythagorean neutrosophic fuzzy graphs. *Journal of Computational Mathematica*, 5(1), 105-116. <https://doi.org/10.26524/cm97>
- Ajay, D., Chellamani, P., Rajchakit, G., Boonsatit, N., & Hammachukiattikul, P. (2022). Regularity of Pythagorean neutrosophic graphs with an illustration in MCDM. *AIMS Mathematics*, 7(5), 9424-9442. doi: 10.3934/math.2022523

- Ajay, D., John Borg, S., & Chellamani, P. (2022a). Domination in Pythagorean Neutrosophic Graphs with an Application in Fuzzy Intelligent Decision Making. In *International Conference on Intelligent and Fuzzy Systems* (pp. 667-675). Springer, Cham. https://doi.org/10.1007/978-3-031-09176-6_74
- Al-Hamido, R. K. (2022). A New Neutrosophic Algebraic Structures. *Journal of Computational and Cognitive Engineering*. <https://doi.org/10.47852/bonviewJCCE2202213>
- Akram, M., & Siddique, S. (2017). Neutrosophic competition graphs with applications. *Journal of Intelligent & Fuzzy Systems*, 33(2), 921-935. DOI: 10.3233/JIFS-162207
- Atanasov, K. T. (1986). Intuitionistic fuzzy sets. *Fuzzy Sets and Systems*, 87-96. [https://doi.org/10.1016/S0165-0114\(86\)80034-3](https://doi.org/10.1016/S0165-0114(86)80034-3)
- Broumi, S., Talea, M., Bakali, A., & Smarandache, F. (2016). Single valued neutrosophic graphs. *Journal of New theory*, (10), 86-101.
- Broumi, S., Bakali, A., Talea, M., & Smarandache, F. (2016a). Isolated single valued neutrosophic graphs. *Neutrosophic Sets Syst* 11:74–78
- Broumi, S., Smarandache, F., Talea, M., & Bakali, A. (2016b). An introduction to bipolar single valued neutrosophic graph theory. In *Applied Mechanics and Materials*, 841, 184-191. DOI:10.4028/www.scientific.net/AMM.841.184
- Broumi, S., Talea, M., Bakali, A., & Smarandache, F. (2016c). On bipolar single valued neutrosophic graphs. *Journal of New Theory*, (11), 84-102.
- Broumi, S., Talea, M., Bakali, A., & Smarandache, F. (2016d). Interval valued neutrosophic graphs. *Critical Review*, XII, 2016, 5-33.
- Broumi, S., Talea, M., Bakali, A., Hassan, A., & Smarandache, F. (2017). Generalized Interval valued Neutrosophic graphs of first type. In *2017 IEEE International Conference on Innovations in Intelligent SysTems and Applications (INISTA)*, Gdynia Maritime University, Gdynia, Poland (Vol. 306). <http://doi.org/10.5281/zenodo.888864>
- Chatterjee, R., Majumdar, P., & Samanta, S. K. (2016). On some similarity measures and entropy on quadripartitioned single valued neutrosophic sets. *Journal of Intelligent & Fuzzy Systems*, 30(4), 2475-2485. DOI: 10.3233/JIFS-152017
- Chatterjee, R., Majumdar, P., & Samanta, S. K. (2016a). Interval-valued Possibility Quadripartitioned Single Valued Neutrosophic Soft Sets and some uncertainty based measures on them. *Neutrosophic Sets Syst Neutrosophic Sets and Systems*, vol. 14, , pp. 35-43. doi.org/10.5281/zenodo.570879.
- Chatterjee, R., Majumdar, P., & Samanta, S. K. (2016b). On some similarity measures and entropy on quadripartitioned single valued neutrosophic sets. *Journal of Intelligent & Fuzzy Systems*, 30(4), 2475-2485. DOI: 10.3233/JIFS-152017
- Chatterjee, R., Majumdar, P., & Samanta, S. K. (2020). A multi-criteria group decision making algorithm with quadripartitioned neutrosophic weighted aggregation operators using quadripartitioned neutrosophic numbers in IPQSVNSS environment. *Soft Computing*, 24(12), 8857-8880. DOI: 10.1007/s00500-019-04417-1
- Chellamani, P., & Ajay, D. (2021). Pythagorean neutrosophic Dombi fuzzy graphs with an application to MCDM. *Neutrosophic Sets and Systems*, 47, 411-431. [10.5281/zenodo.5775162](https://doi.org/10.5281/zenodo.5775162)

- Chellamani, P., Ajay, D., Broumi, S., & Ligori, T. (2021b). An approach to decision-making via picture fuzzy soft graphs. *Granular Computing*, 1-22. . <https://doi.org/10.1007/s41066-021-00282-2>
- Das, S., Das, R., & Pramanik, S. (2022). Single valued bipolar pentapartitioned neutrosophic set and its application in MADM strategy. *Neutrosophic Sets and Systems*, 49, 145-163. DOI: 10.5281/zenodo.6426381
- Das, S., Das, R., & Pramanik, S. (2022a). Single Valued Pentapartitioned Neutrosophic Graphs. *Neutrosophic Sets and Systems*, Vol. 50, pp. 225-238. DOI: 10.5281/zenodo.6774779
- Das, K., Samanta, S., & De, K. (2020). Generalized neutrosophic competition graphs. *Neutrosophic Sets and Systems*, 31, 156-171. 10.5281/zenodo.3639608
- Das, S. K., & Edalatpanah, S. A. (2020). A new ranking function of triangular neutrosophic number and its application in integer programming. *International Journal of Neutrosophic Science*, 4(2), 82-92. DOI: 10.5281/zenodo.3767107
- Das, S., Shil, B., & Pramanik, S. (2021). SVPNS-MADM strategy based on GRA in SVPNS Environment. *Neutrosophic Sets and Systems*, 47, 50-65. DOI: 10.5281/zenodo.5775091
- Hussain, S., Hussain, J., Rosyida, I., & Broumi, S. (2022). Quadripartitioned Neutrosophic Soft Graphs. In *Handbook of Research on Advances and Applications of Fuzzy Sets and Logic* (pp. 771-795). IGI Global. DOI:10.4018/978-1-7998-7979-4.ch034
- Hussain, S. S., Durga, N., Hossein, R., Ganesh, G., & Oscar, C. (2022a). New Concepts on Quadripartitioned Single-Valued Neutrosophic Graph with Real-Life Application. *International Journal of Fuzzy Systems*, 24(3), 1515-1529. <https://doi.org/10.1007/s40815-021-01205-8>
- Kumar, R., Edalatpanah, S. A., Jha, S., & Singh, R. (2019). A novel approach to solve gaussian valued neutrosophic shortest path problems. *International Journal of Engineering and Advanced Technology*, 2019, 8(3), pp. 347–353. 10.5281/zenodo.3047525
- Kauffman A (1973) *Introduction a la Theorie des Sous-emsembles Flous*, 1, Masson et Cie.
- Mao, X., Guoxi, Z., Fallah, M., & Edalatpanah, S. A. (2020). A neutrosophic-based approach in data envelopment analysis with undesirable outputs. *Mathematical problems in engineering*, 7626102. <https://doi.org/10.1155/2020/7626102>
- Mohamed, S. Y., & Ali, A. M. (2020). Energy of spherical fuzzy graphs. *Advances in Mathematics: Scientific Journal*, 9(1), 321-332. DOI:10.37418/amsj.9.1.26
- Mordeson, J. N., & Chang-Shyh, P. (1994). Operations on fuzzy graphs. *Information sciences*, 79(3-4), 159-170. [https://doi.org/10.1016/0020-0255\(94\)90116-3](https://doi.org/10.1016/0020-0255(94)90116-3)
- Mallick, R., & Pramanik, S. (2020). Pentapartitioned neutrosophic set and its properties. *Neutrosophic Sets and Systems*, 36, 184-192. DOI: 10.5281/zenodo.4065431
- Majumder, P., Das, S., Das, R., & Tripathy, B. C. (2021). Identification of the most significant risk factor of COVID-19 in economy using cosine similarity measure under SVPNS-environment. *Neutrosophic Sets and Systems*, *Neutrosophic Sets and Systems*, vol. 46, 2021, pp. 112-127. DOI: 10.5281/zenodo.5553497

- Naz, S., Rashmanlou, H., & Malik, M. A. (2017). Operations on single valued neutrosophic graphs with application. *Journal of Intelligent & Fuzzy Systems*, 32(3), 2137-2151. DOI: 10.3233/JIFS-161944
- Naz, S., Akram, M., & Smarandache, F. (2018). Certain notions of energy in single-valued neutrosophic graphs. *Axioms*, 7(3), 50. <https://doi.org/10.3390/axioms7030050>
- Pramanik, S. (2022). Interval quadripartitioned neutrosophic set, *Neutrosophic Sets and Systems*, 50. In press.
- Polymenis, A. (2021). A neutrosophic Student's-t-type of statistic for AR (1) random processes. *Journal of Fuzzy Extension and Applications*, 2(4), 388-393. 10.22105/JFEA.2021.287294.1149
- Quek, S. G., Broumi, S., Selvachandran, G., Bakali, A., Talea, M., & Smarandache, F. (2018). Some results on the graph theory for complex neutrosophic sets. *Symmetry*, 10(6), 190. <https://doi.org/10.3390/sym10060190>
- Quek, S. G., Selvachandran, G., Ajay, D., Chellamani, P., Taniar, D., Fujita, H., ... & Giang, N. L. (2022). New concepts of pentapartitioned neutrosophic graphs and applications for determining safest paths and towns in response to COVID-19. *Computational and Applied Mathematics*, 41(4), 1-27. <https://doi.org/10.1007/s40314-022-01823-4>
- Radha, R., & Arul Mary, A. S. (2021). Quadripartitioned neutrosophic pythagorean lie subalgebra. *Journal of Fuzzy Extension and Applications*, 2(3), 283-296. <https://doi.org/10.22105/jfea.2021.293069.1157>
- Saha, P., Majumder, P., Das, S., Das, P. K., & Tripathy, B. C. (2022). Single-valued pentapartitioned neutrosophic dice similarity measure and its application in the selection of suitable metal oxide nano-additive for biodiesel blend on environmental aspect. *Neutrosophic Sets and Systems*, Vol. 48, 2022, pp. 154-171, DOI: 10.5281/zenodo.6041439.
- Rosenfeld, A. (1975). Fuzzy graphs. In *Fuzzy Sets and Their Applications to Cognitive and Decision Processes*, Zadeh, L.A., Fu, K.S., Shimura, M., Eds.; Academic Press: Cambridge, MA, USA, 77–95. . <https://doi.org/10.1016/B978-0-12-775260-0.50008-6>
- Smarandache, F. (2013). n-Valued refined neutrosophic logic and its applications to physics. *Infinite Study*, 4, 143-146. DOI: 10.5281/zenodo.49149
- Smarandache, F. (1999). A unifying field in logics, neutrosophy: neutrosophic probability, set and logic. American Research Press, Rehoboth.
- Smarandache, F. (2022). Plithogeny, Plithogenic Set, Logic, Probability and Statistics: A Short Review. *Journal of Computational and Cognitive Engineering*, 47-50. <https://doi.org/10.47852/bonviewJCCE2202213>
- Tan, R. P., & Zhang, W. D. (2021). Decision-making method based on new entropy and refined single-valued neutrosophic sets and its application in typhoon disaster assessment. *Applied Intelligence*, 51(1), 283-307. <https://doi.org/10.1007/s10489-020-01706-3>
- Voskoglou, M., & Broumi, S. (2022). A hybrid method for the assessment of analogical reasoning skills. *Journal of fuzzy extension and applications*, 3(2), 152-157. 10.22105/JFEA.2022.342139.1220

Vellapandi, R., & Gunasekaran, S. (2020). A new decision making approach for winning strategy based on multi soft set logic. *Journal of fuzzy extension and applications*, 1(2), 119-129. [10.22105/JFEA.2020.249373.1013](https://doi.org/10.22105/JFEA.2020.249373.1013)

Wang, H., Smarandache, F., Zhang, Y. Q., & Sunderraman, R. (2010). Single valued neutrosophic sets. *Multispace Multistructure* 4:410–413

Yager, R.R. (2013). Pythagorean fuzzy subsets. In *Proceedings of the 2013 Joint IFSAWorld Congress and NAFIPS Annual Meeting (IFSA/NAFIPS)*, Edmonton, AB, Canada, 57–61. <https://doi.org/10.1109/IFSANAFIPS.2013.6608375>

Zhang, K., Xie, Y., Noorkhah, S. A., Imeni, M., & Das, S. K. (2022). Neutrosophic management evaluation of insurance companies by a hybrid TODIM-BSC method: a case study in private insurance companies. *Management Decision*. <https://doi.org/10.1108/MD-01-2022-0120>

Zadeh, L. A. (1965). Fuzzy sets. *Inf Control* 8, 38–353. <https://doi.org/10.2307/2272014>

Zadeh, L. A. (2020). Similarity relations and fuzzy orderings. *Inf. Sci.*, vol. 3, 177 – 200. [https://doi.org/10.1016/S0020-0255\(71\)80005-1](https://doi.org/10.1016/S0020-0255(71)80005-1)

© 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

