

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/350022240>

# On Some Algebraic Properties of $n$ -Refined Neutrosophic Elements and $n$ -Refined Neutrosophic Linear Equations

Article in *Mathematical Problems in Engineering* · March 2021

DOI: 10.1155/2021/5573072

CITATIONS

6

READS

82

1 author:



Mohammad Abobala

Tishreen University

33 PUBLICATIONS 260 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



RG Achievement [View project](#)



Contemporary Research [View project](#)

## Research Article

# On Some Algebraic Properties of $n$ -Refined Neutrosophic Elements and $n$ -Refined Neutrosophic Linear Equations

Mohammad Abobala 

Tishreen University, Faculty of Science, Department of Mathematics, Lattakia, Syria

Correspondence should be addressed to Mohammad Abobala; mohammadabobala777@gmail.com

Received 2 February 2021; Revised 19 February 2021; Accepted 3 March 2021; Published 12 March 2021

Academic Editor: R. Ali Borzooei

Copyright © 2021 Mohammad Abobala. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper studies the problem of determining invertible elements (units) in any  $n$ -refined neutrosophic ring. It presents the necessary and sufficient condition for any  $n$ -refined neutrosophic element to be invertible, idempotent, and nilpotent. Also, this work introduces some of the elementary algebraic properties of  $n$ -refined neutrosophic matrices with a direct application in solving  $n$ -refined neutrosophic algebraic equations.

## 1. Introduction

Neutrosophy is a new kind of generalized logic proposed by Smarandache [1]. It becomes a useful tool in many areas of science such as number theory [2, 3], solving equations [4], and medical studies [5, 6]. Also, we find many applications of neutrosophic structures in statistics [7], optimization [8], topology [9], and decision making [10, 11].

On the other hand, neutrosophic algebra began in [12], where Smarandache and Kandasamy defined concepts such as neutrosophic groups and neutrosophic rings. These notions were handled widely by Agboola et al. in [13, 14], where homomorphisms and AH-substructures were studied [15].

Recently, there is an increasing interest by the generalizations of neutrosophic algebraic structures. Smarandache and Abobala proposed  $n$ -refined neutrosophic rings [16], modules [17, 18], and spaces [19–22].

Neutrosophic algebraic equations are useful in many scientific areas; there is a full description of their solutions in neutrosophic fields and refined neutrosophic fields [23]. In particular, the relations between neutrosophic matrices and equations are defined in [24].

From this point of view, we are motivated to generalize the previous studies so that we study some of the algebraic properties of  $n$ -refined neutrosophic elements such as invertibility, nilpotency, and idempotency. Also, we study

elementary properties of  $n$ -refined neutrosophic matrices and their application in solving the  $n$ -refined neutrosophic linear system of equations as a new generalization of previous efforts in [23–25].

## 2. Preliminaries

*Definition 1* (see [16])

Let  $(R, +, \times)$  be a ring and  $I_k, 1 \leq k \leq n$  be  $n$  indeterminacies. We define  $R_n(I) = \{a_0 + a_1I + \dots + a_nI_n; a_i \in R\}$  to be  $n$ -refined neutrosophic ring. If  $n=2$ , we get a ring which is isomorphic to 2-refined neutrosophic ring  $R(I_1, I_2)$ .

Addition and multiplication on  $R_n(I)$  are defined as follows:

$$\begin{aligned} \sum_{i=0}^n x_i I_i + \sum_{i=0}^n y_i I_i &= \sum_{i=0}^n (x_i + y_i) I_i, \quad \sum_{i=0}^n x_i I_i \times \sum_{i=0}^n y_i I_i \\ &= \sum_{i,j=0}^n (x_i \times y_j) I_i I_j, \end{aligned} \quad (1)$$

where  $\times$  is the multiplication defined on the ring  $R$  and  $xI_0 = x$  for all  $x \in R$   $I_j I_i = I_i I_j = I_{\min(i,j)}, I_0 I_j = I_j$ .

It is easy to see that  $R_n(I)$  is a ring in the classical concept and contains a proper ring  $R$ .

**Definition 2** (see [16])

Let  $R_n(I)$  be an  $n$ -refined neutrosophic ring, and it is said to be commutative if  $xy = yx$  for each  $x, y \in R_n(I)$ ; if there is  $1 \in R_n(I)$  such  $1 \cdot x = x \cdot 1 = x$ , then it is called an  $n$ -refined neutrosophic ring with unity.

**Theorem 1** (see [16]). Let  $R_n(I)$  be an  $n$ -refined neutrosophic ring. Then,

- (a)  $R$  is commutative if and only if  $R_n(I)$  is commutative
- (b)  $R$  has unity if and only if  $R_n(I)$  has unity
- (c)  $R_n(I) = \sum_{i=0}^n RI_i = \{\sum_{i=0}^n x_i I_i; x_i \in R\}$

**Definition 3** (see [16])

- (a) Let  $R_n(I)$  be an  $n$ -refined neutrosophic ring and  $P = \sum_{i=0}^n P_i I_i = \{a_0 + a_1 I + \dots + a_n I_n; a_i \in P_i\}$  where  $P_i$  is a subset of  $R$ ; we define  $P$  to be an AH-subring if  $P_i$  is a subring of  $R$  for all  $i$ ; AHS-subring is defined by the condition  $P_i = P_j$  for all  $i, j$ .
- (b)  $P$  is an AH-ideal if  $P_i$  is a two-sided ideal of  $R$  for all  $i$ , and the AHS-ideal is defined by the condition  $P_i = P_j$  for all  $i, j$ .
- (c) The AH-ideal  $P$  is said to be null if  $P_i = R$  or  $P_i = \{0\}$  for all  $i$ .

**Definition 4** (see [16])

Let  $R_n(I)$  be an  $n$ -refined neutrosophic ring and  $P = \sum_{i=0}^n P_i I_i$  be an AH-ideal; we define AH-factor  $R(I)/P = \sum_{i=0}^n (R/P_i)I_i = \sum_{i=0}^n (x_i + P_i)I_i; x_i \in R$ .

**Theorem 2** (see [16])

Let  $R_n(I)$  be an  $n$ -refined neutrosophic ring and  $P = \sum_{i=0}^n P_i I_i$  be an AH-ideal;  
 $R_n(I)/P$  is a ring with the following two binary operations:

$$\begin{aligned} & \sum_{i=0}^n (x_i + P_i)I_i + \sum_{i=0}^n (y_i + P_i)I_i \\ &= \sum_{i=0}^n (x_i + y_i + P_i)I_i, \sum_{i=0}^n (x_i + P_i)I_i \times \sum_{i=0}^n (y_i + P_i)I_i \\ &= \sum_{i=0}^n (x_i \times y_i + P_i)I_i. \end{aligned} \quad (2)$$

### 3. Main Discussion

In this section, we study the invertibility of any element in any  $n$ -refined neutrosophic ring, and we show the conditions of idempotency and nilpotency in these rings. All rings in this section are considered with unity 1.

**Definition 5** Let  $X = A_0 + A_1 I_1 + \dots + A_n I_n$  be an  $n$ -refined neutrosophic element; we define its canonical sequence as follows:

$$\begin{aligned} M_0 &= A_0, \\ M_j &= A_0 + A_j + A_{j+1} + \dots + A_n, \quad 1 \leq j \leq n. \end{aligned} \quad (3)$$

For example,

$$M_3 = A_0 + A_3 + A_4 + \dots + A_n. \quad (4)$$

**Remark 1**

The multiplication operation between two  $n$ -refined neutrosophic elements can be represented by the following equation:

$$\begin{aligned} (A_0 + A_1 I_1 + \dots + A_n I_n)(B_0 + B_1 I_1 + \dots + B_n I_n) &= M_0 N_0 \\ &+ (M_n N_n - M_0 N_0)I_n + \sum_{i=1}^{n-1} (M_i N_i - M_{i+1} N_{i+1})I_i, \end{aligned} \quad (5)$$

where  $M_i$  and  $N_i$  are the canonical sequences of  $A_0 + A_1 I_1 + \dots + A_n I_n$  and  $B_0 + B_1 I_1 + \dots + B_n I_n$ , respectively.

**Proof.** For  $n = 0$ , the statement is true easily. Suppose that it is true for  $n = k$ , we must prove it for  $n = k + 1$ . We compute the multiplication  $L = (A_0 + A_1 I_1 + \dots + A_{k+1} I_{k+1})(B_0 + B_1 I_1 + \dots + B_{k+1} I_{k+1})$ .

$$\begin{aligned} & (A_0 + A_1 I_1 + \dots + A_{k+1} I_{k+1})(B_0 + B_1 I_1 + \dots + B_{k+1} I_{k+1}) = (A_0 + A_1 I_1 + \dots + A_k I_k)(B_0 + B_1 I_1 + \dots + B_k I_k) \\ & + A_{k+1} I_{k+1} (B_0 + B_1 I_1 + \dots + B_k I_k) + (A_0 + A_1 I_1 + \dots + A_k I_k) B_{k+1} I_{k+1} + A_{k+1} I_{k+1} B_{k+1} I_{k+1} \\ & = M_0 N_0 + (M_k N_k - M_0 N_0)I_k + \sum_{i=1}^k (M_i N_i - M_{i+1} N_{i+1})I_i + I_1 [A_{k+1} B_1 + A_1 B_{k+1}] \\ & + I_2 [A_{k+1} B_2 + A_2 B_{k+1}] + \dots + I_k [A_{k+1} B_k + A_k B_{k+1}] + I_{k+1} [A_0 B_{k+1} + A_{k+1} B_0 + A_{k+1} B_{k+1}]. \end{aligned} \quad (6)$$

Thus, the coefficient of  $I_{k+1}$  is  $A_0 B_{k+1} + A_{k+1} B_0 + A_{k+1} B_{k+1} = (A_{k+1} + A_0)(B_{k+1} + B_0) - (A_0)(B_0) = M_{k+1} N_{k+1} - M_0 N_0$ . Also, the coefficient of  $I_i$ ,  $1 \leq i \leq k$  is

$$\begin{aligned} M_i N_i - M_{i+1} N_{i+1} + A_{k+1} B_i + A_i B_{k+1} &= (A_0 + A_i + A_{i+1} + \dots + A_k) \\ & (B_0 + B_i + B_{i+1} + \dots + B_k) - (A_0 + A_{i+1} + A_{i+2} + \dots + A_k) \\ & (B_0 + B_{i+1} + B_{i+2} + \dots + B_k) + A_{k+1} B_i + A_i B_{k+1} = (A_0 + A_i + A_{i+1} \end{aligned}$$

$1 + \dots + A_k + A_{k+1}) (B_0 + B_i + B_{i+1} + \dots + B_k + B_{k+1}) - (A_0 + A_{i+1} + A_{i+2} + \dots + A_k + A_{k+1}) (B_0 + B_{i+1} + B_{i+2} + \dots + B_k + B_{k+1}) = M_i N_i - M_{i+1} N_{i+1}$ , where  $1 \leq i \leq k+1$ . Hence, our proof is completed by induction.

### Theorem 3

Let  $X = A_0 + A_1 I_1 + \dots + A_n I_n$  be an  $n$ -refined neutrosophic element, then it is invertible if and only if  $M_j, 0 \leq j \leq n$  are invertible. The inverse of  $X$  is  $X^{-1} = (M_0)^{-1} + (M_n^{-1} - M_0^{-1})I_n + \sum_{j=1}^{n-1} (M_j^{-1} - M_{j+1}^{-1})I_j = (A_0)^{-1} + ((A_0 + A_1 + \dots + A_n)^{-1} - (A_0 + A_2 + \dots + A_n)^{-1})I_1 + ((A_0 + A_2 + \dots + A_n)^{-1} - (A_0 + A_3 + \dots + A_n)^{-1})I_2 + ((A_0 + A_3 + \dots + A_n)^{-1} - (A_0 + A_4 + \dots + A_n)^{-1})I_3 + \dots + ((A_0 + A_n)^{-1} - (A_0)^{-1})I_n$ .

*Proof.*  $X$  is invertible if and only if there exists  $Y = B_0 + B_1 I_1 + \dots + B_n I_n$ , where  $XY = YX = 1$ . By using Remark 14, we can write the following:

$M_0 N_0 + (M_n N_n - M_0 N_0)I_n + \sum_{i=1}^{n-1} (M_i N_i - M_{i+1} N_{i+1})I_i = 1$ . This implies that  $M_0 N_0 = 1$  and  $M_i N_i - M_{i+1} N_{i+1} = 0$  for all  $i$ , where 0 is the zero element. Hence, we get  $M_i N_i = M_{i+1} N_{i+1} = M_0 N_0 = 1$ . So,  $M_j, 0 \leq j \leq n$  are invertible.

On the other hand, we put  $X^{-1} = (M_0)^{-1} + (M_n^{-1} - M_0^{-1})I_n + \sum_{j=1}^{n-1} (M_j^{-1} - M_{j+1}^{-1})I_j$ , and now we compute  $XX^{-1}$  as follows:

$$\begin{aligned} XX^{-1} &= M_0 M_0^{-1} + (M_1 M_1^{-1} - M_2 M_2^{-1})I_1 \\ &+ (M_2 M_2^{-1} - M_3 M_3^{-1})I_2 + \dots + (M_n M_n^{-1} - M_0 M_0^{-1})I_n = 1. \end{aligned} \quad (7)$$

### Example 1

Considering  $Z(I) = \{a + bI_1 + cI_2; a, b, c \in Z_2\}$  the 2-refined neutrosophic ring of integers, the set of invertible elements in  $Z_2$  is  $\{-1, 1\}$ . Hence, the set of all invertible elements in the corresponding 2-refined neutrosophic ring is  $\{1, -1, 1 - 2I_2, -1 + 2I_2, 1 - 2I_1, -1 + 2I_1, 1 + 2I_1 - 2I_2, -1 - 2I_1 + 2I_2\}$ .

### Theorem 4

Let  $X = A_0 + A_1 I_1 + \dots + A_n I_n$  be an  $n$ -refined neutrosophic element, and we have the following:

- (a)  $X$  is nilpotent if and only if  $M_j$  for all  $j$  are nilpotent
- (b)  $X$  is idempotent if and only if  $M_j$  for all  $j$  are idempotent

### Proof

- (a) First of all we will prove that  $X^r = M_0^r + I_n[(M_n)^r - (M_0)^r] + \sum_{i=1}^{n-1} ((M_i)^r - (M_{i+1})^r)I_i$ .

We use the induction, for  $r = 1$  it is clear. Suppose that it is true for  $r = k$ , we prove it for  $k + 1$ .

$$\begin{aligned} X^{k+1} &= X^k X = \left( M_0^k + I_n[(M_n)^k - (M_0)^k] + \sum_{i=1}^{n-1} ((M_i)^k - (M_{i+1})^k)I_i \right) (A_0 + A_1 I_1 + \dots + A_n I_n) \\ &= \left( M_0^k + I_n[(M_n)^k - (M_0)^k] + \sum_{i=1}^{n-1} ((M_i)^k - (M_{i+1})^k)I_i \right) \left( M_0 + (M_n - M_0)I_n + \sum_{i=1}^{n-1} (M_i - M_{i+1})I_i \right) \\ &= M_0^k M_0 + I_n[(M_n)^k M_n - M_0^k M_0] + \sum_{i=1}^{n-1} ((M_i^k M_i) - (M_{i+1}^k M_{i+1}))I_i \\ &= M_0^{k+1} + I_n[(M_n)^{k+1} - (M_0)^{k+1}] + \sum_{i=1}^{n-1} ((M_i)^{k+1} - (M_{i+1})^{k+1})I_i. \end{aligned} \quad (8)$$

$X$  is nilpotent if there is a positive integer  $r$  such that  $X^r = 0$ . This is equivalent to

$$\begin{aligned} M_0^r &= (M_n)^k \\ &= (M_j)^k \\ &= 0 \quad \text{for all } j, \text{ which implies the proof.} \end{aligned} \quad (9)$$

- (b) The proof is similar to (a).

## 4. $n$ -Refined Neutrosophic Linear Algebraic Equations

This section is dedicated to introduce an algorithm to solve  $n$ -refined neutrosophic linear equations over any  $n$ -refined neutrosophic field by turning them into classical systems of numbers.

Also, we discuss some elementary properties of  $n$ -refined neutrosophic matrices.

**Definition 6**

Let  $F_n(I)$  be any  $n$ -refined neutrosophic field. The  $n$ -refined linear neutrosophic equation with one variable over  $F_n(I)$  is defined as follows:

$$\begin{aligned} AX + B &= 0, \\ A, B, X &\in F_n(I), \end{aligned} \quad (10)$$

where

$$\begin{aligned} A &= a_0 + a_1 I_1 + \cdots + a_n I_n, \\ B &= b_0 + b_1 I_1 + \cdots + b_n I_n, \\ X &= x_0 + x_1 I_1 + \cdots + x_n I_n. \end{aligned} \quad (11)$$

**Theorem 5**

Let  $F_n(I)$  be any  $n$ -refined neutrosophic field and  $(*)AX + B = 0$  be any  $n$ -refined linear neutrosophic equation over  $F_n(I)$ . Then,  $(*)$  is solvable over  $F_n(I)$  if and only if the following classical system is solvable over the classical field  $F$ :

$$\begin{aligned} (1) \quad & a_0 x_0 + b_0 = 0 \\ (2) \quad & (a_0 + a_n)(x_0 + x_n) + (b_0 + b_n) = 0 \\ (3) \quad & (a_0 + a_n + a_{n-1})(x_0 + x_n + x_{n-1}) + (b_0 + b_n + b_{n-1}) = 0 \\ & \vdots \\ (n+1) \quad & (a_0 + a_1 + \cdots + a_n)(x_0 + x_1 + \cdots + x_n) + (b_0 + b_1 + \cdots + b_n) = 0 \end{aligned}$$

*Proof.* We will show that Equation (18) is equivalent to the previous classical system of equations.

We compute Equation (18) by using the canonical form, and we get

$$\begin{aligned} M_0 N_0 + (M_n N_n - M_0 N_0) I_n + \sum_{i=1}^{n-1} (M_i N_i - M_{i+1} N_{i+1}) I_i \\ = -b_0 - b_1 I_1 - \cdots - b_n I_n, \end{aligned} \quad (12)$$

where  $M_i$  and  $N_i$  are the canonical forms of  $A$  and  $X$ , respectively.

From (12), we get the following classical system:

$$\begin{aligned} M_0 N_0 &= -b_0, \\ M_n N_n - M_0 N_0 &= -b_n, \\ M_i N_i - M_{i+1} N_{i+1} &= -b_i, \quad \text{for all } 1 \leq i \leq n-1. \end{aligned} \quad (13)$$

The equation  $M_0 N_0 = -b_0$  equivalents  $a_0 x_0 + b_0 = 0$ . The equation  $M_n N_n - M_0 N_0 = -b_n$  equivalents  $(a_0 + a_n)(x_0 + x_n) + (b_0 + b_n) = 0$ .

Also, any equation with form  $M_i N_i - M_{i+1} N_{i+1} = -b_i$  for all  $1 \leq i \leq n-1$  equivalents  $(a_0 + a_n + a_{n-1} + \cdots + a_i)(x_0 + x_n + x_{n-1} + \cdots + x_i) + (b_0 + b_n + b_{n-1} + \cdots + b_i) = 0$  by mathematical induction; thus, our proof is complete.

Now, we can apply the previous theorem to solve  $n$ -refined neutrosophic linear equations, and we illustrate an example.

**Example 2**

Let  $R$  be the real field and  $R_3(I)$  be its corresponding 3-refined neutrosophic field. Consider the following 3-refined neutrosophic Equation (18)  $(1 + I_2 + I_3)X + (I_1 + 2I_2) = 0$ . To solve it, we turn it into the classical equivalent system.

- (1)  $1 \cdot x_0 + 0 = 0$ ; its solution  $x_0 = 0$ .
- (2)  $(1 + 1)(x_0 + x_3) + (0 + 0) = 0$ ; its solution is  $x_0 + x_3 = 0$ ; thus  $x_3 = 0$ .
- (3)  $(1 + 1 + 1)(x_0 + x_3 + x_2) + (0 + 0 + 2) = 0$ ; its solution is  $3(x_0 + x_3 + x_2) = -2$ ; thus  $x_2 = -2/3$ .
- (4)  $(1 + 1 + 1 + 0)(x_0 + x_3 + x_2 + x_1) + (0 + 1 + 2 + 0) = 0$ ; its solution is  $x_0 + x_3 + x_2 + x_1 = -1$ ; thus  $x_1 = -1/3$ .

Hence, the solution of Equation (18) is

$$X = -\frac{2}{3}I_2 - \frac{1}{3}I_1. \quad (14)$$

**Definition 7** Let  $A = \begin{pmatrix} a_{11} & \cdots & a_{1m} \\ \vdots & \ddots & \vdots \\ a_{n1} & \cdots & a_{nm} \end{pmatrix}$  be an  $m \times n$  matrix; if

$a_{ij} = x + yI_1 + zI_2 + \cdots + tI_n \in R_n(I)$ , then it is called an  $n$ -refined neutrosophic matrix, where  $R_n(I)$  is an  $n$ -refined neutrosophic ring.

**Remark 2**

If  $A$  is an  $m \times n$  matrix, then it can be represented as an element of the  $n$ -refined neutrosophic ring of matrices like the following:  $A = B + CI_1 + DI_2 + \cdots + KI_n$  where  $D, B, C, \dots, K$  are classical matrices with elements from the ring  $R$  and from size  $m \times n$ .

For example,  $A = \begin{pmatrix} 2 + I_1 + 3I_2 - I_3 & 1 - I_1 - I_2 \\ 3 + 4I_2 + 2I_3 & 1 + I_1 \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 3 & 1 \end{pmatrix} + \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} I_1 + \begin{pmatrix} 3 & -1 \\ 4 & 0 \end{pmatrix} I_2 + I_3 \begin{pmatrix} -1 & 0 \\ 2 & 0 \end{pmatrix}$  is a 3-

refined neutrosophic matrix.

**Remark 3**

The identity with respect to multiplication is the normal unitary matrix.

**Definition 8**

Let  $A$  be a square  $m \times m$   $n$ -refined neutrosophic matrix, then it is called invertible if there exists an  $n$ -refined square  $m \times m$  neutrosophic matrix  $B$  such that  $AB = U_{m \times m}$  where  $U_{m \times m}$  is the unitary classical matrix.

**Remark 4**

Let  $X = A_0 + A_1 I_1 + \cdots + A_n I_n$  be a square  $m \times m$   $n$ -refined neutrosophic matrix, then it is invertible if and only if  $M_j$ ,  $0 \leq j \leq n$  are invertible. The inverse of  $X$  is

$$\begin{aligned}
X^{-1} &= (M_0)^{-1} + (M_n^{-1} - M_0^{-1})I_n + \sum_{j=1}^{n-1} (M_j^{-1} - M_{j+1}^{-1})I_j \\
&= (A_0)^{-1} + ((A_0 + A_1 + \dots + A_n)^{-1} - (A_0 + A_2 + \dots + A_n)^{-1})I_1 \\
&\quad + ((A_0 + A_2 + \dots + A_n)^{-1} - (A_0 + A_3 + \dots + A_n)^{-1})I_2 \\
&\quad + ((A_0 + A_3 + \dots + A_n)^{-1} - (A_0 + A_4 + \dots + A_n)^{-1})I_3 + \dots + ((A_0 + A_n)^{-1} - (A_0)^{-1})I_n.
\end{aligned} \tag{15}$$

The proof holds directly as a special case of Theorem 3.

We defined the determinant of a square  $m \times m$   $n$ -refined neutrosophic matrix as

**Definition 9**

$$\begin{aligned}
\det X &= \det A_0 + [\det(A_0 + A_1 + \dots + A_n) - \det(A_0 + A_2 + \dots + A_n)]I_1 \\
&\quad + [\det(A_0 + A_2 + \dots + A_n) - \det(A_0 + A_3 + \dots + A_n)]I_2 + \dots \\
&\quad + [\det(A_0 + A_n) - \det(A_0)]I_n = \det(M_0) + (\det(M_n) - \det(M_0))I_n + \sum_{i=1}^{n-1} (\det(M_i) - \det(M_{i+1}))I_i.
\end{aligned} \tag{16}$$

This definition is supported by the condition of invertibility.

$$(c) \det X^{-1} = (\det X)^{-1}$$

**Theorem 6**

Let  $X = A_0 + A_1I_1 + \dots + A_nI_n$  be a square  $m \times m$   $n$ -refined neutrosophic matrix, and we have the following:

- (a)  $X$  is invertible if and only if  $\det X \neq 0$
- (b) If  $Y = B_0 + B_1I_1 + \dots + B_nI_n$  is a square  $m \times m$   $n$ -refined neutrosophic matrix, then  $\det XY = \det X \det Y$

*Proof.*

- (a) If  $\det X \neq 0$ , this will be equivalent to  $\det M_j \neq 0$  for all  $j$ , i.e.,  $M_j$  are invertible; thus,  $X$  is invertible according to Theorem 3.
- (b)  $XY = M_0N_0 + (M_nN_n - M_0N_0)I_n + \sum_{i=1}^{n-1} (M_iN_i - M_{i+1}N_{i+1})I_i$ . Hence,

$$\begin{aligned}
\det XY &= \det(M_0N_0) + I_n[\det(M_nN_n) - \det(M_0N_0)] \\
&\quad + \sum_{i=1}^{n-1} [(\det(M_iN_i) - \det(M_{i+1}N_{i+1}))I_i] \\
&= \det M_0 \det N_0 + I_n[\det(M_n) \det(N_n) - \det(M_0) \det(N_0)] \\
&\quad + \sum_{i=1}^{n-1} (\det(M_i) \det(N_i) - \det(M_{i+1}) \det(N_{i+1}))I_i \\
&= \left[ \det(M_0) + (\det(M_n) - \det(M_0))I_n + \sum_{i=1}^{n-1} (\det(M_i) - \det(M_{i+1}))I_i \right] \\
&\quad \cdot \left[ \det(N_0) + (\det(N_n) - \det(N_0))I_n + \sum_{i=1}^{n-1} (\det(N_i) - \det(N_{i+1}))I_i \right] = \det X \det Y.
\end{aligned} \tag{17}$$

(c) It holds directly from (b).

Now, we can find an easy algorithm to solve a linear system of  $n$ -refined neutrosophic algebraic equations over any  $n$ -refined neutrosophic field by using the inverse matrix method.

We construct an example.

**Example 3**

Consider the following system of 2-refined neutrosophic linear equations:

$$(2 + I_1 + 3I_2)X + (1 - I_1 - I_2)Y = -I_1, \tag{18}$$

$$(3 + 4I_2)X + (1 + I_1)Y = I_2. \tag{19}$$

The corresponding refined neutrosophic matrix is

$$A = \begin{pmatrix} 2 + I_1 + 3I_2 & 1 - I_1 - I_2 \\ 3 + 4I_2 & 1 + I_1 \end{pmatrix}.$$

We have the following:

$$(a) A = \begin{pmatrix} 2 + I_1 + 3I_2 & 1 - I_1 - I_2 \\ 3 + 4I_2 & 1 + I_1 \end{pmatrix} = \begin{pmatrix} 21 \\ 31 \end{pmatrix} + \begin{pmatrix} 1 - 1 \\ 0 \ 1 \end{pmatrix} I_1 + \begin{pmatrix} 3 - 1 \\ 4 \ 0 \end{pmatrix} I_2 \quad \text{where } B = \begin{pmatrix} 21 \\ 31 \end{pmatrix}, \quad C = \begin{pmatrix} 1 - 1 \\ 0 \ 1 \end{pmatrix}, \quad \text{and}$$

$$D = \begin{pmatrix} 3 - 1 \\ 4 \ 0 \end{pmatrix}, \quad B + D = \begin{pmatrix} 50 \\ 71 \end{pmatrix}, \quad B + C + D = \begin{pmatrix} 6 - 1 \\ 7 \ 2 \end{pmatrix}.$$

$$(b) B^{-1} = \begin{pmatrix} -11 \\ 3 - 2 \end{pmatrix}, \quad (B + D)^{-1} = \begin{pmatrix} 1/50 \\ -7/51 \end{pmatrix}, \quad (B + C + D)^{-1} = \begin{pmatrix} 2/19 \ 1/19 \\ -7/196/19 \end{pmatrix}.$$

$$(c) A^{-1} = B^{-1} + I_1 \quad [(B + C + D)^{-1} - (B + D)^{-1}] \quad + I_2$$

$$[(B + D)^{-1} - B^{-1}] = \begin{pmatrix} -1 \ 1 \\ 3 \ -2 \end{pmatrix} + I_1 \begin{pmatrix} -9/95 \ 1/19 \\ 98/95 - 13/19 \end{pmatrix} +$$

$$I_2 \begin{pmatrix} 6/5 \ -1 \\ -22/5 \ 3 \end{pmatrix} =$$

$$\begin{pmatrix} -1 - (9/95)I_1 + (6/5)I_2 & 1 + (1/19)I_1 - I_2 \\ 3 + (98/95)I_1 - (22/5)I_2 & -2 - (13/19)I_1 + 3I_2 \end{pmatrix}.$$

$$\text{It is easy to find that } A^{-1}A = AA^{-1} = \begin{pmatrix} 1 \ 0 \\ 0 \ 1 \end{pmatrix}.$$

$$(d) \det B = -1, \det (B + D) = 5, \det (B + C + D) = 19, \det A = -1 + I_1[19 - 5] + I_2[5 - (-1)] = -1 + 14I_1 + 6I_2.$$

Since  $A$  is invertible, we get the solution of the previous system of the 2-refined linear system by computing the product:

$$\begin{aligned} A^{-1} \begin{pmatrix} -I_1 \\ I_2 \end{pmatrix} &= \begin{pmatrix} -1 - \frac{9}{95}I_1 + \frac{6}{5}I_2 & 1 + \frac{1}{19}I_1 - I_2 \\ 3 + \frac{98}{95}I_1 - \frac{22}{5}I_2 & -2 - \frac{13}{19}I_1 + 3I_2 \end{pmatrix}, \\ \begin{pmatrix} -I_1 \\ I_2 \end{pmatrix} &= \begin{pmatrix} I_1 \left[ 1 + \frac{9}{95} - \frac{6}{5} + \frac{1}{19} \right] \\ I_1 \left[ -3 - \frac{98}{95} + \frac{22}{5} - \frac{13}{19} \right] + I_2[-2 + 3] \end{pmatrix}, \\ &= \begin{pmatrix} -I_1 \frac{1}{19} \\ -\frac{6}{19}I_1 + I_2 \end{pmatrix}. \end{aligned} \quad (20)$$

Thus,

$$X = \frac{1}{19}I_1, \quad (21)$$

$$Y = -\frac{6}{19}I_1 + I_2.$$

## 5. Conclusion

In this paper, we have determined the necessary and sufficient conditions for the invertibility, nilpotency, and idempotency of elements in a refined neutrosophic ring. In particular, we have studied some of algebraic properties of  $n$ -refined neutrosophic matrices such as determinants and inverses with an application solving the  $n$ -refined neutrosophic linear algebraic system of equations.

As a future research direction, we aim to study  $n$ -refined neutrosophic Diophantine equations.

## Data Availability

No data were used to support this study.

## Conflicts of Interest

The author declares that there are no conflicts of interest.

## References

- [1] F. Smarandache, “ $n$ -Valued refined neutrosophic logic and its applications in physics,” *Progress in Physics*, vol. 4, pp. 143–146, 2013.
- [2] H. Sankari and M. Abobala, “Neutrosophic linear diophantine equations with two variables,” *Neutrosophic Sets and Systems*, vol. 38, 2020.
- [3] M. Abobala, “Foundations of neutrosophic number theory,” *Neutrosophic Sets and Systems*, vol. 39, pp. 120–132, 2021.
- [4] S. A. Edalatpanah, “Systems of neutrosophic linear equations,” *Neutrosophic Sets and Systems*, vol. 33, pp. 92–104, 2020.
- [5] M. Abdel-Basset, M. Gunasekaran, G. Abdullah, and C. Victor, “A novel intelligent medical decision support model based on soft computing and IoT,” *IEEE Internet of Things Journal*, vol. 7, 2019.
- [6] M. Abdel-Basset, M. Mohamed, M. Elhoseny, L. H. Son, F. Chiclana, and A. E.-N. H. Zaied, “Cosine similarity measures of bipolar neutrosophic set for diagnosis of bipolar disorder diseases,” *Artificial Intelligence in Medicine*, vol. 101, Article ID 101735, 2019.
- [7] R. Alhabib and A. A. Salama, “The neutrosophic time series-study its models (Linear-Logarithmic) and test the coefficients significance of its linear model,” *Neutrosophic Sets and Systems*, vol. 33, pp. 105–115, 2020.
- [8] M. Abdel-Basset, R. Mohamed, A. E. N. H. Zaied, A. Gamal, and F. Smarandache, “Solving the supply chain problem using the best-worst method based on a novel Plithogenic model,” in *Optimization Theory Based on Neutrosophic and Plithogenic Sets*, pp. 1–19, Academic Press, Cambridge, MA, USA, 2020.
- [9] K. Ramesh, “Ngbr homomorphisms in neutrosophic topological spaces,” *Neutrosophic Sets and Systems*, vol. 32, pp. 25–37, 2020.
- [10] M. Abdel-Basset, A. Gamal, L. H. Son, and F. Smarandache, “A bipolar neutrosophic multi criteria decision making

- framework for professional selection,” *Applied Sciences*, vol. 10, no. 4, p. 1202, 2020.
- [11] A. Chakraborty, B. Banik, S. P. Mondal, and S. Alam, “Arithmetic and geometric operators of pentagonal neutrosophic number and its application in mobile communication service based MCGDM problem,” *Neutrosophic Sets and Systems*, vol. 32, pp. 61–79, 2020.
  - [12] V. W. B. Kandasamy and F. Smarandache, *Some Neutrosophic Algebraic Structures and Neutrosophic N-Algebraic Structures*, Hexis, Phonex, Arizona, 2006.
  - [13] A. A. A. Agboola, A. D. Akwu, and Y. T. Oyebo, “Neutrosophic groups and subgroups,” *International Journal of Mathematical Combinatorics*, vol. 3, pp. 1–9, 2012.
  - [14] T. Chalapathi and L. Madhavi, “Neutrosophic boolean rings,” *Neutrosophic Sets and Systems*, vol. 33, pp. 57–66, 2020.
  - [15] H. Sankari and M. Abobala, “AH-homomorphisms in neutrosophic rings and refined neutrosophic rings,” *Neutrosophic Sets and Systems*, vol. 38, 2020.
  - [16] F. Smarandache and M. Abobala, “ $n$ -Refined neutrosophic Rings,” *International Journal of Neutrosophic Science*, vol. 5, pp. 83–90, 2020.
  - [17] H. Sankari and M. Abobala, “ $n$ -Refined neutrosophic modules,” *Neutrosophic Sets and Systems*, vol. 36, pp. 1–11, 2020.
  - [18] M. Abobala, “Semi homomorphisms and algebraic relations between strong refined neutrosophic modules and strong neutrosophic modules,” *Neutrosophic Sets and Systems*, vol. 39, pp. 107–119, 2021.
  - [19] F. Smarandache and M. Abobala, “ $n$ -Refined neutrosophic vector spaces,” *International Journal of Neutrosophic Science*, vol. 7, pp. 47–54, 2020.
  - [20] H. Sankari and M. Abobala, “Solving three conjectures about neutrosophic quadruple vector spaces,” *Neutrosophic Sets and Systems*, vol. 38, 2020.
  - [21] M. A. Ibrahim, A. A. A. Agboola, B. S. Badmus, and S. A. Akinleye, “On refined neutrosophic vector spaces I,” *International Journal of Neutrosophic Science*, vol. 7, pp. 97–109, 2020.
  - [22] M. A. Ibrahim, A. A. A. Agboola, B. S. Badmus, and S. A. Akinleye, “On refined neutrosophic vector spaces II,” *International Journal of Neutrosophic Science*, vol. 9, pp. 22–36, 2020.
  - [23] M. Abobala, “On some neutrosophic algebraic equations,” *Journal of New Theory*, vol. 89, 2020.
  - [24] M. Abobala, “On refined neutrosophic matrices and their applications in refined neutrosophic algebraic equations,” *Journal of Mathematics*, vol. 2021, Article ID 5531093, 5 pages, 2021.
  - [25] R. Das, F. Smarandache, and B. Tripathy, “Neutrosophic fuzzy matrices and some algebraic operations,” *Neutrosophic Sets and Systems*, vol. 32, pp. 401–409, 2020.