

Neutrosophic quadruple algebraic hyperstructures

A. A. A. AGBOOLA, B. DAVVAZ, F. SMARANDACHE

Received 13 February 2017; Revised 23 February 2017; Accepted 12 March 2017

ABSTRACT. The objective of this paper is to develop neutrosophic quadruple algebraic hyperstructures. Specifically, we develop neutrosophic quadruple semihypergroups, neutrosophic quadruple canonical hypergroups and neutrosophic quadruple hyperrings and we present elementary properties which characterize them.

2010 AMS Classification: 03B60, 20A05, 20N20, 97H40

Keywords: Neutrosophy, Neutrosophic quadruple number, Neutrosophic quadruple semihypergroup, Neutrosophic quadruple canonical hypergroup, Neutrosophic quadruple hyperring.

Corresponding Author: A. A. A. Agboola (aaaola2003@yahoo.com)

1. INTRODUCTION

The concept of neutrosophic quadruple numbers was introduced by Florentin Smarandache [18]. It was shown in [18] how arithmetic operations of addition, subtraction, multiplication and scalar multiplication could be performed on the set of neutrosophic quadruple numbers. In [1], Akinleye et.al. introduced the notion of neutrosophic quadruple algebraic structures. Neutrosophic quadruple rings were studied and their basic properties were presented. In the present paper, two hyperoperations $\hat{+}$ and $\hat{\times}$ are defined on the neutrosophic set NQ of quadruple numbers to develop new algebraic hyperstructures which we call neutrosophic quadruple algebraic hyperstructures. Specifically, it is shown that $(NQ, \hat{\times})$ is a neutrosophic quadruple semihypergroup, $(NQ, \hat{+})$ is a neutrosophic quadruple canonical hypergroup and $(NQ, \hat{+}, \hat{\times})$ is a neutrosophic quadruple hyperring and their basic properties are presented.

Definition 1.1 ([18]). A neutrosophic quadruple number is a number of the form (a, bT, cI, dF) where T, I, F have their usual neutrosophic logic meanings and $a, b, c, d \in \mathbb{R}$ or \mathbb{C} . The set NQ defined by

$$(1.1) \quad NQ = \{(a, bT, cI, dF) : a, b, c, d \in \mathbb{R} \text{ or } \mathbb{C}\}$$

is called a neutrosophic set of quadruple numbers. For a neutrosophic quadruple number (a, bT, cI, dF) representing any entity which may be a number, an idea, an object, etc, a is called the known part and (bT, cI, dF) is called the unknown part.

Definition 1.2. Let $a = (a_1, a_2T, a_3I, a_4F), b = (b_1, b_2T, b_3I, b_4F) \in NQ$. We define the following:

$$(1.2) \quad a + b = (a_1 + b_1, (a_2 + b_2)T, (a_3 + b_3)I, (a_4 + b_4)F),$$

$$(1.3) \quad a - b = (a_1 - b_1, (a_2 - b_2)T, (a_3 - b_3)I, (a_4 - b_4)F).$$

Definition 1.3. Let $a = (a_1, a_2T, a_3I, a_4F) \in NQ$ and let α be any scalar which may be real or complex, the scalar product $\alpha.a$ is defined by

$$(1.4) \quad \alpha.a = \alpha.(a_1, a_2T, a_3I, a_4F) = (\alpha a_1, \alpha a_2T, \alpha a_3I, \alpha a_4F).$$

If $\alpha = 0$, then we have $0.a = (0, 0, 0, 0)$ and for any non-zero scalars m and n and $b = (b_1, b_2T, b_3I, b_4F)$, we have:

$$(m + n)a = ma + na,$$

$$m(a + b) = ma + mb,$$

$$mn(a) = m(na),$$

$$-a = (-a_1, -a_2T, -a_3I, -a_4F).$$

Definition 1.4 ([18]). [Absorbance Law] Let X be a set endowed with a total order $x < y$, named " x prevailed by y " or " x less stronger than y " or " x less preferred than y ". $x \leq y$ is considered as " x prevailed by or equal to y " or " x less stronger than or equal to y " or " x less preferred than or equal to y ".

For any elements $x, y \in X$, with $x \leq y$, absorbance law is defined as

$$(1.5) \quad x.y = y.x = \text{absorb}(x, y) = \max\{x, y\} = y$$

which means that the bigger element absorbs the smaller element (the big fish eats the small fish). It is clear from (1.5) that

$$(1.6) \quad x.x = x^2 = \text{absorb}(x, x) = \max\{x, x\} = x \quad \text{and}$$

$$(1.7) \quad x_1.x_2 \cdots x_n = \max\{x_1, x_2, \cdots, x_n\}.$$

Analogously, if $x > y$, we say that " x prevails to y " or " x is stronger than y " or " x is preferred to y ". Also, if $x \geq y$, we say that " x prevails or is equal to y " or " x is stronger than or equal to y " or " x is preferred or equal to y ".

Definition 1.5. Consider the set $\{T, I, F\}$. Suppose in an optimistic way we consider the prevalence order $T > I > F$. Then we have:

$$(1.8) \quad TI = IT = \max\{T, I\} = T,$$

$$(1.9) \quad TF = FT = \max\{T, F\} = T,$$

$$(1.10) \quad IF = FI = \max\{I, F\} = I,$$

$$(1.11) \quad TT = T^2 = T,$$

$$(1.12) \quad II = I^2 = I,$$

$$(1.13) \quad FF = F^2 = F.$$

71 Analogously, suppose in a pessimistic way we consider the prevalence order $T <$
72 $I < F$. Then we have:

$$73 \quad (1.14) \quad TI = IT = \max\{T, I\} = I,$$

$$74 \quad (1.15) \quad TF = FT = \max\{T, F\} = F,$$

$$75 \quad (1.16) \quad IF = FI = \max\{I, F\} = F,$$

$$76 \quad (1.17) \quad TT = T^2 = T,$$

$$77 \quad (1.18) \quad II = I^2 = I,$$

$$78 \quad (1.19) \quad FF = F^2 = F.$$

79 Except otherwise stated, we will consider only the prevalence order $T < I < F$
80 in this paper.

81 **Definition 1.6.** Let $a = (a_1, a_2T, a_3I, a_4F), b = (b_1, b_2T, b_3I, b_4F) \in NQ$. Then

$$\begin{aligned} 82 \quad a.b &= (a_1, a_2T, a_3I, a_4F).(b_1, b_2T, b_3I, b_4F) \\ 83 &= (a_1b_1, (a_1b_2 + a_2b_1 + a_2b_2)T, (a_1b_3 + a_2b_3 + a_3b_1 + a_3b_2 + a_3b_3)I, \\ 84 \quad (1.20) \quad &(a_1b_4 + a_2b_4, a_3b_4 + a_4b_1 + a_4b_2 + a_4b_3 + a_4b_4)F). \end{aligned}$$

85 **Theorem 1.7** ([1]). $(NQ, +)$ is an abelian group.

86 **Theorem 1.8** ([1]). (NQ, \cdot) is a commutative monoid.

87 **Theorem 1.9** ([1]). (NQ, \cdot) is not a group.

88 **Theorem 1.10** ([1]). $(NQ, +, \cdot)$ is a commutative ring.

89 **Definition 1.11.** Let NQR be a neutrosophic quadruple ring and let NQS be a
90 nonempty subset of NQR . Then NQS is called a neutrosophic quadruple subring of
91 NQR , if $(NQS, +, \cdot)$ is itself a neutrosophic quadruple ring. For example, $NQR(n\mathbb{Z})$
92 is a neutrosophic quadruple subring of $NQR(\mathbb{Z})$ for $n = 1, 2, 3, \dots$.

93 **Definition 1.12.** Let NQJ be a nonempty subset of a neutrosophic quadruple
94 ring NQR . NQJ is called a neutrosophic quadruple ideal of NQR , if for all $x, y \in$
95 $NQJ, r \in NQR$, the following conditions hold:

- 96 (i) $x - y \in NQJ$,
- 97 (ii) $xr \in NQJ$ and $rx \in NQJ$.

98 **Definition 1.13** ([1]). Let NQR and NQS be two neutrosophic quadruple rings
99 and let $\phi : NQR \rightarrow NQS$ be a mapping defined for all $x, y \in NQR$ as follows:

- 100 (i) $\phi(x + y) = \phi(x) + \phi(y)$,
- 101 (ii) $\phi(xy) = \phi(x)\phi(y)$,
- 102 (iii) $\phi(T) = T, \phi(I) = I$ and $\phi(F) = F$,
- 103 (iv) $\phi(1, 0, 0, 0) = (1, 0, 0, 0)$.

104 Then ϕ is called a neutrosophic quadruple homomorphism. Neutrosophic quadruple
105 monomorphism, endomorphism, isomorphism, and other morphisms can be defined
106 in the usual way.

107 **Definition 1.14.** Let $\phi : NQR \rightarrow NQS$ be a neutrosophic quadruple ring homo-
108 morphism.

(i) The image of ϕ denoted by $Im\phi$ is defined by the set

$$Im\phi = \{y \in NQS : y = \phi(x), \text{ for some } x \in NQR\}.$$

(ii) The kernel of ϕ denoted by $Ker\phi$ is defined by the set

$$Ker\phi = \{x \in NQR : \phi(x) = (0, 0, 0, 0)\}.$$

Theorem 1.15 ([1]). *Let $\phi : NQR \rightarrow NQS$ be a neutrosophic quadruple ring homomorphism. Then:*

- (1) $Im\phi$ is a neutrosophic quadruple subring of NQS ,
- (2) $Ker\phi$ is not a neutrosophic quadruple ideal of NQR .

Theorem 1.16 ([1]). *Let $\phi : NQR(\mathbb{Z}) \rightarrow NQR(\mathbb{Z})/NQR(n\mathbb{Z})$ be a mapping defined by $\phi(x) = x + NQR(n\mathbb{Z})$ for all $x \in NQR(\mathbb{Z})$ and $n = 1, 2, 3, \dots$. Then ϕ is not a neutrosophic quadruple ring homomorphism.*

Definition 1.17. Let H be a non-empty set and let $+$ be a hyperoperation on H . The couple $(H, +)$ is called a canonical hypergroup if the following conditions hold:

- (i) $x + y = y + x$, for all $x, y \in H$,
- (ii) $x + (y + z) = (x + y) + z$, for all $x, y, z \in H$,
- (iii) there exists a neutral element $0 \in H$ such that $x + 0 = \{x\} = 0 + x$, for all $x \in H$,
- (iv) for every $x \in H$, there exists a unique element $-x \in H$ such that $0 \in x + (-x) \cap (-x) + x$,
- (v) $z \in x + y$ implies $y \in -x + z$ and $x \in z - y$, for all $x, y, z \in H$.

A nonempty subset A of H is called a subcanonical hypergroup, if A is a canonical hypergroup under the same hyperaddition as that of H that is, for every $a, b \in A$, $a - b \in A$. If in addition $a + A - a \subseteq A$ for all $a \in H$, A is said to be normal.

Definition 1.18. A hyperring is a tripple $(R, +, \cdot)$ satisfying the following axioms:

- (i) $(R, +)$ is a canonical hypergroup,
- (ii) (R, \cdot) is a semihypergroup such that $x \cdot 0 = 0 \cdot x = 0$ for all $x \in R$, that is, 0 is a bilaterally absorbing element,
- (iii) for all $x, y, z \in R$,

$$x \cdot (y + z) = x \cdot y + x \cdot z \text{ and } (x + y) \cdot z = x \cdot z + y \cdot z.$$

That is, the hyperoperation \cdot is distributive over the hyperoperation $+$.

Definition 1.19. Let $(R, +, \cdot)$ be a hyperring and let A be a nonempty subset of R . A is said to be a subhyperring of R if $(A, +, \cdot)$ is itself a hyperring.

Definition 1.20. Let A be a subhyperring of a hyperring R . Then

- (i) A is called a left hyperideal of R if $r \cdot a \subseteq A$ for all $r \in R, a \in A$,
- (ii) A is called a right hyperideal of R if $a \cdot r \subseteq A$ for all $r \in R, a \in A$,
- (iii) A is called a hyperideal of R if A is both left and right hyperideal of R .

Definition 1.21. Let A be a hyperideal of a hyperring R . A is said to be normal in R , if $r + A - r \subseteq A$, for all $r \in R$.

For full details about hypergroups, canonical hypergroups, hyperrings, neutrosophic canonical hypergroups and neutrosophic hyperrings, the reader should see [3, 14]

2. DEVELOPMENT OF NEUTROSOPHIC QUADRUPLE CANONICAL HYPERGROUPS AND NEUTROSOPHIC QUADRUPLE HYPERRINGS

In this section, we develop two neutrosophic hyperquadruple algebraic hyperstructures namely neutrosophic quadruple canonical hypergroup and neutrosophic quadruple hyperring. In what follows, all neutrosophic quadruple numbers will be real neutrosophic quadruple numbers i.e $a, b, c, d \in \mathbb{R}$ for any neutrosophic quadruple number $(a, bT, cI, dF) \in NQ$.

Definition 2.1. Let $+$ and \cdot be hyperoperations on \mathbb{R} that is $x + y \subseteq \mathbb{R}, x \cdot y \subseteq \mathbb{R}$ for all $x, y \in \mathbb{R}$. Let $\hat{+}$ and $\hat{\cdot}$ be hyperoperations on NQ . For $x = (x_1, x_2T, x_3I, x_4F), y = (y_1, y_2T, y_3I, y_4F) \in NQ$ with $x_i, y_i \in \mathbb{R}, i = 1, 2, 3, 4$, define:

$$(2.1) \quad \begin{aligned} x \hat{+} y &= \{(a, bT, cI, dF) : a \in x_1 + y_1, b \in x_2 + y_2, \\ &c \in x_3 + y_3, d \in x_4 + y_4\}, \end{aligned}$$

$$(2.2) \quad \begin{aligned} x \hat{\cdot} y &= \{(a, bT, cI, dF) : a \in x_1 \cdot y_1, b \in (x_1 \cdot y_2) \cup (x_2 \cdot y_1) \cup (x_2 \cdot y_2), c \in (x_1 \cdot y_3) \\ &\cup (x_2 \cdot y_3) \cup (x_3 \cdot y_1) \cup (x_3 \cdot y_2) \cup (x_3 \cdot y_3), d \in (x_1 \cdot y_4) \cup (x_2 \cdot y_4) \\ &\cup (x_3 \cdot y_4) \cup (x_4 \cdot y_1) \cup (x_4 \cdot y_2) \cup (x_4 \cdot y_3) \cup (x_4 \cdot y_4)\}. \end{aligned}$$

Theorem 2.2. $(NQ, \hat{+})$ is a canonical hypergroup.

Proof. Let $x = (x_1, x_2T, x_3I, x_4F), y = (y_1, y_2T, y_3I, y_4F), z = (z_1, z_2T, z_3I, z_4F) \in NQ$ be arbitrary with $x_i, y_i, z_i \in \mathbb{R}, i = 1, 2, 3, 4$.

(i) To show that $x \hat{+} y = y \hat{+} x$, let

$$\begin{aligned} x \hat{+} y &= \{a = (a_1, a_2T, a_3I, a_4F) : a_1 \in x_1 + y_1, a_2 \in x_2 + y_2, a_3 \in x_3 + y_3, \\ &a_4 \in x_4 + y_4\}, \\ y \hat{+} x &= \{b = (b_1, b_2T, b_3I, b_4F) : b_1 \in y_1 + x_1, b_2 \in y_2 + x_2, b_3 \in y_3 + x_3, \\ &b_4 \in y_4 + x_4\}. \end{aligned}$$

Since $a_i, b_i \in \mathbb{R}, i = 1, 2, 3, 4$, it follows that $x \hat{+} y = y \hat{+} x$.

(ii) To show that $x \hat{+} (y \hat{+} z) = (x \hat{+} y) \hat{+} z$, let

$$\begin{aligned} y \hat{+} z &= \{w = (w_1, w_2T, w_3I, w_4F) : w_1 \in y_1 + z_1, w_2 \in y_2 + z_2, \\ &w_3 \in y_3 + z_3, w_4 \in y_4 + z_4\}. \end{aligned}$$

$$\begin{aligned} x \hat{+} (y \hat{+} z) &= x \hat{+} w \\ &= \{p = (p_1, p_2T, p_3I, p_4F) : p_1 \in x_1 + w_1, p_2 \in x_2 + w_2, p_3 \in x_3 + w_3, \\ &p_4 \in x_4 + w_4\} \\ &= \{p = (p_1, p_2T, p_3I, p_4F) : p_1 \in x_1 + (y_1 + z_1), p_2 \in x_2 + (y_2 + z_2), \\ &p_3 \in x_3 + (y_3 + z_3), p_4 \in x_4 + (y_4 + z_4)\}. \end{aligned}$$

Also, let $x \hat{+} y = \{u = (u_1, u_2T, u_3I, u_4F) : u_1 \in x_1 + y_1, u_2 \in x_2 + y_2, u_3 \in x_3 + y_3, u_4 \in x_4 + y_4\}$ so that

$$\begin{aligned} (x \hat{+} y) \hat{+} z &= u \hat{+} z \\ &= \{q = (q_1, q_2T, q_3I, q_4F) : q_1 \in u_1 + z_1, q_2 \in u_2 + z_2, q_3 \in u_3 + z_3, \\ &q_4 \in u_4 + z_4\} \\ &= \{q = (q_1, q_2T, q_3I, q_4F) : q_1 \in (x_1 + y_1) + z_1, q_2 \in (x_2 + y_2) + z_2, \\ &q_3 \in (x_3 + y_3) + z_3, q_4 \in (x_4 + y_4) + z_4\}. \end{aligned}$$

Since $u_i, p_i, q_i, w_i, x_i, y_i, z_i \in \mathbb{R}, i = 1, 2, 3, 4$, it follows that $x \hat{+} (y \hat{+} z) = (x \hat{+} y) \hat{+} z$.

(iii) To show that $0 = (0, 0, 0, 0) \in NQ$ is a neutral element, consider

$$\begin{aligned} x \hat{+} (0, 0, 0, 0) &= \{a = (a_1, a_2T, a_3I, a_4F) : a_1 \in x_1 + 0, a_2 \in x_2 + 0, a_3 \in x_3 + 0, \\ &\quad a_4 \in x_4 + 0\} \\ &= \{a = (a_1, a_2T, a_3I, a_4F) : a_1 \in \{x_1\}, a_2 \in \{x_2\}, a_3 \in \{x_3\}, \\ &\quad a_4 \in \{x_4\}\} \\ &= \{x\}. \end{aligned}$$

Similarly, it can be shown that $(0, 0, 0, 0) \hat{+} x = \{x\}$. Hence $0 = (0, 0, 0, 0) \in NQ$ is a neutral element.

(iv) To show that for every $x \in NQ$, there exists a unique element $\hat{-}x \in NQ$ such that $0 \in x \hat{+} (\hat{-}x) \cap (\hat{-}x) \hat{+} x$, consider

$$\begin{aligned} x \hat{+} (\hat{-}x) \cap (\hat{-}x) \hat{+} x &= \{a = (a_1, a_2T, a_3I, a_4F) : a_1 \in x_1 - x_1, a_2 \in x_2 - x_2, \\ &\quad a_3 \in x_3 - x_3, a_4 \in x_4 - x_4\} \cap \{b = (b_1, b_2T, b_3I, b_4F) : \\ &\quad b_1 \in -x_1 + x_1, b_2 \in -x_2 + x_2, b_3 \in -x_3 + x_3, b_4 \in -x_4 + x_4\} \\ &= \{(0, 0, 0, 0)\}. \end{aligned}$$

This shows that for every $x \in NQ$, there exists a unique element $\hat{-}x \in NQ$ such that $0 \in x \hat{+} (\hat{-}x) \cap (\hat{-}x) \hat{+} x$.

(v) Since for all $x, y, z \in NQ$ with $x_i, y_i, z_i \in \mathbb{R}, i = 1, 2, 3, 4$, it follows that $z \in x \hat{+} y$ implies $y \in \hat{-}x \hat{+} z$ and $x \in z \hat{+} (\hat{-}y)$. Hence, $(NQ, \hat{+})$ is a canonical hypergroup. \square

Lemma 2.3. Let $(NQ, \hat{+})$ be a neutrosophic quadruple canonical hypergroup. Then

- (1) $\hat{-}(\hat{-}x) = x$ for all $x \in NQ$,
- (2) $0 = (0, 0, 0, 0)$ is the unique element such that for every $x \in NQ$, there is an element $\hat{-}x \in NQ$ such that $0 \in x \hat{+} (\hat{-}x)$,
- (3) $\hat{-}0 = 0$,
- (4) $\hat{-}(x \hat{+} y) = \hat{-}x \hat{-} y$ for all $x, y \in NQ$.

Example 2.4. Let $NQ = \{0, x, y\}$ be a neutrosophic quadruple set and let $\hat{+}$ be a hyperoperation on NQ defined in the table below.

$\hat{+}$	0	x	y
0	0	x	y
x	x	$\{0, x, y\}$	y
y	y	y	$\{0, y\}$

Then $(NQ, \hat{+})$ is a neutrosophic quadruple canonical hypergroup.

Theorem 2.5. $(NQ, \hat{\times})$ is a semihypergroup.

Proof. Let $x = (x_1, x_2T, x_3I, x_4F), y = (y_1, y_2T, y_3I, y_4F), z = (z_1, z_2T, z_3I, z_4F) \in NQ$ be arbitrary with $x_i, y_i, z_i \in \mathbb{R}, i = 1, 2, 3, 4$.

(i)

$$\begin{aligned} x \hat{\times} y &= \{a = (a_1, a_2T, a_3I, a_4F) : a_1 \in x_1y_1, a_2 \in x_1y_2 \cup x_2y_1 \cup x_2y_2, a_3 \in x_1y_3 \\ &\cup x_2y_3 \cup x_3y_1 \cup x_3y_2 \cup x_3y_3, a_4 \in x_1y_4 \cup x_2y_4 \\ &\cup x_3y_4 \cup x_4y_1 \cup x_4y_2 \cup x_4y_3 \cup x_4y_4\} \\ &\subseteq NQ. \end{aligned}$$

(ii) To show that $x \hat{\times} (y \hat{\times} z) = (x \hat{\times} y) \hat{\times} z$, let

$$\begin{aligned} y \hat{\times} z &= \{w = (w_1, w_2T, w_3I, w_4F) : w_1 \in y_1z_1, w_2 \in y_1z_2 \cup y_2z_1 \cup y_2z_2, \\ &w_3 \in y_1z_3 \cup y_2z_3 \cup y_3z_1 \cup y_3z_2 \cup y_3z_3, w_4 \in y_1z_4 \cup y_2z_4 \\ &\cup y_3z_4 \cup y_4z_1 \cup y_4z_2 \cup y_4z_3 \cup y_4z_4\} \end{aligned}$$

so that

$$\begin{aligned} x \hat{\times} (y \hat{\times} z) &= x \hat{\times} w \\ &= \{p = (p_1, p_2T, p_3I, p_4F) : p_1 \in x_1w_1, p_2 \in x_1w_2 \cup x_2w_1 \cup x_2w_2, \\ &p_3 \in x_1w_3 \cup x_2w_3 \cup x_3w_1 \cup x_3w_2 \cup x_3w_3, p_4 \in x_1w_4 \cup x_2w_4 \\ &\cup x_3w_4 \cup x_4w_1 \cup x_4w_2 \cup x_4w_3 \cup x_4w_4\}. \end{aligned}$$

Also, let

$$\begin{aligned} x \hat{\times} y &= \{u = (u_1, u_2T, u_3I, u_4F) : u_1 \in x_1y_1, u_2 \in x_1y_2 \cup x_2y_1 \cup x_2y_2, u_3 \in x_1y_3 \\ &\cup x_2y_3 \cup x_3y_1 \cup x_3y_2 \cup x_3y_3, u_4 \in x_1y_4 \cup x_2y_4 \\ &\cup x_3y_4 \cup x_4y_1 \cup x_4y_2 \cup x_4y_3 \cup x_4y_4\} \end{aligned}$$

so that

$$\begin{aligned} (x \hat{\times} y) \hat{\times} z &= u \hat{\times} z \\ &= \{q = (q_1, q_2T, q_3I, q_4F) : q_1 \in u_1z_1, q_2 \in u_1z_2 \cup u_2z_1 \cup u_2z_2, \\ &q_3 \in u_1z_3 \cup u_2z_3 \cup u_3z_1 \cup u_3z_2 \cup u_3z_3, q_4 \in u_1z_4 \cup u_2z_4 \\ &\cup u_3z_4 \cup u_4z_1 \cup u_4z_2 \cup u_4z_3 \cup u_4z_4\}. \end{aligned}$$

Substituting w_i of (2.3) in (2.4) and also substituting u_i of (2.5) in (2.6), where $i = 1, 2, 3, 4$ and since $p_i, q_i, u_i, w_i, x_i, z_i \in \mathbb{R}$, it follows that $x \hat{\times} (y \hat{\times} z) = (x \hat{\times} y) \hat{\times} z$. Consequently, $(NQ, \hat{\times})$ is a semihypergroup which we call neutrosophic quadruple semihypergroup. \square

Remark 2.6. $(NQ, \hat{\times})$ is not a hypergroup.

Definition 2.7. Let $(NQ, \hat{+})$ be a neutrosophic quadruple canonical hypergroup. For any subset NH of NQ , we define

$$\hat{-}NH = \{\hat{-}x : x \in NH\}.$$

A nonempty subset NH of NQ is called a neutrosophic quadruple subcanonical hypergroup, if the following conditions hold:

- (i) $0 = (0, 0, 0, 0) \in NH$,
- (ii) $x \hat{-}y \subseteq NH$ for all $x, y \in NH$.

A neutrosophic quadruple subcanonical hypergroup NH of a neutrosophic quadruple canonical hypergroup NQ is said to be normal, if $x \hat{+}NH \hat{-}x \subseteq NH$ for all $x \in NQ$.

Definition 2.8. Let $(NQ, \hat{+})$ be a neutrosophic quadruple canonical hypergroup. For $x_i \in NQ$ with $i = 1, 2, 3, \dots, n \in \mathbb{N}$, the heart of NQ denoted by NQ_ω is defined by

$$NQ_\omega = \bigcup_{i=1}^n (x_i \hat{-} x_i).$$

In Example 2.4, $NQ_\omega = NQ$.

Definition 2.9. Let $(NQ_1, \hat{+})$ and $(NQ_2, \hat{+}')$ be two neutrosophic quadruple canonical hypergroups. A mapping $\phi : NQ_1 \rightarrow NQ_2$ is called a neutrosophic quadruple strong homomorphism, if the following conditions hold:

- (i) $\phi(x \hat{+} y) = \phi(x) \hat{+}' \phi(y)$ for all $x, y \in NQ_1$,
- (ii) $\phi(T) = T$,
- (iii) $\phi(I) = I$,
- (iv) $\phi(F) = F$,
- (v) $\phi(0) = 0$.

If in addition ϕ is a bijection, then ϕ is called a neutrosophic quadruple strong isomorphism and we write $NQ_1 \cong NQ_2$.

Definition 2.10. Let $\phi : NQ_1 \rightarrow NQ_2$ be a neutrosophic quadruple strong homomorphism of neutrosophic quadruple canonical hypergroups. Then the set $\{x \in NQ_1 : \phi(x) = 0\}$ is called the kernel of ϕ and it is denoted by $\text{Ker}\phi$. Also, the set $\{\phi(x) : x \in NQ_1\}$ is called the image of ϕ and it is denoted by $\text{Im}\phi$.

Theorem 2.11. $(NQ, \hat{+}, \hat{\times})$ is a hyperring.

Proof. That $(NQ, \hat{+})$ is a canonical hypergroup follows from Theorem 2.2. Also, that $(NQ, \hat{\times})$ is a semihypergroup follows from Theorem 2.4.

Next, let $x = (x_1, x_2T, x_3I, x_4F) \in NQ$ be arbitrary with $x_i, y_i, z_i \in \mathbb{R}, i = 1, 2, 3, 4$. Then

$$\begin{aligned} x \hat{\times} 0 &= \{u = (u_1, u_2T, u_3I, u_4F) : u_1 \in x_1.0, u_2 \in x_1.0 \cup x_2.0 \cup x_2.0, u_3 \in x_1.0 \\ &\quad \cup x_2.0 \cup x_3.0 \cup x_3.0 \cup x_3.0, u_4 \in x_1.0 \cup x_2.0 \cup x_3.0 \cup x_4.0 \cup x_4.0 \\ &\quad \cup x_4.0 \cup x_4.0\} \\ &= \{u = (u_1, u_2T, u_3I, u_4F) : u_1 \in \{0\}, u_2 \in \{0\}, u_3 \in \{0\}, u_4 \in \{0\}\} \\ &= \{0\}. \end{aligned}$$

Similarly, it can be shown that $0 \hat{\times} x = \{0\}$. Since x is arbitrary, it follows that $x \hat{\times} 0 = 0 \hat{\times} x = \{0\}$, for all $x \in NQ$. Hence, $0 = (0, 0, 0, 0)$ is a bilaterally absorbing element.

To complete the proof, we have to show that $x \hat{\times} (y \hat{+} z) = (x \hat{\times} y) \hat{+} (x \hat{\times} z)$, for all $x, y, z \in NQ$. To this end, let $x = (x_1, x_2T, x_3I, x_4F), y = (y_1, y_2T, y_3I, y_4F), z = (z_1, z_2T, z_3I, z_4F) \in NQ$ be arbitrary with $x_i, y_i, z_i \in \mathbb{R}, i = 1, 2, 3, 4$. Let

$$(2.7) \quad \begin{aligned} y \hat{+} z &= \{w = (w_1, w_2T, w_3I, w_4F) : w_1 \in y_1 + z_1, w_2 \in y_2 + z_2, w_3 \in y_3 + z_3, \\ &\quad w_4 \in y_4 + z_4\} \end{aligned}$$

so that

$$\begin{aligned} x \hat{\times} (y \hat{+} z) &= x \hat{\times} w \\ &= \{p = (p_1, p_2 T, p_3 I, p_4 F) : p_1 \in x_1 w_1, p_2 \in x_1 w_2 \cup x_2 w_1 \cup x_2 w_2, \\ &\quad p_3 \in x_1 w_3 \cup x_2 w_3 \cup x_3 w_1 \cup x_3 w_2 \cup x_3 y_3, p_4 \in x_1 w_4 \cup x_2 w_4 \\ &\quad \cup x_3 w_4 \cup x_4 w_1 \cup x_4 w_2 \cup x_4 w_3 \cup x_4 w_4\}. \end{aligned} \quad (2.8)$$

Substituting $w_i, i = 1, 2, 3, 4$ of (2.7) in (2.8), we obtain the following:

$$(2.9) \quad p_1 \in x_1(y_1 + z_1),$$

$$(2.10) \quad p_2 \in x_1(y_2 + z_2) \cup x_2(y_1 + z_1) \cup x_2(y_2 + z_2),$$

$$(2.11) \quad p_3 \in x_1(y_3 + z_3) \cup x_2(y_3 + z_3) \cup x_3(y_1 + z_1) \cup x_3(y_2 + z_2) \cup x_3(y_3 + z_3),$$

$$(2.12) \quad p_4 \in x_1(y_4 + z_4) \cup x_2(y_4 + z_4) \cup x_3(y_4 + z_4) \cup x_4(y_1 + z_1) \cup x_4(y_2 + z_2),$$

$$(2.12) \quad \cup x_4(y_3 + z_3) \cup x_4(y_4 + z_4).$$

Also, let

$$\begin{aligned} x \hat{\times} y &= \{u = (u_1, u_2 T, u_3 I, u_4 F) : u_1 \in x_1 y_1, u_2 \in x_1 y_2 \cup x_2 y_1 \cup x_2 y_2, \\ &\quad u_3 \in x_1 y_3 \cup x_2 y_3 \cup x_3 y_1 \cup x_3 y_2 \cup x_3 y_3, u_4 \in x_1 y_4 \cup x_2 y_4 \\ &\quad \cup x_3 y_4 \cup x_4 y_1 \cup x_4 y_2 \cup x_4 y_3 \cup x_4 y_4\} \end{aligned} \quad (2.13)$$

$$\begin{aligned} x \hat{\times} z &= \{v = (v_1, v_2 T, v_3 I, v_4 F) : v_1 \in x_1 z_1, v_2 \in x_1 z_2 \cup x_2 z_1 \cup x_2 z_2, \\ &\quad v_3 \in x_1 z_3 \cup x_2 z_3 \cup x_3 z_1 \cup x_3 z_2 \cup x_3 z_3, v_4 \in x_1 z_4 \cup x_2 z_4 \\ &\quad \cup x_3 z_4 \cup x_4 z_1 \cup x_4 z_2 \cup x_4 z_3 \cup x_4 z_4\} \end{aligned} \quad (2.14)$$

so that

$$\begin{aligned} (x \hat{\times} y) \hat{+} (x \hat{\times} z) &= u \hat{+} v \\ &= \{q = (q_1, q_2 T, q_3 I, q_4 F) : q_1 \in u_1 + v_1, q_2 \in u_2 + v_2, \\ &\quad q_3 \in u_3 + v_3, q_4 \in u_4 + v_4\}. \end{aligned} \quad (2.15)$$

Substituting u_i of (2.13) and v_i of (2.14) in (2.15), we obtain the following:

$$(2.16) \quad q_1 \in u_1 + v_1 \subseteq x_1 y_1 + x_1 z_1 \subseteq x_1(y_1 + z_1),$$

$$(2.17) \quad q_2 \in u_2 + v_2 \subseteq (x_1 y_2 \cup x_2 y_1 \cup x_2 y_2) \\ + (x_1 z_2 \cup x_2 z_1 \cup x_2 z_2)$$

$$(2.17) \quad \subseteq x_1(y_2 + z_2) \cup x_2(y_1 + z_1) \cup x_2(y_2 + z_2),$$

$$(2.18) \quad q_3 \in u_3 + v_3 \subseteq (x_1 y_3 \cup x_2 y_3 \cup x_3 y_1) \cup x_3 y_2 \cup x_3 y_3 \\ + (x_1 z_3 \cup x_2 z_3 \cup x_3 z_1) \cup x_3 z_2 \cup x_3 z_3$$

$$(2.18) \quad \subseteq x_1(y_3 + z_3) \cup x_2(y_3 + z_3) \cup x_3(y_1 + z_1) \cup x_3(y_2 + z_2) \cup x_3(y_3 + z_3).$$

$$(2.18) \quad q_4 \in u_4 + v_4 \subseteq (x_1 y_4 \cup x_2 y_4 \cup x_3 y_4) \cup x_4 y_1 \cup x_4 y_2 \cup x_4 y_3 \cup x_4 y_4 \\ + (x_1 z_4 \cup x_2 z_4 \cup x_3 z_4) \cup x_4 z_1 \cup x_4 z_2 \cup x_4 z_3 \cup x_4 z_4$$

$$(2.18) \quad \subseteq x_1(y_4 + z_4) \cup x_2(y_4 + z_4) \cup x_3(y_4 + z_4) \cup x_4(y_1 + z_1) \cup x_4(y_2 + z_2) \\ \cup x_4(y_3 + z_3) \cup x_4(y_4 + z_4).$$

$$(2.19) \quad \cup x_4(y_3 + z_3) \cup x_4(y_4 + z_4).$$

$$(2.19) \quad \cup x_4(y_3 + z_3) \cup x_4(y_4 + z_4).$$

Comparing (2.9), (2.10), (2.11) and (2.12) respectively with (2.16), (2.17), (2.18) and (2.19), we obtain $p_i = q_i, i = 1, 2, 3, 4$. Hence, $x \hat{\times} (y \hat{+} z) = (x \hat{\times} y) \hat{+} (x \hat{\times} z)$, for all

333 $x, y, z \in NQ$. Thus, $(NQ, \hat{+}, \hat{\times})$ is a hyperring which we call neutrosophic quadruple
334 hyperring. \square

335 **Theorem 2.12.** $(NQ, \hat{+}, \circ)$ is a Krasner hyperring where \circ is an ordinary multi-
336 plicative binary operation on NQ .

337 **Definition 2.13.** Let $(NQ, \hat{+}, \hat{\times})$ be a neutrosophic quadruple hyperring. A nonempty
338 subset NJ of NQ is called a neutrosophic quadruple subhyperring of NQ , if $(NJ, \hat{+}, \hat{\times})$
339 is itself a neutrosophic quadruple hyperring.

340 NJ is called a neutrosophic quadruple hyperideal if the following conditions hold:

- 341 (i) $(NJ, \hat{+})$ is a neutrosophic quadruple subcanonical hypergroup.
- 342 (ii) For all $x \in NJ$ and $r \in NQ$, $x \hat{\times} r, r \hat{\times} x \subseteq NJ$.

343 A neutrosophic quadruple hyperideal NJ of NQ is said to be normal in NQ , if
344 $x \hat{+} NJ \hat{-} x \subseteq NJ$, for all $x \in NQ$.

345 **Definition 2.14.** Let $(NQ_1, \hat{+}, \hat{\times})$ and $(NQ_2, \hat{+}', \hat{\times}')$ be two neutrosophic quadruple
346 hyperrings. A mapping $\phi : NQ_1 \rightarrow NQ_2$ is called a neutrosophic quadruple
347 strong homomorphism, if the following conditions hold:

- 348 (i) $\phi(x \hat{+} y) = \phi(x) \hat{+}' \phi(y)$, for all $x, y \in NQ_1$,
- 349 (ii) $\phi(x \hat{\times} y) = \phi(x) \hat{\times}' \phi(y)$, for all $x, y \in NQ_1$,
- 350 (iii) $\phi(T) = T$,
- 351 (iv) $\phi(I) = I$,
- 352 (v) $\phi(F) = F$,
- 353 (vi) $\phi(0) = 0$.

354 If in addition ϕ is a bijection, then ϕ is called a neutrosophic quadruple strong
355 isomorphism and we write $NQ_1 \cong NQ_2$.

356 **Definition 2.15.** Let $\phi : NQ_1 \rightarrow NQ_2$ be a neutrosophic quadruple strong homo-
357 morphism of neutrosophic quadruple hyperrings. Then the set $\{x \in NQ_1 : \phi(x) = 0\}$
358 is called the kernel of ϕ and it is denoted by $Ker\phi$. Also, the set $\{\phi(x) : x \in NQ_1\}$
359 is called the image of ϕ and it is denoted by $Im\phi$.

360 **Example 2.16.** Let $(NQ, \hat{+}, \hat{\times})$ be a neutrosophic quadruple hyperring and let
361 NX be the set of all strong endomorphisms of NQ . If \oplus and \odot are hyperoperations
362 defined for all $\phi, \psi \in NX$ and for all $x \in NQ$ as

$$\begin{aligned} 363 \quad \phi \oplus \psi &= \{\nu(x) : \nu(x) \in \phi(x) \hat{+} \psi(x)\}, \\ 364 \quad \phi \odot \psi &= \{\nu(x) : \nu(x) \in \phi(x) \hat{\times} \psi(x)\}, \end{aligned}$$

365 then (NX, \oplus, \odot) is a neutrosophic quadruple hyperring.

366 3. CHARACTERIZATION OF NEUTROSOPHIC QUADRUPLE CANONICAL 367 HYPERGROUPS AND NEUTROSOPHIC HYPERRINGS

368 In this section, we present elementary properties which characterize neutrosophic
369 quadruple canonical hypergroups and neutrosophic quadruple hyperrings.

370 **Theorem 3.1.** Let NG and NH be neutrosophic quadruple subcanonical hyper-
371 groups of a neutrosophic quadruple canonical hypergroup $(NQ, \hat{+})$. Then

- 372 (1) $NG \cap NH$ is a neutrosophic quadruple subcanonical hypergroup of NQ ,

(2) $NG \times NH$ is a neutrosophic quadruple subcanonical hypergroup of NQ .

Theorem 3.2. Let NH be a neutrosophic quadruple subcanonical hypergroup of a neutrosophic quadruple canonical hypergroup $(NQ, \hat{+})$. Then

- (1) $NH \hat{+} NH = NH$,
- (2) $x \hat{+} NH = NH$, for all $x \in NH$.

Theorem 3.3. Let $(NQ, \hat{+})$ be a neutrosophic quadruple canonical hypergroup. NQ_ω , the heart of NQ is a normal neutrosophic quadruple subcanonical hypergroup of NQ .

Theorem 3.4. Let NG and NH be neutrosophic quadruple subcanonical hypergroups of a neutrosophic quadruple canonical hypergroup $(NQ, \hat{+})$.

- (1) If $NG \subseteq NH$ and NG is normal, then NG is normal.
- (2) If NG is normal, then $NG \hat{+} NH$ is normal.

Definition 3.5. Let NG and NH be neutrosophic quadruple subcanonical hypergroups of a neutrosophic quadruple canonical hypergroup $(NQ, \hat{+})$. The set $NG \hat{+} NH$ is defined by

$$(3.1) \quad NG \hat{+} NH = \{x \hat{+} y : x \in NG, y \in NH\}.$$

It is obvious that $NG \hat{+} NH$ is a neutrosophic quadruple subcanonical hypergroup of $(NQ, \hat{+})$.

If $x \in NH$, the set $x \hat{+} NH$ is defined by

$$(3.2) \quad x \hat{+} NH = \{x \hat{+} y : y \in NH\}.$$

If x and y are any two elements of NH and τ is a relation on NH defined by $x \tau y$ if $x \in y \hat{+} NH$, it can be shown that τ is an equivalence relation on NH and the equivalence class of any element $x \in NH$ determined by τ is denoted by $[x]$.

Lemma 3.6. For any $x \in NH$, we have

- (1) $[x] = x \hat{+} NH$,
- (2) $[\hat{-}x] = \hat{-}[x]$.

Proof. (1)

$$\begin{aligned} [x] &= \{y \in NH : x \tau y\} \\ &= \{y \in NH : y \in x \hat{+} NH\} \\ &= x \hat{+} NH. \end{aligned}$$

(2) Obvious. □

Definition 3.7. Let NQ/NH be the collection of all equivalence classes of $x \in NH$ determined by τ . For $[x], [y] \in NQ/NH$, we define the set $[x] \hat{\oplus} [y]$ as

$$(3.3) \quad [x] \hat{\oplus} [y] = \{[z] : z \in x \hat{+} y\}.$$

Theorem 3.8. $(NQ/NH, \hat{\oplus})$ is a neutrosophic quadruple canonical hypergroup.

Proof. Same as the classical case and so omitted. □

Theorem 3.9. Let $(NQ, \hat{+})$ be a neutrosophic quadruple canonical hypergroup and let NH be a normal neutrosophic quadruple subcanonical hypergroup of NQ . Then, for any $x, y \in NH$, the following are equivalent:

- (1) $x \in y \hat{+} NH$,
- (2) $y \hat{-} x \subseteq NH$,
- (3) $(y \hat{-} x) \cap NH \neq \emptyset$

Proof. Same as the classical case and so omitted. \square

Theorem 3.10. Let $\phi : NQ_1 \rightarrow NQ_2$ be a neutrosophic quadruple strong homomorphism of neutrosophic quadruple canonical hypergroups. Then

- (1) $\text{Ker}\phi$ is not a neutrosophic quadruple subcanonical hypergroup of NQ_1 ,
- (2) $\text{Im}\phi$ is a neutrosophic quadruple subcanonical hypergroup of NQ_2 .

Proof. (1) Since it is not possible to have $\phi((0, T, 0, 0)) = \phi((0, 0, 0, 0))$, $\phi((0, 0, I, 0)) = \phi((0, 0, 0, 0))$ and $\phi((0, 0, 0, F)) = \phi((0, 0, 0, 0))$, it follows that $(0, T, 0, 0)$, $(0, 0, I, 0)$ and $(0, 0, 0, F)$ cannot be in the kernel of ϕ . Consequently, $\text{Ker}\phi$ cannot be a neutrosophic quadruple subcanonical hypergroup of NQ_1 .

(2) Obvious. \square

Remark 3.11. If $\phi : NQ_1 \rightarrow NQ_2$ is a neutrosophic quadruple strong homomorphism of neutrosophic quadruple canonical hypergroups, then $\text{Ker}\phi$ is a subcanonical hypergroup of NQ_1 .

Theorem 3.12. Let $\phi : NQ_1 \rightarrow NQ_2$ be a neutrosophic quadruple strong homomorphism of neutrosophic quadruple canonical hypergroups. Then

- (1) $NQ_1/\text{Ker}\phi$ is not a neutrosophic quadruple canonical hypergroup,
- (2) $NQ_1/\text{Ker}\phi$ is a canonical hypergroup.

Theorem 3.13. Let NH be a neutrosophic quadruple subcanonical hypergroup of the neutrosophic quadruple canonical hypergroup $(NQ, \hat{+})$. Then the mapping $\phi : NQ \rightarrow NQ/NH$ defined by $\phi(x) = x \hat{+} NH$ is not a neutrosophic quadruple strong homomorphism.

Remark 3.14. Isomorphism theorems do not hold in the class of neutrosophic quadruple canonical hypergroups.

Lemma 3.15. Let NJ be a neutrosophic quadruple hyperideal of a neutrosophic quadruple hyperring $(NQ, \hat{+}, \hat{\times})$. Then

- (1) $\hat{-}NJ = NJ$,
- (2) $x \hat{+} NJ = NJ$, for all $x \in NJ$,
- (3) $x \hat{\times} NJ = NJ$, for all $x \in NJ$.

Theorem 3.16. Let NJ and NK be neutrosophic quadruple hyperideals of a neutrosophic quadruple hyperring $(NQ, \hat{+}, \hat{\times})$. Then

- (1) $NJ \cap NK$ is a neutrosophic quadruple hyperideal of NQ ,
- (2) $NJ \times NK$ is a neutrosophic quadruple hyperideal of NQ ,
- (3) $NJ \hat{+} NK$ is a neutrosophic quadruple hyperideal of NQ .

Theorem 3.17. Let NJ be a normal neutrosophic quadruple hyperideal of a neutrosophic quadruple hyperring $(NQ, \hat{+}, \hat{\times})$. Then

- 450 (1) $(x \hat{+} NJ) \hat{+} (y \hat{+} NJ) = (x \hat{+} y) \hat{+} NJ$, for all $x, y \in NJ$,
 451 (2) $(x \hat{+} NJ) \hat{\times} (y \hat{+} NJ) = (x \hat{\times} y) \hat{+} NJ$, for all $x, y \in NJ$,
 452 (3) $x \hat{+} NJ = y \hat{+} NJ$, for all $y \in x \hat{+} NJ$.

453 **Theorem 3.18.** Let NJ and NK be neutrosophic quadruple hyperideals of a neu-
 454 trosophic quadruple hyperring $(NQ, \hat{+}, \hat{\times})$ such that NJ is normal in NQ . Then

- 455 (1) $NJ \cap NK$ is normal in NJ ,
 456 (2) $NJ \hat{+} NK$ is normal in NQ ,
 457 (3) NJ is normal in $NJ \hat{+} NK$.

458 Let NJ be a neutrosophic quadruple hyperideal of a neutrosophic quadruple
 459 hyperring $(NQ, \hat{+}, \hat{\times})$. For all $x \in NQ$, the set NQ/NJ is defined as

$$460 \quad (3.4) \quad NQ/NJ = \{x \hat{+} NJ : x \in NQ\}.$$

461 For $[x], [y] \in NQ/NJ$, we define the hyperoperations $\hat{\oplus}$ and $\hat{\otimes}$ on NQ/NJ as follows:

$$462 \quad (3.5) \quad [x] \hat{\oplus} [y] = \{[z] : z \in x \hat{+} y\},$$

$$464 \quad (3.6) \quad [x] \hat{\otimes} [y] = \{[z] : z \in x \hat{\times} y\}.$$

465 It can easily be shown that $(NQ/NH, \hat{\oplus}, \hat{\otimes})$ is a neutrosophic quadruple hyperring.

466 **Theorem 3.19.** Let $\phi : NQ \rightarrow NR$ be a neutrosophic quadruple strong homomor-
 467 phism of neutrosophic quadruple hyperrings and let NJ be a neutrosophic quadruple
 468 hyperideal of NQ . Then

- 469 (1) $\text{Ker} \phi$ is not a neutrosophic quadruple hyperideal of NQ ,
 470 (2) $\text{Im} \phi$ is a neutrosophic quadruple hyperideal of NR ,
 471 (3) $NQ/\text{Ker} \phi$ is not a neutrosophic quadruple hyperring,
 472 (4) $NQ/\text{Im} \phi$ is a neutrosophic quadruple hyperring,
 473 (5) The mapping $\psi : NQ \rightarrow NQ/NJ$ defined by $\psi(x) = x \hat{+} NJ$, for all $x \in NQ$
 474 is not a neutrosophic quadruple strong homomorphism.

475 **Remark 3.20.** The classical isomorphism theorems of hyperrings do not hold in
 476 neutrosophic quadruple hyperrings.

477 4. CONCLUSION

478 We have developed neutrosophic quadruple algebraic hyperstructures in this pa-
 479 per. In particular, we have developed new neutrosophic algebraic hyperstructures
 480 namely neutrosophic quadruple semihypergroups, neutrosophic quadruple canonical
 481 hypergroups and neutrosophic quadruple hyperrings. We have presented elementary
 482 properties which characterize the new neutrosophic algebraic hyperstructures.

483 **Acknowledgements.** The authors thank all the anonymous reviewers for useful
 484 observations and critical comments which have improved the quality of the paper.

485 REFERENCES

- 486 [1] S. A. Akinleye, F. Smarandache and A. A. A. Agboola, On Neutrosophic Quadruple Algebraic
 487 Structures, Neutrosophic Sets and Systems 12 (2016) 122–126.
 488 [2] S. A. Akinleye, E. O. Adeleke and A. A. A. Agboola, Introduction to Neutrosophic Nerrings,
 489 Ann. Fuzzy Math. Inform. 12 (3) (2016) 7–19.

- 490 [3] A. A. A. Agboola, On Refined Neutrosophic Algebraic Structures I, Neutrosophic Sets and
491 Systems 10 (2015) 99–101.
- 492 [4] A. A. A. Agboola and B. Davvaz, On Neutrosophic Canonical Hypergroups and Neutrosophic
493 Hyperrings, Neutrosophic Sets and Systems 2 (2014) 34–41.
- 494 [5] A. Asokkumar and M. Velrajan, Characterization of regular hyperrings, Italian Journal of
495 Pure and Applied Mathematics 22 (2007) 115–124.
- 496 [6] A. R. Bargi, A class of hyperrings, J. Disc. Math. Sc. and Cryp. (6) (2003) 227–233.
- 497 [7] P. Corsini, Prolegomena of Hypergroup Theory, Second edition, Aviani editore 1993.
- 498 [8] P. Corsini and V. Leoreanu, Applications of Hyperstructure Theory, Advances in Mathemat-
499 ics, Kluwer Academic Publishers, Dordrecht 2003.
- 500 [9] B. Davvaz, Polygroup Theory and Related Systems, World Sci. Publ. 2013.
- 501 [10] B. Davvaz, Isomorphism theorems of hyperrings, Indian J. Pure Appl. Math. 35 (3) (2004)
502 321–333.
- 503 [11] B. Davvaz, Approximations in hyperrings, J. Mult.-Valued Logic Soft Comput. 15 (5-6)
504 (2009) 471–488.
- 505 [12] B. Davvaz and A. Salasi, A realization of hyperrings, Comm. Algebra 34 (2006) 4389–4400.
- 506 [13] B. Davvaz and T. Vougiouklis, Commutative rings obtained from hyperrings (H_v -rings) with
507 α^* -relations, Comm. Algebra 35 (2007) 3307–3320.
- 508 [14] B. Davvaz and V. Leoreanu-Fotea, Hyperring Theory and Applications, International Aca-
509 demic Press, USA 2007.
- 510 [15] B. Davvaz, Isomorphism theorems of hyperrings, Indian Journal of Pure and Applied Math-
511 ematics 23 (3) (2004) 321–331.
- 512 [16] M. De Salvo, Hyperrings and hyperfields, Annales Scientifiques de l’Universite de Clermont-
513 Ferrand II, 22 (1984) 89–107.
- 514 [17] F. Smarandache, Neutrosophy/Neutrosophic Probability, Set, and Logic, American Research
515 Press, Rehoboth, USA 1998.
- 516 <http://fs.gallup.unm.edu/eBook-otherformats.htm>
- 517 [18] F. Smarandache, Neutrosophic Quadruple Numbers, Refined Neutrosophic Quadruple Num-
518 bers, Absorbance Law, and the Multiplication of Neutrosophic Quadruple Numbers, Neutro-
519 sophic Sets and Systems 10 (2015) 96–98.
- 520 [19] F. Smarandache, (t,i,f) - Neutrosophic Structures and I-Neutrosophic Structures, Neutro-
521 sophic Sets and Systems 8 (2015) 3–10.
- 522 [20] F. Smarandache, n-Valued Refined Neutrosophic Logic and Its Applications in Physics,
523 Progress in Physics 4 (2013) 143–146.
- 524 [21] M. Velrajan and A. Asokkumar, Note on Isomorphism Theorems of Hyperrings, Int. J. Math.
525 & Math. Sc. (2010) ID 376985 1–12.

526 A. A. A. AGBOOLA (agboolaaaa@funaab.edu.ng)

527

528 Department of Mathematics, Federal University of Agriculture, Abeokuta, Nige-
529 ria

530 B. DAVVAZ (davvaz@yazd.ac.ir)

531

532 Department of Mathematics, Yazd University, Yazd, Iran

533 F. SMARANDACHE (smarand@unm.edu)

534

535 Department of Mathematics and Science, University of New Mexico, 705 Gurley
536 Ave., Gallup, NM 87301, USA

537