

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/358692583>

The Structure Of Imperfect Triplets In Several Refined Neutrosophic Rings

Article · February 2022

DOI: 10.54216/JNFS.020103

CITATIONS

0

READS

4

4 authors, including:



Mikail Bal

Gaziantep University

21 PUBLICATIONS 86 CITATIONS

[SEE PROFILE](#)



Rozina Ali

Cairo University

33 PUBLICATIONS 10 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Neutrosophic [View project](#)



The Structure Of Imperfect Triplets In Several Refined Neutrosophic Rings

Mikail Bal, Gaziantep University, Turkey

Katy D. Ahmad, Islamic University Of Gaza, Palestine

Arwa A. Hajjari, Cairo University, Egypt

Rozina Ali, Cairo university, Egypt

Corrsepondence: katyon765@gmail.com

rozyyy123n@gmail.com

Abstract: This paper solves the imperfect triplets problem in refined neutrosophic rings, where it presents the necessary and sufficient conditions for a triple (x, y, z) to be an imperfect triplet in any refined neutrosophic ring. Also, this work introduces a full description of the structure of imperfect triplets in numerical refined neutrosophic rings such as refined neutrosophic ring of integers $Z(I_1, I_2)$, refined neutrosophic ring of rationales $Q(I_1, I_2)$, and refined neutrosophic ring of real numbers $R(I_1, I_2)$.

Keywords: Refined Neutrosophic Ring, imperfect Duplet, Imperfect triplet

1. Introduction

Neutrosophy is a generalization of intuitionistic fuzzy logic founded by F.Smarandache to deal with indeterminacy in science and real life problems.

Neutrosophic algebra began with the efforts of Kandasamy and Smarandache where the concept of neutrosophic ring was presented in [7] as a generalization of classical rings. These rings were handled by many authors such as [1,2,9,11-19].

Recently, there is an increasing interest in the generalizations of neutrosophic rings, where refined neutrosophic rings were defined by Agboola et.al [5,6].

If $(R, +, \cdot)$ is a ring, then the corresponding refined neutrosophic ring $R(I_1, I_2)$ is defined as follows:

$R(I_1, I_2) = \{(a, bI_1, cI_2); a, b, c \in R\}$. The operations on $R(I_1, I_2)$ are defined as follows:

Addition: $(a, bI_1, cI_2) + (x, yI_1, zI_2) = (a + x, (b + y)I_1, (c + z)I_2)$.

Multiplication: $(a, bI_1, cI_2) \cdot (x, yI_1, zI_2) = (ax, (ay + bx + by + bz + cy)I_1, (az + cz + cx)I_2)$

The notion of neutrosophic duplets and neutrosophic triplets was defined and handled by Smarandache et.al in [3,8,10], where they opened an interesting research direction about finding these elements in rings.

The structure of imperfect refined neutrosophic duplets was discussed in [38].

Through this paper, we extend the previous efforts to solve the problem of triplets into the case of refined neutrosophic rings, where we present the condition of imperfect triplets in any refined neutrosophic ring even when it is not commutative. In particular, we determine all possible imperfect triplets in the refined neutrosophic ring of integers $Z(I_1, I_2)$, refined neutrosophic ring of rationales $Q(I_1, I_2)$, and refined neutrosophic ring of real numbers $R(I_1, I_2)$.

2. Preliminaries

Definition 2.1: [5]

(a) If X is a set then $X(I_1, I_2) = \{(a, bI_1, cI_2); a, b, c \in X\}$ is called the refined neutrosophic set generated by X, I_1, I_2 .

(b) Let $(R, +, \cdot)$ be a ring, $(R(I_1, I_2), +, \cdot)$ is called the refined neutrosophic ring generated by R, I_1, I_2 .

Definition 2.2: [12]

Let R be any ring, x, y are two arbitrary elements in R . We call them a duplet with y acts as an identity if and only if

$$xy = yx = x.$$

Definition 2.3: [10]

Let R be any ring, x, y, z three arbitrary elements in R . We call them a triplet with y acts as an identity if and only if

$$xy = yx = x, zy = yz = z, xz = zx = y.$$

Theorem 2.4: [38]

Let $x = (x_0, x_1 I_1, x_2 I_2), y = (y_0, y_1 I_1, y_2 I_2)$ be any two elements in $R(I_1, I_2)$, then (x, y) is an imperfect refined neutrosophic duplet with y acts as an identity if and only if

$(x_0, y_0), (x_0 + x_2, y_0 + y_2), (x_0 + x_1 + x_2, y_0 + y_1 + y_2)$ are three imperfect duplets in the classical ring R with $y_0, y_0 + y_2, y_0 + y_1 + y_2$ acts as identities.

For the structure of refined neutrosophic imperfect duplets. See [38].

3. Main discussion :

Theorem 3.1:

Let $Q(I_1, I_2), R(I_1, I_2)$ be the neutrosophic ring of rationales and reals respectively. Consider the following two forms of imperfect duplets

$$1-) \{(0, x_1 I_1, 0), (y_0, y_1 I_1, y_2 I_2); y_0 + y_1 + y_2 = 1, x_1 \neq 0\},$$

$$2-) \{(0, x_1 I_1, -x_1 I_2), (y_0, y_1 I_1, y_2 I_2); y_0 + y_2 = 1 \text{ and } x_1 \neq 0\}.$$

The corresponding imperfect triplets derived from the previous forms are

$$(a-) \{(0, x_1 I_1, 0), (0, I_1, 0), (0, \frac{1}{x_1} I_1, 0); x_1 \neq 0\},$$

$$(b-) \{(0, x_1 I_1, -x_1 I_2), (0, -I_1, I_2), (0, \frac{1}{x_1} I_1, -\frac{1}{x_1} I_2); x_1 \neq 0\}.$$

Proof:

Suppose that (x, y, z) is an imperfect triplet, then $xy = yx = x, xz = zx = y, yz = zy = z$. It is clear that $(x, y), (y, z)$ are imperfect duplets with y acts as an identity.

From the form (1), we have $x = (0, x_1 I_1, 0)$, $y = (y_0, y_1 I_1, y_2 I_2)$, $z = (0, z_1 I_1, 0)$ with $y_0 + y_1 + y_2 = 1$, $x_1 \neq 0$. From the equation $xz = y$, we get that $y_0 = y_2 = 0$, $y_1 = 1$ and $z_1 = \frac{1}{x_1}$, thus the corresponding imperfect triplet is (a).

From the form (2), we have $x = (0, x_1 I_1, -x_1 I_2)$, $y = (y_0, y_1 I_1, y_2 I_2)$, $z = (0, z_1 I_1, -z_1 I_2)$ with $y_0 + y_2 = 1$. From the equation $xz = y$, we get that $y_2 = 1$, $y_1 = -1$, $y_0 = 0$, $z_1 = \frac{1}{x_1}$.

And the corresponding imperfect triplet is (b).

Example 3.2:

We construct an example about an imperfect triplet with form (b).

Put $x_1 = \frac{1}{3}$. The corresponding triplet is $x = (0, \frac{1}{3} I_1, -\frac{1}{3} I_2)$, $y = (0, -I_1, I_2)$, $z = (0, 3 I_1, -3 I_2)$.

Theorem 3.3:

Let $Q(I_1, I_2)$, $R(I_1, I_2)$ be the neutrosophic ring of rationales and reals respectively. Consider the following two forms of imperfect duplets

$$3-) \{(0, x_1 I_1, x_2 I_1), (y_0, 0, y_2 I_2); y_0 + y_2 = 1\},$$

$$4-) \{(x_0, 0, -x_0 I_2), (1, y_1 I_1, y_2 I_2)\},$$

The corresponding imperfect triplets has the forms

$$(c) \{(0, x_1 I_1, x_2 I_2), (0, 0, I_2), (0, \frac{-x_1}{x_2(x_1+x_2)} I_1, \frac{1}{x_2} I_2); x_2, x_1 + x_2 \neq 0\}$$

$$(d) \{(x_0, 0, -x_0 I_2), (1, 0, -I_2), (\frac{1}{x_0}, 0, \frac{-1}{x_0} I_2); x_0 \neq 0\}.$$

Proof:

Suppose that (x, y, z) is an imperfect triplet, then $xy = yx = x$, $xz = zx = y$, $yz = zy = z$. It is clear that (x, y) , (y, z) are imperfect duplets with y acts as an identity.

From the form (3), we have $x = (0, x_1 I_1, x_2 I_1)$, $y = (y_0, 0, y_2 I_2)$, $z = (0, z_1 I_1, z_2 I_1)$ with $y_0 + y_2 = 1$. From the equation $xz = y$, we get:

$y_0 = 0, y_2 = 1, z_2 = \frac{1}{x_2}, z_1 = \frac{-x_1}{x_2(x_1+x_2)}$. Thus the corresponding imperfect triplet is (c) .

From the form (4), we have $x=(x_0, 0, -x_0I_2), y=(1, y_1I_1, y_2I_2), z=(z_0, 0, -z_0I_2)$. From equation $xz = y$, we get $z_0 = \frac{1}{x_0}, y_2 = -1, y_1 = 0$. Thus we get the imperfect triplet (d).

Example 3.4:

We construct an example about an imperfect triplet with form (c).

Put $x_1 = \frac{1}{2}, x_2 = \frac{5}{2}$,

We get the following triplet $x = \left(0, \frac{1}{2}I_1, \frac{5}{2}I_1\right), y = (0, 0, I_2), z = \left(0, -\frac{1}{15}I_1, \frac{2}{5}I_1\right)$

Theorem 3.5:

Let $Q(I_1, I_2), R(I_1, I_2)$ be the neutrosophic ring of rationales and reals respectively. Consider the following two forms of imperfect duplets

5-) $\{(x_0, x_1I_1, -x_0I_2), (1, y_1I_1, -y_1I_2); x_0 \neq 0\}$,

6-) $\{(x_0, x_1I_1, x_2I_2), (1, y_1I_1, 0); x_0 + x_1 + x_2 = 0\}$.

The corresponding imperfect triplets are:

(e) $\{(x_0, x_1I_1, -x_0I_2), (1, I_1, -I_2), (\frac{1}{x_0}, \frac{1}{x_1}I_1, \frac{-1}{x_0}I_2); x_0, x_1 \neq 0\}$,

(f) $\{(x_0, x_1I_1, x_2I_2), (1, -I_1, 0), (\frac{1}{x_0}, z_1I_1, \frac{-x_2}{x_0(x_0+x_2)}I_2); x_0 + x_1 + x_2 = 0, x_0, x_0 + x_2 \neq 0, z_1 \text{ is arbitrary}\}$.

Proof:

The proof holds by a similar discussion.

Example 3.6:

We construct an example about an imperfect triplet with form (e).

Put $x_0 = 1, x_1 = \frac{1}{2}$. Thus we get the following triplet

$$x = \left(1, \frac{1}{2} I_1, -I_2\right), y = (1, I_1, -I_2), z = (1, 2I_1, -I_2).$$

Theorem 3.7:

Let $Q(I_1, I_2)$, $R(I_1, I_2)$ be the neutrosophic ring of rationales and reals respectively. Imperfect triplets has the following 6 forms (a), (b), (c), (d), (e), (f).

The proof holds directly from previous Theorems .

Now, we check the existence of imperfect triplets in the refined neutrosophic ring of integers $Z(I_1, I_2)$.

Remark 3.8:

A triple (x, y, z) is an imperfect triplet in $Z(I_1, I_2)$ if it can be represented by one of the forms (a-f) in Theorem 3.14 as a result of the inclusion between $Z(I_1, I_2)$ and $Q(I_1, I_2)$.

This means that the all imperfect triplets in $Z(I_1, I_2)$ are:

$$(a-) \{(0, x_1 I_1, 0), (0, I_1, 0 I_2), \left(0, \frac{1}{x_1} I_1, 0\right); x_1 \in \{-1, 1\}\},$$

$$(b-) \{(0, x_1 I_1, -x_1 I_1), (0, -I_1, I_2), \left(0, \frac{1}{x_1} I_1, -\frac{1}{x_1} I_1\right); x_1 \in \{-1, 1\}\}.$$

$$(c) \{(0, x_1 I_1, x_2 I_1), (0, 0, I_2), \left(0, \frac{-x_1}{x_2(x_1+x_2)} I_1, \frac{1}{x_2} I_2\right); x_2 \in \{-1, 1\} \text{ and } \frac{-x_1}{x_2(x_1+x_2)} \in Z\}$$

$$(d) \{(x_0, 0, -x_0 I_2), (1, 0, -I_2), \left(\frac{1}{x_0}, 0, \frac{-1}{x_0} I_2\right); x_0 \in \{-1, 1\}\}.$$

$$(e) \{(x_0, x_1 I_1, -x_0 I_2), (1, I_1, -I_2), \left(\frac{1}{x_0}, \frac{1}{x_1} I_1, \frac{-1}{x_0} I_2\right); x_0, x_1 \in \{-1, 1\}\},$$

$$(f) \{(x_0, x_1 I_1, x_2 I_2), (1, -I_1, 0), \left(\frac{1}{x_0}, z_1 I_1, \frac{-x_2}{x_0(x_0+x_2)} I_2\right); x_0 + x_1 + x_2 = 0, x_0 \in \{-1, 1\} \text{ and } z_1, \frac{-x_2}{x_0(x_0+x_2)} \in Z\}.$$

Example 3.9:

The forms (a),(b),(d),(e) are clear. We illustrate an example about an imperfect triplet of form (c).

We put $x_2 = 1$, $x_1 = -2$, so that we get the following triplet

$$x = (0, -2I_1, I_2), y = (0, 0, I_2), z = (0, -2I_1, I_2).$$

Doi : <https://doi.org/10.54216/JNFS.020103>

Received July 25, 2021 Accepted: Jan 05, 2022

Another example about imperfect triplets with form (f).

Put $x_0 = x_1 = 1, x_2 = -2, z_1 = 5$, we get the following triplet

$$x = (1, I_1, -2I_2), y = (1, -I_1, 0), z = (1, 5I_1, -2I_2).$$

Conclusion

In this paper, we have studied the problem of finding imperfect triplets in a refined neutrosophic ring. Where we have determined a general condition for imperfect duplets in such rings. In particular, we have presented the structure of all imperfect triplets in the refined neutrosophic rings of integers, reals, and rationales respectively.

References

- [1] T.Chalapathi and L. Madhavi,. "Neutrosophic Boolean Rings", Neutrosophic Sets and Systems, Vol. 33, pp. 57-66, 2020.
- [2] S. A. Edalatpanah., " Systems of Neutrosophic Linear Equations", Neutrosophic Sets and Systems, Vol. 33, pp. 92-104, 2020.
- [3] Kandasamy, V., Kandasamy, I., and Smarandache, F., " Algebraic Structure of Neutrosophic Duplets In Neutrosophic Rings $\langle Z \cup I \rangle, \langle Q \cup I \rangle$, and $\langle R \cup I \rangle$ ", Neutrosophic Sets and Systems, Vol.23, pp.85-95, 2018.
- [4] Kandasamy, V. W. B., Ilanthenral, K., and Smarandache, F., " Semi-Idempotents in Neutrosophic Rings", Mathematics Journal (MDPI), Vol. 7, 2019.
- [5] Adeleke, E.O., Agboola, A.A.A.,and Smarandache, F., "Refined Neutrosophic Rings I", International Journal of Neutrosophic Science, Vol. 2(2), pp. 77-81, 2020.
- [6] Adeleke, E.O., Agboola, A.A.A., and Smarandache, F., "Refined Neutrosophic Rings II", International Journal of Neutrosophic Science, Vol. 2(2), pp. 89-94, 2020.
- [7] Kandasamy, V.W.B., and Smarandache, F., "Some Neutrosophic Algebraic Structures and Neutrosophic N-Algebraic Structures", Hexis, Phonex, Arizona 2006.

- [8] Kandasamy, V., Kandasamy, I., and Smarandache, F., "Neutrosophic Duplets of $\{Z_p^n, \times\}$ and $\{Z_{pq}, \times\}$ and Their Properties", Symmetry, 2018.
- [9] Yingcang, Ma., Xiaohong Zhang ., Smarandache, F., and Juanjuan, Z., The Structure of Idempotents in Neutrosophic Rings and Neutrosophic Quadruple Rings, Symmetry, Vol. 11, 2019.
- [10] Kandasamy, V., Kandasamy, I., and Smarandache, F., "Neutrosophic Triplets In Neutrosophic Rings", Mathematics, 2019.
- [11] M. Ali, F. Smarandache, M. Shabir and L. Vladareanu, "Generalization of Neutrosophic Rings and Neutrosophic Fields," Neutrosophic Sets and Systems, vol. 5, pp. 9-14, 2014.
- [12] Florentin Smarandache, *Neutrosophic Duplet Structures*, Joint Fall 2017 Meeting of the Texas Section of the APS, Texas Section of the AAPT, and Zone 13 of the Society of Physics Students, The University of Texas at Dallas, Richardson, TX, USA, October 20-21, 2017, <http://meetings.aps.org/Meeting/TSF17/Session/F1.32>
- [13] Abobala, M., "On The Characterization of Maximal and Minimal Ideals In Several Neutrosophic Rings", Neutrosophic Sets and Systems, Vol. 45, 2021.
- [14] Abobala, M., "Neutrosophic Real Inner Product Spaces", Neutrosophic Sets and Systems, Vol. 43, 2021.
- [15] Abobala, M., On Refined Neutrosophic Matrices and Their Applications In Refined Neutrosophic Algebraic Equations, Journal Of Mathematics, Hindawi, 2021
- [16] Abobala, M., On The Representation of Neutrosophic Matrices by Neutrosophic Linear Transformations, Journal of Mathematics, Hindawi, 2021.
- [17] Abobala, M., "On Some Algebraic Properties of n-Refined Neutrosophic Elements and n-Refined Neutrosophic Linear Equations", Mathematical Problems in Engineering, Hindawi, 2021
- [18] Abobala, M., Partial Foundation of Neutrosophic Number Theory, Neutrosophic Sets and Systems, Vol. 39 , 2021.

- [19] Abobala, M., "A Study Of Nil Ideals and Kothe's Conjecture In Neutrosophic Rings", International Journal of Mathematics and Mathematical Sciences, hindawi, 2021
- [20] Abobala, M., and Hatip, A., "An Algebraic Approach To Neutrosophic Euclidean Geometry", Neutrosophic Sets and Systems, Vol. 43, 2021.
- [21] Abobala, M., Bal, M., and Hatip, A., "A Review On Recent Advantages In Algebraic Theory Of Neutrosophic Matrices", International Journal of Neutrosophic Science, Vol. 17, 2021.
- [22] Hajjari, A., and Ali, R., "A Contribution To Kothe's Conjecture In Refined Neutrosophic Rings", International Journal of Neutrosophic Science", Vol. 16, 2021.
- [23] Olgun, N., Hatip, A., Bal, M., and Abobala, M., "A Novel Approach To Necessary and Sufficient Conditions For The Diagonalization of Refined Neutrosophic Matrices", International Journal of Neutrosophic Science, Vol. 16, pp. 72-79, 2021.
- [24] Abobala, M., Hatip, A., Bal, M., "A Study Of Some Neutrosophic Clean Rings", International journal of neutrosophic science, 2022.
- [25] Abobala, M., Bal, M., Aswad, M., "A Short Note On Some Novel Applications of Semi Module Homomorphisms", International journal of neutrosophic science, 2022.
- [26] Ibrahim, M., and Abobala, M., "An Introduction To Refined Neutrosophic Number Theory", Neutrosophic Sets and Systems, Vol. 45, 2021.
- [27] Chalapathi, T., and Kumar, K., "Neutrosophic Units of Neutrosophic Rings and Fields", Neutrosophic Sets and Systems, Vol. 21, 2018.
- [28] Abobala, M., "On Some Special Elements In Neutrosophic Rings and Refined Neutrosophic Rings", Journal of New Theory, vol. 33, 2020.
- [29] Smarandache, F., and Broumi, M., "Neutro-Intelligent Set is a particular case of the refined neutrosophic set", Journal of Neutrosophic and Fuzzy Systems, Vol. 1, 2022.
- [30] Ahmad, K., Bal, M., and Aswad, M., "The kernel of Fuzzy and Anti Fuzzy Groups", Journal of Neutrosophic and Fuzzy Systems, Vol. 1, 2022.
- [31] Ahmad, K., Bal, M., and Aswad, M., "A Short Note on The Solution Of Fermat's Diophantine Equation In Some Neutrosophic Rings", Journal of Neutrosophic and Fuzzy Systems, Vol. 1, 2022.

- [32] Singh, P, K., Ahmad, K., Bal, M., Aswad, M., " On The Symbolic Turiyam Rings", Journal of Neutrosophic and Fuzzy Systems, 2022.
- [33] Sankari, H, and Abobala, M, " A Contribution to m-Power Closed Groups", UMM-Alqura University Journal for Applied Sciences, KSA, 2020.
- [34] Singh, P,K., " Data With Turiyam Set for Fourth Dimension Quantum Information Processing", Journal of Neutrosophic and Fuzzy Systems, vol.1, 2022.
- [35] Singh, P,K., " Anti-geometry and NeutroGeometry Characterization of Non-Euclidean Data", Journal of Neutrosophic and Fuzzy Systems, vol. 1, 2022
- [36] Abobala, M., "On Some Neutrosophic Algebraic Equations", Journal of New Theory, Vol. 33, 2020.
- [37] Abobala, M., A Study of Maximal and Minimal Ideals of n-Refined Neutrosophic Rings, Journal of Fuzzy Extension and Applications, Vol. 2, pp. 16-22, 2021.
- [38] Ahmad, K., Bal, M., Hajjari, A., Ali, R., " On Imperfect Duplets In Some refined Neutrosophic Rings", Journal of Neutrosophic and Fuzzy Systems, 2022.
- [37] **Prem Kumar Singh**, NeutroAlgebra and NeutroGeometry for Dealing Heteroclinic Patterns. In: Theory and Applications of NeutroAlgebras as Generalizations of Classical Algebras, IGI Global Publishers, April 2022, Chapter 6, DOI: 10.4018/978-1-6684-3495-6, ISBN13: 9781668434956
- [38] **Prem Kumar Singh**, AntiGeometry and NeutroGeometry Characterization of Non-Euclidean Data Sets, Journal of Neutrosophic and Fuzzy Systems, Nov 2021, Volume 1, Issue 1, pp. 24-33, ISSN: , DOI: <https://doi.org/10.54216/JNFS.0101012>
- [39] **Prem Kumar Singh**, Three-way n-valued neutrosophic concept lattice at different granulation, International Journal of Machine Learning and Cybernetics, *Springer*, November 2018, Vol 9, Issue 11, pp. 1839-1855,