

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/375632505>

Neutrosophic Treatment of Duality Linear Models and the Binary Simplex Algorithm

Article in *PADMA* · November 2023

DOI: 10.54216/PAMDA.020202

CITATIONS

0

READ

1

2 authors:



Maissam Ahmad Jdid

Damascus University

40 PUBLICATIONS 356 CITATIONS

[SEE PROFILE](#)



Florentin Smarandache

University of New Mexico Gallup

3,783 PUBLICATIONS 46,705 CITATIONS

[SEE PROFILE](#)



Neutrosophic Treatment of Duality Linear Models and the Binary Simplex Algorithm

Maissam Jdid^{*1}, Florentin Smarandache²

¹Faculty of Science, Damascus University, Damascus, Syria

²University of New Mexico •Mathematics, Physics and Natural Sciences Division
705 Gurley Ave., Gallup, NM 87301, USA

Emails: maissam.jdid66@damascusuniversity.edu.sy; smarand@unm.edu

Abstract

One of the most important theories in linear programming is the dualistic theory and its basic idea is that for every linear model has dual linear model, so that solving the original linear model gives a solution to the dual model. Therefore, when we solving the linear programming model, we actually obtain solutions for two linear models. In this research, we present a study of the models. The neutrosophic dual and the binary simplex algorithm, which works to find the optimal solution for the original and dual models at the same time. The importance of this algorithm is evident in that it is relied upon in several operations research topics, such as integer programming algorithms, some nonlinear programming algorithms, and sensitivity analysis in linear programming...

Keywords: Neutrosophic Science; Neutrosophic Linear Models; Dual Models; Neutrosophic dual Linear Models; binary Simplex Algorithm.

1. Introduction:

Most companies and institutions rely on studies provided by experts and researchers using operations research methods in order to ensure a safe work environment away from danger and danger and to give decision makers a margin of freedom, after the studies and research presented using the concepts of neutrosophic science in most fields of science have proven their ability to provide more accurate results. From the results that we were obtaining using classical data-driven studies, we reformulated many operations research topics using neutrosophic concepts [1-20], and to complement what we have done, we present in this research a study of dual linear models and the binary simplex algorithm that gives us a solution for the original and dual models at the same time so that we can provide a clear study that helps decision makers in companies and institutions develop plans and programs through which the greatest profit and lowest cost are achieved.

Discussion:

In our practical life, we encounter many problems that are formulated in the form of linear mathematical models consisting of an objective function and a set of constraints in the form of equations or inequalities. The linear model is written in many formulas that are distinguished by the type of the objective function and the form of the constraints. Many references have been provided that studies of linear and dual linear models are based on classical data, and the appropriate algorithms for solving each of them and how to find the dual model for any model [21-25]. Since classical data results in optimal solutions, which are classical values, restricted values appropriate to the conditions in which the data were collected, and for any change in these conditions, these solutions are inappropriate and may cause the facility a large unexpected loss, so we found it appropriate to reformulate these studies using concepts neutrosophic science is done by taking data with neutrosophic values suitable for all conditions. In previous research, we presented neutrosophic linear models, the direct neutrosophic simplex algorithm, and the graphical method for finding the optimal solution for linear models,[18,19,20] . As a

continuation of what we have done previously, we present in this research a study whose purpose is to reformulate the binary simplex algorithm to find the optimal solution for the original model and the dual model at the same time.

We present this study in two stages:

The first stage:

Finding the neutrosophic dual models using the binary table.

When we want to find the model associated with any linear model, we start by placing this model in the symmetrical form, so we mention the symmetrical form.

The symmetrical form of the neutrosophic linear model: [19]

We say of a linear model that it is in the symmetrical form if all variables are constrained to be non-negative and if all constraints are given in the form of inequalities (and the inequalities of the maximization model constraints must be in the form (\leq less than or equal to) while the inequalities of the minimization model constraints must be in the form (\geq is greater than or equal to), then the linear model is written in the neutrosophic symmetric form in one of two cases:

First case:

$$NZ = \sum_{j=1}^n (c_j \pm \varepsilon_j) x_j \rightarrow \text{Max}$$

Constraints:

$$\sum_{j=1}^n Na_{ij} x_j \leq b_i \pm \delta_i \quad ; \quad i = 1, 2, \dots, m$$

$$x_j \geq 0$$

Second case:

$$NL = \sum_{j=1}^n (c_j \pm \varepsilon_j) x_j \rightarrow \text{Min}$$

Constraints:

$$\sum_{j=1}^n Na_{ij} x_j \geq b_i \pm \delta_i \quad ; \quad i = 1, 2, \dots, m$$

$$x_j \geq 0$$

In both cases, we have $x_j \geq 0$, which are the decision variables, unknown values that we obtain after solving the linear model.

$Nc_j = c_j \pm \varepsilon_j$ and $Nb_i = b_i \pm \delta_i$ and $Na_{ij} = a_{ij} \pm \mu_{ij}$ where ($j=1,2,\dots,n$, $i=1,2,\dots,m$) are the data of the problem under study, and they are neutrosophic values, indefinite values that enjoy a margin of freedom and are taken according to The nature of the situation represented by the linear model

First method:

Constructing neutrosophic dual linear models using tables:

If the original model is given in detailed form, we put it in the symmetrical form as stated in the previous paragraph, then we draw a binary table for the original and dual models according to the following steps:

- We explain the above through the following two cases:**

Find

$$NZ = Nc_1x_1 + Nc_2x_2 + \dots + Nc_nx_n \rightarrow Max$$

Constants:

$$Na_{11}x_1 + Na_{12}x_2 + \cdots + Na_{1n}x_n \leq Nb_1$$

$$Na_{21}x_1 + Na_{22}x_2 + \cdots + Na_{2n}x_n \leq Nb_2$$

$$Na_{m1}x_1 + Na_{m2}x_2 + \cdots + Na_{mn}x_n \leq Nb_m$$

$$x_1, x_2, \dots, x_n \geq 0$$

Table No. (1) Objective follower of the maximization type

Dual vibrable		Original model				
		<div>objective function</div>	$Nc_1x_1 + Nc_2x_2 + \cdots + Nc_nx_n$		<div>Max</div>	
		constants			Constants column	
		y_1	1	$Na_{11}x_1 + Na_{12}x_2 + \cdots + Na_{1n}x_n$	\leq	Nb_1
		y_2	2	$Na_{21}x_1 + Na_{22}x_2 + \cdots + Na_{2n}x_n$	\leq	Nb_2
		\leq	...
		y_m	m	$Na_{m1}x_1 + Na_{m2}x_2 + \cdots + Na_{mn}x_n$	\leq	Nb_m
		Non-negativity constraints		x_1, x_2, \dots, x_n	\geq	0
		Dual model				
		<div>Objective function constraints</div>	$Nb_1y_1 + Nb_2y_2 + \cdots + Nb_iy_m$		<div>Min</div>	Constants column
	1	$Na_{11}y_1 + Na_{21}y_2 + \cdots + Na_{m1}y_m$	\geq	Nc_1		
	2	$Na_{12}y_1 + Na_{22}y_2 + \cdots + Na_{m2}y_m$	\geq	Nc_2		
	\geq	...		
	n	$Na_{1n}y_1 + Na_{2n}y_2 + \cdots + Na_{mn}y_m$	\geq	Nc_n		

Non-negativity constraints	y_1, y_2, \dots, y_m	\geq	0
----------------------------	------------------------	--------	---

The second case: The original model is symmetrical and of the reduction type:

Find

$$NL = Nc_1x_1 + Nc_2x_2 + \dots + Nc_nx_n \rightarrow \text{Min}$$

Constants:

$$Na_{11}x_1 + Na_{12}x_2 + \dots + Na_{1n}x_n \geq Nb_1$$

$$Na_{21}x_1 + Na_{22}x_2 + \dots + Na_{2n}x_n \geq Nb_2$$

$$\dots \dots \dots$$

$$Na_{m1}x_1 + Na_{m2}x_2 + \dots + Na_{mn}x_n \geq Nb_m$$

$$x_1, x_2, \dots, x_n \geq 0$$

The binary table for the original model and the accompanying model is as follows:

Table No. (2) objective follower in the original model of the reduce type

Original model					
Dual vibrable		objective function	$Nc_1x_1 + Nc_2x_2 + \dots + Nc_nx_n$		Min Constants column
		constants			
	y_1	1	$Na_{11}x_1 + Na_{12}x_2 + \dots + Na_{1n}x_n$	\geq	Nb_1
	y_2	2	$Na_{21}x_1 + Na_{22}x_2 + \dots + Na_{2n}x_n$	\geq	Nb_2
	$\dots \dots \dots$	\geq	...
	y_m	m	$Na_{m1}x_1 + Na_{m2}x_2 + \dots + Na_{mn}x_n$	\geq	Nb_m
Dual model					
		objective function	$Nb_1y_1 + Nb_2y_2 + \dots + Nb_my_m$		Max Constants column
		constants			
		1	$Na_{11}y_1 + Na_{21}y_2 + \dots + Na_{m1}y_m$	\leq	Nc_1
		2	$Na_{12}y_1 + Na_{22}y_2 + \dots + Na_{m2}y_m$	\leq	Nc_2
		...	$\dots \dots \dots$	\leq	...
		n	$Na_{1n}y_1 + Na_{2n}y_2 + \dots + Na_{mn}y_m$	\leq	Nc_n
		Non-negativity constraints	y_1, y_2, \dots, y_m	\geq	0

The second stage: formulation of the binary neutrosophic algorithm.

The binary simplex algorithm (for the original and dual models) neutrosophic. Through this algorithm, we can find the two ideal solutions for both the original and dual models at the same time. Before starting the binary simplex algorithm, we must mention the modified simplex algorithm that we will use within the steps of the binary algorithm.

$$\begin{aligned} Na_{11}y_1 + Na_{21}y_2 + \dots + Na_{m1}y_m - v_1 &= Nc_1 \\ Na_{12}y_1 + Na_{22}y_2 + \dots + Na_{m2}y_m - v_2 &= Nc_2 \\ &\vdots \\ Na_{1n}y_1 + Na_{2n}y_2 + \dots + Na_{mn}y_m - v_n &= Nc_n \\ y_i &\geq 0 ; i = 1, 2, \dots, m \\ v_j &\geq 0 ; j = 1, 2, \dots, n \end{aligned}$$

The two models have the same coefficients, and the matrix of instances of the dual model is the transpose of the matrix of instances of the original model. We write the two models in the following binary table:

Dual vibrable		Original model			
		<div> <div>objective function</div> <div>constants</div> </div>	$Nc_1x_1 + Nc_2x_2 + \dots + Nc_nx_n + 0u_1 + 0u_2 + \dots + 0u_m$		<div> <div>Max</div> <div>Constants column</div> </div>
		1	$Na_{11}x_1 + Na_{12}x_2 + \dots + Na_{1n}x_n + u_1$	=	Nb_1
		y_2	$Na_{21}x_1 + Na_{22}x_2 + \dots + Na_{2n}x_n + u_2$	=	Nb_2
		...	$\dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots$	=	...
		y_m	$Na_{m1}x_1 + Na_{m2}x_2 + \dots + Na_{mn}x_n + u_m$	=	Nb_m
		Non-negativity constraints	$x_1, x_2, \dots, x_n, u_1, u_2, \dots, u_m$	\geq	0
Dual model					
		<div> <div>objective function</div> <div>constants</div> </div>	$Nb_1y_1 + Nb_2y_2 + \dots + Nb_iy_m + 0v_1 + 0v_2 + \dots + 0v_n$		<div> <div>Min</div> <div>Constants column</div> </div>
		1	$Na_{11}y_1 + Na_{21}y_2 + \dots + Na_{m1}y_m - v_1$	=	Nc_1
		2	$Na_{12}y_1 + Na_{22}y_2 + \dots + Na_{m2}y_m - v_2$	=	Nc_2
		...	$\dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots \dots$	=	...
		n	$Na_{1n}y_1 + Na_{2n}y_2 + \dots + Na_{mn}y_m - v_n$	=	Nc_n
		Non-negativity constraints	$y_1, y_2, \dots, y_m, v_1, v_2, \dots, v_n$	\geq	0

- Table No. (4): The binary table for the original and dual models according to the modified Simplex algorithm

		Basic variables with a (-) sign in the dual model				Follow the objective of the dual model NB_i
		$-v_1$	$-v_2$...	$-v_n$	
<div> <div>Non-basic vibrable</div> <div>basic vibrable</div> </div>		x_1	x_2	...	x_n	
		Na_{11}	Na_{12}	...	Na_{1n}	Nb_1
No n-	y_1	u_1				

	y_2	u_2	Na_{21}	Na_{22}	...	Na_{2n}	Nb_2

	y_m	u_m	Na_{m1}	Na_{m2}	...	Na_{mn}	Nb_m
	objective of the original model		Nc_1	Nc_2	...	Nc_n	$L - 0 \rightarrow Min$ $Z - 0 \rightarrow Max$

Binary simplex algorithm for the original and dual models:

From the modified simplex algorithm of the original model, we obtain the optimal solution of the original model when all the elements are in the last row (the objective function row of the original model) $Nc_j \leq 0$; $j = 1, 2, \dots, n$ and at the same time all the elements are in the column The last (associated objective function column) $Nb_i \geq 0$; $i = 1, 2, \dots, m$ and we get the optimal solution for the dual model when all elements in the last column (associated objective function column) are $Nb_i \geq 0$; $i = 1, 2, \dots, m$ and at the same time the last row (the objective function row of the original model) $Nc_j \leq 0$; $j = 1, 2, \dots, n$ (because it will correspond to $Nc_j = -v_j$) which are the two conditions Same for both models. Therefore, when searching for the optimal solution for both models together, we must work to make all elements $Nb_i \geq 0$; $i = 1, 2, \dots, m$ and to make all elements $Nc_j \leq 0$; $j = 1, 2, \dots, n$ to achieve this, we rely on one of the two models, put its variables and coefficients in a table, and place the dual model in an external frame of that table. In general, we find that the necessity of placing the two models in a short table does not allow us to get rid of the negative constants on the right side, and therefore the general case of the previous binary table can it must include negative constants $Nb_i < 0$, and the elements of the last row can include positive elements $Nc_j > 0$, so when searching for the optimal solution for the two models, we must work to address these elements based on one of the two models.

Depending on the original model, we do this in two stages:

The first stage:

We make the constant Nb_i non-negative, which corresponds to obtaining a non-negative basic solution for the original model.

The second stage:

We make all elements of the objective function row non-positive (in the case of the objective function, maximization), and this corresponds to obtaining the optimal solution required for the original model.

Based on the dual model, we do this in two stages:

The first stage:

We must make the elements of the dual model objective function column $Nb_i \geq 0$; $i = 1, 2, \dots, m$,The last row is non-negative

The second stage:

We must make the free constants for the dual model $-Nc_j$ non-positive, and this corresponds to obtaining the optimal solution for the dual model. We explain the above through the following example:

Find the optimal solution for both the following neutrosophic linear model and its dual using the binary algorithm

Example:

Find:

$$[5,8] x_1 + [3,6] x_2 \rightarrow Max$$

Constans:

$$2x_1 + 3x_2 \leq [14,20]$$

$$2x_1 + x_2 \leq [10,16]$$

$$3x_2 \leq [12,18]$$

$$3x_1 \leq [15,21]$$

$$x_1 \geq 0, x_2 \geq 0$$

We form the binary table of the model and the dual model:

Table No. (5) The original model and its dual mode

Dual vibrable		Original model				
		objective function constants	[5,8] x_1 + [3,6] x_2		<div>Max Constants column</div>	
		y_1	1	$2x_1 + 3x_2$	\leq	[14,20]
		y_2	2	$2x_1 + x_2$	\leq	[10,16]
		y_3	3	$3x_2$	\leq	[12,18]
		y_4	4	$3x_1$	\leq	[15,21]
		Non-negativity constraints		x_1, x_2	\geq	0
		Dual model				
		objective function constants	[14,20] y_1 + [10,16] y_2 + [12,18] y_3 + [15,21] y_4		<div>Min Constants column</div>	
		1	$2y_1 + 2y_2 + 3y_4$	\geq	[5,8]	
		2	$3y_1 + y_2 + 3y_4$	\geq	[3,6]	
		Non-negativity constraints		y_1, y_2, y_3, y_4	\geq	0

In the following table, we wrote the two models in standard form:

		Original model			
		objective function constants	$[5,8] \, x_1 + [3,6] \, x_2 + 0u_1 + 0u_2 + 0u_3 + 0u_4$		Max Constants column
Dual vibrable	y_1	1	$2x_1 + 3x_2 + u_1$	=	[14,20]
	y_2	2	$2x_1 + x_2 + u_2$	=	[10,16]
	y_3	3	$3x_2 + u_3$	=	[12,18]
	y_4	4	$3x_1 + u_4$	=	[15,21]
	Non-negativity constraints		$x_1, x_2, u_1, u_2, u_3, u_4$	\geq	0
	Dual model				
		objective function constants	$[14,20]y_1 + [10,16]y_2 + [12,18]y_3 + [15,21]y_4 + 0v_1 + 0v_2$		Min Constants column
		1	$2y_1 + 2y_2 + 3y_4 - v_1$	=	[5,8]
		2	$3y_1 + y_2 + 3y_4 - v_2$	=	[3,6]

Non-negativity constraints	$y_1, y_2, y_3, y_4, v_1, v_2$	\geq	0
----------------------------	--------------------------------	--------	---

Table No. (6) Standard format for the original model and the dual model

We notice from the table that the standard form of the original model includes a ready-made base of additional variables u_1, u_2, u_3, u_4 , but for the dual model there is no ready-made base. Therefore, we multiply the two restrictions by (-1) and we obtain the basic form of the dual model.

The following table shows the basic form of the original and dual models:

Table No. (7): The basic shape of the original model and the dual model

Dual vibrable		Original model					
		objective function constants	$[5,8] \, x_1 + [3,6] \, x_2 + 0u_1 + 0u_2 + 0u_3 + 0u_4$		<i>Max</i> Constants column		
		y_1	1	$2x_1 + 3x_2 + u_1$	=	[14,20]	
		y_2	2	$2x_1 + x_2 + u_2$	=	[10,16]	
		y_3	3	$3x_2 + u_3$	=	[12,18]	
		y_4	4	$3x_1 + u_4$	=	[15,21]	
		Non-negativity constraints		$x_1, x_2, u_1, u_2, u_3, u_4$		\geq	0
Dual model							
		objective function constants	$[14,20]y_1 + [10,16]y_2 + [12,18]y_3 + [15,21]y_4 + 0v_1 + 0v_2$		<i>Min</i> Constants column		
		1	$-2y_1 - 2y_2 - 3y_4 + v_1$	=	-[5,8]		
		2	$-3y_1 - y_2 - 3y_4 + v_2$	=	-[3,6]		
		Non-negativity constraints		$y_1, y_2, y_3, y_4, v_1, v_2$		\geq	0

We put the two models in the modified Simplex algorithm table and we get the following table:

Table No. (8): The binary table for the original and dual models according to the modified Simplex algorithm

According to the original model					
		$-v_1$	$-v_2$		
		Non-basic vibrable	basic vibrable	objective function	
				NB_i	
Non-basic vibrable	y_1	u_1	2	3	[14,20]
	y_2	u_2	2	1	[10,16]
	y_3	u_3	0	3	[12,18]
	y_4	u_4	3	0	[15,21]
	objective function		[5,8]	[3,6]	$L - 0$
		NC_i			$Z - 0$

		According to the dual model					
		u_1	u_2	u_3	u_4		
<div>Non-basic vibrable basic vibrable</div>		y_1	y_2	y_3	y_4	objective function Original model	
		Nc_i					
Non-basic	x_1	v_1	-2	-2	0	-3	-[5,8]
	x_2	v_2	-3	-1	-3	0	-[3,6]
	objective function Dual model NB_i		[14,20]	[10,16]	[12,18]	[15,21]	<div>$Z - 0$</div> <div>$L - 0$</div>

The first stage:

1- For the original model:

Since the values in the constant's column are all positive, we study the values in the objective function row and determine the largest positive value. We find:

$$\max([5,8], [3,6]) = [5,8]$$

It is an expression of the variable x_1 . This means that it will enter the base. To determine the element that will exit from the base, we calculate the index $N\theta$, where:

$$N\theta \in \min \left[\frac{[14,20]}{2}, \frac{[10,16]}{2}, \frac{[15,21]}{3} \right] = \frac{[15,21]}{3} = [5,7]$$

We find that the pivot column is the column of the non-base variable x_1 , meaning that the variable x_1 will enter the base instead of the variable u_4 , and the pivot element is the element resulting from the intersection of the pivot row and the pivot column, which is (3)

We perform the switching between variables using a modified simplex algorithm.

2- For the dual model:

We study the elements of the objective function row. We notice that all the values are positive. Therefore, we study the elements of the constant's column. We find that they are all negative values. We choose the most negative of them, which is $-[5,8]$ which is the row of the base variable v_1 , so its row is the pivot row. To determine the pivot column and the pivot element, we calculate the index $N\theta'$ where:

$$N\theta' \in \max \left[\frac{[14,20]}{-2}, \frac{[10,16]}{-2}, \frac{[15,21]}{-3} \right] = \frac{[15,21]}{-3}$$

So, the column of the non-base variable u_4 is the pivot column, meaning that the variable u_4 will enter the base instead of the variable v_1 , and the pivot element is the element resulting from the intersection of the pivot row and the pivot column, which is (-3). We perform the switching between the variables using the modified simplex algorithm, from (1) and (2) We get the following double table:

Table No. (9): The binary table for the first stage, the solution according to the original and dual models

		According to the original model		objective function Original model NB_i
		$-y_4$	$-v_2$	
Non-basic vibrable basic vibrable		u_4	x_2	

Non-basic vibrable	y_1	u_1	$-\frac{2}{3}$		3	[4,6]	
	y_2	u_2	$-\frac{2}{3}$		1	[0,4]	
	y_3	u_3	0		3	[12,18]	
	v_1	x_1	$\frac{1}{3}$		0	[5,7]	
		objective function Original model Nc_i	$\left[\frac{-8}{3}, \frac{-5}{3}\right]$		[3,6]	<div>$L - [25,56]$ $Z - [25,56]$</div>	
		According to the dual model					
			u_1	u_2	u_3	x_1	
<div>Non-basic vibrable basic vibrable</div>		y_1	y_2	y_3	v_1	objective function Original model	
		Nc_i					
Non-basic vibrable	u_4	y_4	$\frac{2}{3}$	$\frac{2}{3}$	0	$-\frac{1}{3}$	$\left[\frac{8}{3}, \frac{5}{3}\right]$
	x_2	v_2	-3	-1	-3	0	-[3,6]
		objective function Original model Nc_i	[4,6]	[0,4]	[12,18]	[5,7]	<div>$Z - [25,56]$ $L - [25,56]$</div>

The second phase:

We apply the stopping criterion of the algorithm

For the original model:

Since the values in the constant's column are all positive, we study the values in the objective function row. We notice that there is a positive value, which is [3,6], meaning that we have not yet reached the optimal solution. Therefore, we specify the pivot column, which is the column of the variable x_2 corresponding to the only positive value in the objective function row. [3,6] In order to determine the pivot row and the pivot element, we calculate the index $N\theta$, where:

$$N\theta \in \min \left[\frac{[4,6]}{3}, \frac{[0,4]}{1}, \frac{[12,18]}{3} \right] = \frac{[4,6]}{3}$$

It corresponds to the base element u_1 , so its row is the pivot row and the pivot element is (3). We swap between the variables using the modified simplex algorithm.

For the dual model:

We study the elements of the objective function row. We notice that all the values are positive. Therefore, we study the elements of the constant's column. We find that there is a single negative value, which is $-[3,6]$, which is the line of the base variable v_2 , so its row is the pivot row. To determine the pivot column and the pivot element, we calculate the index $N\theta'$ where:

$$N\theta' \in \max \left[\frac{[4,6]}{-3}, \frac{[0,4]}{-1}, \frac{[12,18]}{-3} \right] = \frac{[4,6]}{-3}$$

So, the column of the non-base variable y_1 is the pivot column, meaning that the variable y_1 will enter the base instead of the variable v_2 , and the pivot element is the element resulting from the intersection of the pivot row and

the pivot column, which is (-3) . We perform the switching between the variables using the modified simplex algorithm, from (1) and (2) We get the following double table:

Table No. (10): The binary algorithm table for the second stage

		According to the original model					
		$-y_4$		$-y_1$			
		u_4	u_1		objective function Dual model		
		Non-basic vibrable			NB_i		
		basic vibrable					
Non-basic vibrable	v_2	x_2	$-\frac{2}{9}$	$\frac{1}{3}$	$[\frac{4}{3}, 2]$		
	y_2	u_2	$-\frac{4}{9}$	$-\frac{1}{3}$	$[\frac{4}{3}, 2]$		
	y_3	u_3	$\frac{2}{3}$	-1	$[8, 12]$		
	v_1	x_1	1	0	$[5, 7]$		
		objective function Original model Nc_i	$[-6, -1]$		$[-2, -1]$		
		According to the dual model					
			x_2	u_2	u_3	x_1	
		Non-basic vibrable basic vibrable	v_2	y_2	y_3	v_1	objective function Original model
			Nc_i				
Non-basic vibrable	u_4	y_4	$\frac{2}{9}$	$\frac{4}{9}$	$-\frac{2}{3}$	-1	$[-6, -1]$
	u_1	y_1	$-\frac{1}{3}$	$\frac{1}{3}$	1	0	$[-2, -1]$
		objective function Dual model NB_i	$[\frac{4}{3}, 2]$	$[\frac{4}{3}, 2]$	$[8, 12]$	$[5, 7]$	$Z - [29, 68]$ $L - [29, 68]$

We apply the stopping criterion of the algorithm:

- 1- For the original model, we study the elements of the objective function row until the criterion for stopping the algorithm is met, which is the absence of any positive element
- 2- For the dual model, we also study the elements of the constants column until the criterion for stopping the algorithm is met, which is the absence of any negative element
- 3- We find that the criterion has been met and thus we have reached the optimal solution

The optimal solution of the original model is:

$$x_2^* \in [\frac{4}{3}, 2], u_2^* \in [\frac{4}{3}, 2], u_3^* \in [8, 12], x_1^* \in [5, 7], u_1^* = u_4^* = 0$$

The value of the objective function corresponds to:

$$Z^* = \max Z \in [29,68]$$

The optimal solution of the dual model is:

$$y_1^* \in [1,2] , y_4^* \in [1,6] , v_2^* = y_2^* = y_3^* = v_1^* = 0$$

The value of the objective function corresponds to:

$$L^* = \min L \in [29,166]$$

We note that:

$$Z^* = \max Z \in [29,68] \leq L^* = \min L \in [29,166]$$

This solution is acceptable according to the following theory:

If (x_1, x_2, \dots, x_n) is an acceptable solution to the original model of type *Max* and (y_1, y, \dots, y_m) was an acceptable solution for the dual model of type *Min*, so the value of the objective function of the original model does not exceed the value of the objective function of the dual model for these two solutions, that is, it is

$$\sum_{j=1}^n Nc_j x_j \leq \sum_{i=1}^m Nb_i y_i$$

This is for all acceptable solutions for both models (including the optimal solution)

5. Conclusion and Results:

From the previous study, we arrived at a solution for the original and dual models at the same time, which are neutrosophic values from which we know the minimum and maximum profit that we can obtain, because the interpretation of the optimal solution for the original model is that it gives us the best production plan that makes the value of that production as large as possible, within Available capabilities. As for the optimal solution for the dual model, it gives us the best values for the prices of raw materials, which, if used without waste, will also give us the best production plan, and the result is the maximum profit.

References

- [1] Florentin Smarandache, Maissam Jdid, On Overview of Neutrosophic and Plithogenic Theories and Applications, Applied Mathematics and Data Analysis, Vo .2, No .1, 2023
- [2] Maissam Jdid- Hla Hasan, The state of Risk and Optimum Decision According to Neutrosophic Rules, International Journal of Neutrosophic Science (IJNS), Vol. 20, No.1,2023
- [3] Mohammed Alshikho, Maissam Jdid, Said Broumi, Artificial Intelligence and Neutrosophic Machine learning in the Diagnosis and Detection of COVID 19, Journal Prospects for Applied Mathematics and Data Analysis, Vol 01, No,02 USA,2023
- [4] Maissam Jdid, Neutrosophic Nonlinear Models, Journal Prospects for Applied Mathematics and Data Analysis, Vo .2, No .1, 2023
- [5] Maissam Jdid, Neutrosophic Mathematical Model of Product Mixture Problem Using Binary Integer Mutant, Journal of Neutrosophic and Fuzzy Systems (JNFS), Vo .6, No .2, 2023
- [6] Maissam Jdid, The Use of Neutrosophic linear Programming Method in the Field of Education, Handbook of Research on the Applications of Neutrosophic Sets Theory and Their Extensions in Education, Chapter 15, IGI-Global,2023
- [7] Maissam Jdid, Florentin Smarandache, Said Broumi, Inspection Assignment Form for Product Quality Control, Journal of Neutrosophic Systems with Applications, Vol. 1, 2023

- [8] Maissam Jdid, Said Broumi, Neutrosophic Rejection and Acceptance Method for the Generation of Random Variables, Neutrosophic Sets and Systems, NSS, Vol.56,2023
- [9] Maissam Jdid, Florentin Smarandache, The Use of Neutrosophic Methods of Operation Research in the Management of Corporate Work, Journal of Neutrosophic Systems with Applications, Vol. 3, 2023
- [10] Maissam Jdid, Florentin Smarandache, Lagrange Multipliers and Neutrosophic Nonlinear Programming Problems Constrained by Equality Constraints Journal of Neutrosophic Systems with Applications, Vol. 6, 2023
- [11] Maissam Jdid, Florentin Smarandache, Optimal Neutrosophic Assignment and the Hungarian Method, Journal Neutrosophic Sets and Systems, NSS Vol.57,2023
- [12] Maissam Jdid, Florentin Smarandache, Graphical Method for Solving Neutrosophical Nonlinear Programming Models, Journal of Neutrosophic Systems with Applications, Vol. 9, 2023
- [13] Maissam Jdid, Florentin Smarandache, Optimal Agricultural Land Use: An Efficient Neutrosophic Linear Programming Method, Journal of Neutrosophic Systems with Applications, Vol. 10, 2023
- [14] Maissam Jdid, Nada A Nabeeh Generating Random Variables that follow the Beta Distribution Using the Neutrosophic Acceptance-Rejection Method *1, Journal Neutrosophic Sets and Systems, NSS Vol.58,2023
- [15] Maissam Jdid, Studying Transport Models with the Shortest Time According to the Neutrosophic Logic, Journal Neutrosophic Sets and Systems, NSS Vol.58,2023
- [16] Florentin Smarandache, Maissam Jdid, NEUTROSOPHIC TRANSPORT AND ASSIGNMENT ISSUES, Publisher: Global Knowledge's: 978_1_59973_769_0
- [17] Maissam Jdid, NEUTROSOPHIC TRANSPORT AND ASSIGNMENT ISSUES, Publisher: Global Knowledge's: 978_1_59973_770_6, (Arabic version).
- [18] Maissam Jdid, AA Salama, Huda E Khalid ,Neutrosophic Handling of the Simplex Direct Algorithm to Define the Optimal Solution in Linear Programming ,International Journal of Neutrosophic Science, Vol.18,No. 1, 2022
- [19] Maissam Jdid, Huda E Khalid, Mysterious Neutrosophic Linear Models, International Journal of Neutrosophic Science, Vol.18, No. 2, 2022
- [20] Maissam Jdid, Florentin Smarandache, The Graphical Method for Finding the Optimal Solution for Neutrosophic linear Models and Taking Advantage of Non-Negativity Constraints to Find the Optimal Solution for Some Neutrosophic linear Models in Which the Number of Unknowns is More than Three, Journal Neutrosophic Sets and Systems, NSS Vol.58,2023
- [21] Alali. Ibrahim Muhammad, Operations Research. Tishreen University Publications, 2004. (Arabic version).
- [22] Al Hamid. Mohammed Dabbas, Mathematical programming, Aleppo University, Syria, 2010. (Arabic version).
- [23] Linear and Nonlinear Programming-DavidG. Luenbrgrr.YinyuYe- Springer Science + Business Media- 2015.
- [24] Maissam Jdid, Bi-Linear Model of Natural Resource Use Published in Tver State University, 2002
- [25] Maissam Jdid Operations Research, Faculty of Informatics Engineering, Al-Sham Private University Publications, 2021.