

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/380667701>

The Mathematical Formulas for Inverting Plithogenic Matrices of Special Orders Between 20 and 24

Article in International Journal of Neutrosophic Science · May 2024

DOI: 10.54216/IJNS.240309

CITATIONS

0

READS

13

7 authors, including:



[wael mahmoud Salameh](#)

Abu Dhabi University

6 PUBLICATIONS 3 CITATIONS

[SEE PROFILE](#)



[Heba Alrawashdeh](#)

University of Jordan

13 PUBLICATIONS 4 CITATIONS

[SEE PROFILE](#)



[Amani Shatarah](#)

Taibah University

14 PUBLICATIONS 25 CITATIONS

[SEE PROFILE](#)



The Mathematical Formulas for Inverting Plithogenic Matrices of Special Orders Between 20 and 24

Ahmad A.Abubaker¹, Wael Mahmoud Mohammad Salameh^{2*}, Heba Alrawashdeh³, Amani Shatarah⁴,
Norah Mousa Alrayes⁵, Abdallah Al-Husban⁶

¹Faculty of Computer Studies, Arab Open University, Riyadh 11681, Saudi Arabia

²Faculty of Information Technology, Abu Dhabi university, Abu Dhabi, UAE

³Faculty of Computer Information Systems, the University of Jordan, Aqaba, Jordan

⁴Department of mathematics, college of science, Taibah university, Al-Madinah, Saudi Arabia

⁵Department of mathematics, college of science, Taibah university, Al-Madinah, Saudi Arabia

⁶Department of Mathematics, Faculty of Science and Technology, Irbid National University, P.O. Box: 2600 Irbid, Jordan

Emails: a.abubaker@arabou.edu.sa; wael.salameh@adu.ac.ae; hsrawashdeh@yahoo.com;
ashatarah@taibahu.edu.sa; nrees@taibahu.edu.sa; dralhosban@inu.edu.jo

Abstract

This paper is dedicated to study the Invertibility properties of all plithogenic square matrices of high orders between 20 and 24, where we present the mathematical conditions and formulas for computing the inverses of all plithogenic square matrices with real plithogenic entries. This goal will be completed by proving many theorems that describe the Invertibility of all classic matrix parts of the corresponding symbolic m-plithogenic matrix.

Keywords: 20-plithogenic matrix; invertible plithogenic matrix; plithogenic determinant.

1. Introduction

Plithogenic sets are considered as a new generalization of fuzzy sets presented by Smarandache in [2], and similar to the neutrosophic sets used in the study of commutative and non-commutative algebraic structures [12-16, 20-22], these sets have been used in linear algebra and the theory of spaces [3-4, 23-25], and even in number theory [17-18] and algebraic equations [5-6, 9].

The matrices of order 2 in [7], and of order 3 and order 4 in [8] were studied, where the basic theorems describing the values and special vectors, as well as the determinants associated with these matrices, as well as the necessary and sufficient orthogonality conditions for these matrices, were formulated, through the properties of the partial matrices from which these matrices are formed.

Recently, the studies have been expanded and their results have been generalized by many researchers to include matrices of this type of higher orders starting from order 5 and ending with 19 [19, 26-27].

These generalized studies have motivated us to discuss one of the most important issues related to these matrices, which is the issue of finding algebraic methods, where we find and prove mathematical formulas that express the method of calculating the inverse for the intended high orders.

2. Main Results

Theorem

Let $\aleph = \aleph_0 + \sum_{i=1}^{20} \aleph_i P_i$ be a 20-plithogenic matrix of size $n \times n$, hence:

1. $\aleph$ is invertible if and only if $\det \aleph$ is an invertible 20-plithogenic real number.
2. $\aleph^{-1} = \aleph_0^{-1} + [(\sum_{i=0}^1 \aleph_i)^{-1} - \aleph_0^{-1}]P_1 + [(\sum_{i=0}^2 \aleph_i)^{-1} - (\sum_{i=0}^1 \aleph_i)^{-1}]P_2 + [(\sum_{i=0}^3 \aleph_i)^{-1} - (\sum_{i=0}^2 \aleph_i)^{-1}]P_3 + [(\sum_{i=0}^4 \aleph_i)^{-1} - (\sum_{i=0}^3 \aleph_i)^{-1}]P_4 + [(\sum_{i=0}^5 \aleph_i)^{-1} - (\sum_{i=0}^4 \aleph_i)^{-1}]P_5 + [(\sum_{i=0}^6 \aleph_i)^{-1} - (\sum_{i=0}^5 \aleph_i)^{-1}]P_6 + \dots$

$$\begin{aligned}
& (\sum_{i=0}^5 \mathfrak{K}_i)^{-1} P_6 + [(\sum_{i=1}^7 \mathfrak{K}_i)^{-1} - (\sum_{i=0}^6 \mathfrak{K}_i)^{-1}] P_7 + [(\sum_{i=1}^8 \mathfrak{K}_i)^{-1} - (\sum_{i=0}^7 \mathfrak{K}_i)^{-1}] P_8 + [(\sum_{i=1}^9 \mathfrak{K}_i)^{-1} - \\
& (\sum_{i=0}^8 \mathfrak{K}_i)^{-1}] P_9 + [(\sum_{i=1}^{10} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^9 \mathfrak{K}_i)^{-1}] P_{10} + [(\sum_{i=1}^{11} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{10} \mathfrak{K}_i)^{-1}] P_{11} + [(\sum_{i=1}^{12} \mathfrak{K}_i)^{-1} - \\
& (\sum_{i=0}^{11} \tau_i)^{-1}] P_{12} + [(\sum_{i=1}^{13} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{12} \mathfrak{K}_i)^{-1}] P_{13} + [(\sum_{i=1}^{14} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{13} \mathfrak{K}_i)^{-1}] P_{14} + [(\sum_{i=1}^{15} \mathfrak{K}_i)^{-1} - \\
& (\sum_{i=0}^{14} \mathfrak{K}_i)^{-1}] P_{15} + [(\sum_{i=1}^{16} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{15} \mathfrak{K}_i)^{-1}] P_{16} + [(\sum_{i=1}^{17} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{16} \mathfrak{K}_i)^{-1}] P_{17} + \\
& [(\sum_{i=1}^{18} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{17} \mathfrak{K}_i)^{-1}] P_{18} + [(\sum_{i=1}^{19} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{18} \mathfrak{K}_i)^{-1}] P_{19} + [(\sum_{i=1}^{20} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{19} \mathfrak{K}_i)^{-1}] P_{20}.
\end{aligned}$$

Proof

1). Let $\mathfrak{K} = \mathfrak{K}_0 + \sum_{i=1}^{20} \mathfrak{K}_i P_i$, then $\mathfrak{K}$ is invertible if and only if there exists $G = G_0 + \sum_{i=1}^{20} G_i P_i$ such that:
 $\mathfrak{K} \times G = U_{n \times n}$, hence:

$$\left\{ \begin{array}{l}
 \aleph_0 G_0 = U_{n \times n} \\
 \sum_{i=0}^1 \aleph_i \sum_{i=0}^1 I_i - \aleph_0 G_0 = O_{n \times n} \\
 \sum_{i=0}^2 \aleph_i \sum_{i=0}^2 G_i - \sum_{i=0}^1 \aleph_i \sum_{i=0}^1 G_i = O_{n \times n} \\
 \sum_{i=0}^3 \aleph_i \sum_{i=0}^3 G_i - \sum_{i=0}^2 \aleph_i \sum_{i=0}^2 G_i = O_{n \times n} \\
 \sum_{i=0}^4 \aleph_i \sum_{i=0}^4 G_i - \sum_{i=0}^3 \aleph_i \sum_{i=0}^3 I_i = O_{n \times n} \\
 \sum_{i=0}^5 \aleph_i \sum_{i=0}^5 G_i - \sum_{i=0}^4 \aleph_i \sum_{i=0}^4 I_i = O_{n \times n} \\
 \sum_{i=0}^6 \aleph_i \sum_{i=0}^6 G_i - \sum_{i=0}^5 \aleph_i \sum_{i=0}^5 G_i = O_{n \times n} \\
 \sum_{i=0}^7 \tau_i \sum_{i=0}^7 G_i - \sum_{i=0}^6 \aleph_i \sum_{i=0}^6 G_i = O_{n \times n} \\
 \sum_{i=0}^8 \aleph_i \sum_{i=0}^8 G_i - \sum_{i=0}^7 \aleph_i \sum_{i=0}^7 G_i = O_{n \times n} \\
 \sum_{i=0}^9 \aleph_i \sum_{i=0}^9 G_i - \sum_{i=0}^8 \aleph_i \sum_{i=0}^8 G_i = O_{n \times n} \\
 \sum_{i=0}^{10} \aleph_i \sum_{i=0}^{10} G_i - \sum_{i=0}^9 \aleph_i \sum_{i=0}^9 G_i = O_{n \times n} \\
 \sum_{i=0}^{11} \aleph_i \sum_{i=0}^{11} G_i - \sum_{i=0}^{10} \aleph_i \sum_{i=0}^{10} G_i = O_{n \times n} \\
 \sum_{i=0}^{12} \aleph_i \sum_{i=0}^{12} G_i - \sum_{i=0}^{11} \aleph_i \sum_{i=0}^{11} G_i = O_{n \times n} \\
 \sum_{i=0}^{13} \aleph_i \sum_{i=0}^{13} G_i - \sum_{i=0}^{12} \aleph_i \sum_{i=0}^{12} G_i = O_{n \times n} \\
 \sum_{i=0}^{14} \aleph_i \sum_{i=0}^{14} G_i - \sum_{i=0}^{13} \aleph_i \sum_{i=0}^{13} G_i = O_{n \times n} \\
 \sum_{i=0}^{15} \aleph_i \sum_{i=0}^{15} G_i - \sum_{i=0}^{14} \aleph_i \sum_{i=0}^{14} G_i = O_{n \times n} \\
 \sum_{i=0}^{16} \aleph_i \sum_{i=0}^{16} G_i - \sum_{i=0}^{15} \aleph_i \sum_{i=0}^{15} G_i = O_{n \times n} \\
 \sum_{i=0}^{17} \aleph_i \sum_{i=0}^{17} G_i - \sum_{i=0}^{16} \aleph_i \sum_{i=0}^{16} G_i = O_{n \times n} \\
 \sum_{i=0}^{18} \aleph_i \sum_{i=0}^{18} G_i - \sum_{i=0}^{17} \aleph_i \sum_{i=0}^{17} G_i = O_{n \times n} \\
 \sum_{i=0}^{19} \aleph_i \sum_{i=0}^{19} G_i - \sum_{i=0}^{18} \aleph_i \sum_{i=0}^{18} G_i = O_{n \times n} \\
 \sum_{i=0}^{20} \aleph_i \sum_{i=0}^{20} G_i - \sum_{i=0}^{19} \aleph_i \sum_{i=0}^{19} G_i = O_{n \times n}
 \end{array} \right.$$

This implies that:

$$\begin{cases} \mathfrak{K}_0 G_0 = U_{n \times n} \\ \sum_{i=0}^q \mathfrak{K}_i \sum_{i=0}^q G_i = U_{n \times n} ; 1 \leq q \leq 20 \end{cases}$$

Hence $\det(\sum_{i=0}^q \mathfrak{K}_i) \neq 0$ for all $1 \leq q \leq 20$, so that $\det(\tau)$ is invertible in $20 - SP_R$.

2). $\sum_{i=0}^q G_i = (\sum_{i=0}^q \mathfrak{K}_i)^{-1}$ for $1 \leq q \leq 20$, therefor

$$\begin{aligned} \mathfrak{K}^{-1} = & \mathfrak{K}_0^{-1} + [(\sum_{i=0}^1 \mathfrak{K}_i)^{-1} - \mathfrak{K}_0^{-1}]P_1 + [(\sum_{i=0}^2 \mathfrak{K}_i)^{-1} - (\sum_{i=0}^1 \mathfrak{K}_i)^{-1}]P_2 + [(\sum_{i=0}^3 \mathfrak{K}_i)^{-1} - \\ & (\sum_{i=0}^2 \mathfrak{K}_i)^{-1}]P_3 + [(\sum_{i=0}^4 \mathfrak{K}_i)^{-1} - (\sum_{i=0}^3 \mathfrak{K}_i)^{-1}]P_4 + [(\sum_{i=0}^5 \mathfrak{K}_i)^{-1} - (\sum_{i=0}^4 \mathfrak{K}_i)^{-1}]P_5 + [(\sum_{i=0}^6 \mathfrak{K}_i)^{-1} - \\ & (\sum_{i=0}^5 \mathfrak{K}_i)^{-1}]P_6 + [(\sum_{i=0}^7 \mathfrak{K}_i)^{-1} - (\sum_{i=0}^6 \mathfrak{K}_i)^{-1}]P_7 + [(\sum_{i=0}^8 \mathfrak{K}_i)^{-1} - (\sum_{i=0}^7 \mathfrak{K}_i)^{-1}]P_8 + [(\sum_{i=0}^9 \mathfrak{K}_i)^{-1} - \\ & (\sum_{i=0}^8 \mathfrak{K}_i)^{-1}]P_9 + [(\sum_{i=0}^{10} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^9 \mathfrak{K}_i)^{-1}]P_{10} + [(\sum_{i=0}^{11} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{10} \mathfrak{K}_i)^{-1}]P_{11} + [(\sum_{i=0}^{12} \mathfrak{K}_i)^{-1} - \\ & (\sum_{i=0}^{11} \mathfrak{K}_i)^{-1}]P_{12} + [(\sum_{i=0}^{13} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{12} \mathfrak{K}_i)^{-1}]P_{13} + [(\sum_{i=0}^{14} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{13} \mathfrak{K}_i)^{-1}]P_{14} + [(\sum_{i=0}^{15} \mathfrak{K}_i)^{-1} - \\ & (\sum_{i=0}^{14} \mathfrak{K}_i)^{-1}]P_{15} + [(\sum_{i=0}^{16} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{15} \mathfrak{K}_i)^{-1}]P_{16} + [(\sum_{i=0}^{17} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{16} \mathfrak{K}_i)^{-1}]P_{17} + [(\sum_{i=0}^{18} \mathfrak{K}_i)^{-1} - \\ & (\sum_{i=0}^{17} \mathfrak{K}_i)^{-1}]P_{18} + [(\sum_{i=0}^{19} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{18} \mathfrak{K}_i)^{-1}]P_{19} + [(\sum_{i=0}^{20} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{19} \mathfrak{K}_i)^{-1}]P_{20}. \end{aligned}$$

Theorem

Let $\mathfrak{K} = \mathfrak{K}_0 + \sum_{i=1}^{21} \mathfrak{K}_i P_i$ be a 21-plithogenic matrix of size $n \times n$, hence:

1. $\mathfrak{K}$ is invertible if and only if $\det \mathfrak{K}$ is an invertible 21-plithogenic real number.
2. $\mathfrak{K}^{-1} = \mathfrak{K}_0^{-1} + [(\sum_{i=0}^1 \mathfrak{K}_i)^{-1} - \mathfrak{K}_0^{-1}]P_1 + [(\sum_{i=0}^2 \mathfrak{K}_i)^{-1} - (\sum_{i=0}^1 \mathfrak{K}_i)^{-1}]P_2 + [(\sum_{i=0}^3 \mathfrak{K}_i)^{-1} - (\sum_{i=0}^2 \mathfrak{K}_i)^{-1}]P_3 + [(\sum_{i=0}^4 \mathfrak{K}_i)^{-1} - (\sum_{i=0}^3 \mathfrak{K}_i)^{-1}]P_4 + [(\sum_{i=0}^5 \mathfrak{K}_i)^{-1} - (\sum_{i=0}^4 \mathfrak{K}_i)^{-1}]P_5 + [(\sum_{i=0}^6 \mathfrak{K}_i)^{-1} - (\sum_{i=0}^5 \mathfrak{K}_i)^{-1}]P_6 + [(\sum_{i=0}^7 \mathfrak{K}_i)^{-1} - (\sum_{i=0}^6 \mathfrak{K}_i)^{-1}]P_7 + [(\sum_{i=0}^8 \mathfrak{K}_i)^{-1} - (\sum_{i=0}^7 \mathfrak{K}_i)^{-1}]P_8 + [(\sum_{i=0}^9 \mathfrak{K}_i)^{-1} - (\sum_{i=0}^8 \mathfrak{K}_i)^{-1}]P_9 + [(\sum_{i=0}^{10} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^9 \mathfrak{K}_i)^{-1}]P_{10} + [(\sum_{i=0}^{11} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{10} \mathfrak{K}_i)^{-1}]P_{11} + [(\sum_{i=0}^{12} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{11} \mathfrak{K}_i)^{-1}]P_{12} + [(\sum_{i=0}^{13} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{12} \mathfrak{K}_i)^{-1}]P_{13} + [(\sum_{i=0}^{14} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{13} \mathfrak{K}_i)^{-1}]P_{14} + [(\sum_{i=0}^{15} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{14} \mathfrak{K}_i)^{-1}]P_{15} + [(\sum_{i=0}^{16} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{15} \mathfrak{K}_i)^{-1}]P_{16} + [(\sum_{i=0}^{17} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{16} \mathfrak{K}_i)^{-1}]P_{17} + [(\sum_{i=0}^{18} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{17} \mathfrak{K}_i)^{-1}]P_{18} + [(\sum_{i=0}^{19} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{18} \mathfrak{K}_i)^{-1}]P_{19} + [(\sum_{i=0}^{20} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{19} \mathfrak{K}_i)^{-1}]P_{20} + [(\sum_{i=0}^{21} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{20} \mathfrak{K}_i)^{-1}]P_{21}.$

Proof

- 1). Let $\mathfrak{K} = \mathfrak{K}_0 + \sum_{i=1}^{21} \mathfrak{K}_i P_i$, then $\mathfrak{K}$ is invertible if and only if there exists $G = G_0 + \sum_{i=1}^{21} G_i P_i$ such that: $\mathfrak{K} \times G = U_{n \times n}$, hence:

$$\left\{ \begin{array}{l}
 \aleph_0 G_0 = U_{n \times n} \\
 \sum_{i=0}^1 \aleph_i \sum_{i=0}^1 I_i - \aleph_0 G_0 = O_{n \times n} \\
 \sum_{i=0}^2 \aleph_i \sum_{i=0}^2 G_i - \sum_{i=0}^1 \aleph_i \sum_{i=0}^1 G_i = O_{n \times n} \\
 \sum_{i=0}^3 \aleph_i \sum_{i=0}^3 G_i - \sum_{i=0}^2 \aleph_i \sum_{i=0}^2 G_i = O_{n \times n} \\
 \sum_{i=0}^4 \aleph_i \sum_{i=0}^4 G_i - \sum_{i=0}^3 \aleph_i \sum_{i=0}^3 I_i = O_{n \times n} \\
 \sum_{i=0}^5 \aleph_i \sum_{i=0}^5 G_i - \sum_{i=0}^4 \aleph_i \sum_{i=0}^4 I_i = O_{n \times n} \\
 \sum_{i=0}^6 \aleph_i \sum_{i=0}^6 G_i - \sum_{i=0}^5 \aleph_i \sum_{i=0}^5 G_i = O_{n \times n} \\
 \sum_{i=0}^7 \tau_i \sum_{i=0}^7 G_i - \sum_{i=0}^6 \aleph_i \sum_{i=0}^6 G_i = O_{n \times n} \\
 \sum_{i=0}^8 \aleph_i \sum_{i=0}^8 G_i - \sum_{i=0}^7 \aleph_i \sum_{i=0}^7 G_i = O_{n \times n} \\
 \sum_{i=0}^9 \aleph_i \sum_{i=0}^9 G_i - \sum_{i=0}^8 \aleph_i \sum_{i=0}^8 G_i = O_{n \times n} \\
 \sum_{i=0}^{10} \aleph_i \sum_{i=0}^{10} G_i - \sum_{i=0}^9 \aleph_i \sum_{i=0}^9 G_i = O_{n \times n} \\
 \sum_{i=0}^{11} \aleph_i \sum_{i=0}^{11} G_i - \sum_{i=0}^{10} \aleph_i \sum_{i=0}^{10} G_i = O_{n \times n} \\
 \sum_{i=0}^{12} \aleph_i \sum_{i=0}^{12} G_i - \sum_{i=0}^{11} \aleph_i \sum_{i=0}^{11} G_i = O_{n \times n} \\
 \sum_{i=0}^{13} \aleph_i \sum_{i=0}^{13} G_i - \sum_{i=0}^{12} \aleph_i \sum_{i=0}^{12} G_i = O_{n \times n} \\
 \sum_{i=0}^{14} \aleph_i \sum_{i=0}^{14} G_i - \sum_{i=0}^{13} \aleph_i \sum_{i=0}^{13} G_i = O_{n \times n} \\
 \sum_{i=0}^{15} \aleph_i \sum_{i=0}^{15} G_i - \sum_{i=0}^{14} \aleph_i \sum_{i=0}^{14} G_i = O_{n \times n} \\
 \sum_{i=0}^{16} \aleph_i \sum_{i=0}^{16} G_i - \sum_{i=0}^{15} \aleph_i \sum_{i=0}^{15} G_i = O_{n \times n} \\
 \sum_{i=0}^{17} \aleph_i \sum_{i=0}^{17} G_i - \sum_{i=0}^{16} \aleph_i \sum_{i=0}^{16} G_i = O_{n \times n} \\
 \sum_{i=0}^{18} \aleph_i \sum_{i=0}^{18} G_i - \sum_{i=0}^{17} \aleph_i \sum_{i=0}^{17} G_i = O_{n \times n} \\
 \sum_{i=0}^{19} \aleph_i \sum_{i=0}^{19} G_i - \sum_{i=0}^{18} \aleph_i \sum_{i=0}^{18} G_i = O_{n \times n} \\
 \sum_{i=0}^{20} \aleph_i \sum_{i=0}^{20} G_i - \sum_{i=0}^{19} \aleph_i \sum_{i=0}^{19} G_i = O_{n \times n}
 \end{array} \right.$$

And $\sum_{i=0}^{21} \aleph_i \sum_{i=0}^{21} G_i - \sum_{i=0}^{20} \aleph_i \sum_{i=0}^{20} G_i = O_{n \times n}$

$$\left\{ \begin{array}{l} \aleph_0 G_0 = U_{n \times n} \\ \sum_{i=0}^q \aleph_i \sum_{i=0}^q G_i = U_{n \times n} ; 1 \leq q \leq 21 \end{array} \right.$$

Hence $\det(\sum_{i=0}^q \aleph_i) \neq 0$ for all $1 \leq q \leq 21$, so that $\det(\aleph)$ is invertible in $21 - SP_R$.

2). $\sum_{i=0}^q G_i = (\sum_{i=0}^q \aleph_i)^{-1}$ for $1 \leq q \leq 21$, therefor

$$\begin{aligned} \aleph^{-1} &= \aleph_0^{-1} + [(\sum_{i=0}^1 \aleph_i)^{-1} - \aleph_0^{-1}]P_1 + [(\sum_{i=0}^2 \aleph_i)^{-1} - (\sum_{i=0}^1 \aleph_i)^{-1}]P_2 + [(\sum_{i=0}^3 \aleph_i)^{-1} - \\ &(\sum_{i=0}^2 \aleph_i)^{-1}]P_3 + [(\sum_{i=0}^4 \aleph_i)^{-1} - (\sum_{i=0}^3 \aleph_i)^{-1}]P_4 + [(\sum_{i=0}^5 \aleph_i)^{-1} - (\sum_{i=0}^4 \aleph_i)^{-1}]P_5 + [(\sum_{i=0}^6 \aleph_i)^{-1} - \\ &(\sum_{i=0}^5 \aleph_i)^{-1}]P_6 + [(\sum_{i=0}^7 \aleph_i)^{-1} - (\sum_{i=0}^6 \aleph_i)^{-1}]P_7 + [(\sum_{i=0}^8 \aleph_i)^{-1} - (\sum_{i=0}^7 \aleph_i)^{-1}]P_8 + [(\sum_{i=0}^9 \aleph_i)^{-1} - \\ &(\sum_{i=0}^8 \aleph_i)^{-1}]P_9 + [(\sum_{i=0}^{10} \aleph_i)^{-1} - (\sum_{i=0}^9 \aleph_i)^{-1}]P_{10} + [(\sum_{i=0}^{11} \aleph_i)^{-1} - (\sum_{i=0}^{10} \aleph_i)^{-1}]P_{11} + [(\sum_{i=0}^{12} \aleph_i)^{-1} - \\ &(\sum_{i=0}^{11} \aleph_i)^{-1}]P_{12} + [(\sum_{i=0}^{13} \aleph_i)^{-1} - (\sum_{i=0}^{12} \aleph_i)^{-1}]P_{13} + [(\sum_{i=0}^{14} \aleph_i)^{-1} - (\sum_{i=0}^{13} \aleph_i)^{-1}]P_{14} + [(\sum_{i=0}^{15} \aleph_i)^{-1} - \\ &(\sum_{i=0}^{14} \aleph_i)^{-1}]P_{15} + [(\sum_{i=0}^{16} \aleph_i)^{-1} - (\sum_{i=0}^{15} \aleph_i)^{-1}]P_{16} + [(\sum_{i=0}^{17} \aleph_i)^{-1} - (\sum_{i=0}^{16} \aleph_i)^{-1}]P_{17} + [(\sum_{i=0}^{18} \aleph_i)^{-1} - \\ &(\sum_{i=0}^{17} \aleph_i)^{-1}]P_{18} + [(\sum_{i=0}^{19} \aleph_i)^{-1} - (\sum_{i=0}^{18} \aleph_i)^{-1}]P_{19} + [(\sum_{i=0}^{20} \aleph_i)^{-1} - (\sum_{i=0}^{19} \aleph_i)^{-1}]P_{20} + [(\sum_{i=0}^{21} \aleph_i)^{-1} - \\ &(\sum_{i=0}^{20} \aleph_i)^{-1}]P_{21}. \end{aligned}$$

Theorem

Let $\aleph = \aleph_0 + \sum_{i=1}^{22} \aleph_i P_i$ be a 22-plithogenic matrix of size $n \times n$, hence:

1. $\aleph$ is invertible if and only if $\det \aleph$ is an invertible 22-plithogenic real number.
2. $\aleph^{-1} = \aleph_0^{-1} + [(\sum_{i=0}^1 \aleph_i)^{-1} - \aleph_0^{-1}]P_1 + [(\sum_{i=0}^2 \aleph_i)^{-1} - (\sum_{i=0}^1 \aleph_i)^{-1}]P_2 + [(\sum_{i=0}^3 \aleph_i)^{-1} - \\ (\sum_{i=0}^2 \aleph_i)^{-1}]P_3 + [(\sum_{i=0}^4 \aleph_i)^{-1} - (\sum_{i=0}^3 \aleph_i)^{-1}]P_4 + [(\sum_{i=0}^5 \aleph_i)^{-1} - (\sum_{i=0}^4 \aleph_i)^{-1}]P_5 + [(\sum_{i=0}^6 \aleph_i)^{-1} - \\ (\sum_{i=0}^5 \aleph_i)^{-1}]P_6 + [(\sum_{i=0}^7 \aleph_i)^{-1} - (\sum_{i=0}^6 \aleph_i)^{-1}]P_7 + [(\sum_{i=0}^8 \aleph_i)^{-1} - (\sum_{i=0}^7 \aleph_i)^{-1}]P_8 + [(\sum_{i=0}^9 \aleph_i)^{-1} - \\ (\sum_{i=0}^8 \aleph_i)^{-1}]P_9 + [(\sum_{i=0}^{10} \aleph_i)^{-1} - (\sum_{i=0}^9 \aleph_i)^{-1}]P_{10} + [(\sum_{i=0}^{11} \aleph_i)^{-1} - (\sum_{i=0}^{10} \aleph_i)^{-1}]P_{11} + [(\sum_{i=0}^{12} \aleph_i)^{-1} - \\ (\sum_{i=0}^{11} \aleph_i)^{-1}]P_{12} + [(\sum_{i=0}^{13} \aleph_i)^{-1} - (\sum_{i=0}^{12} \aleph_i)^{-1}]P_{13} + [(\sum_{i=0}^{14} \aleph_i)^{-1} - (\sum_{i=0}^{13} \aleph_i)^{-1}]P_{14} + [(\sum_{i=0}^{15} \aleph_i)^{-1} - \\ (\sum_{i=0}^{14} \aleph_i)^{-1}]P_{15} + [(\sum_{i=0}^{16} \aleph_i)^{-1} - (\sum_{i=0}^{15} \aleph_i)^{-1}]P_{16} + [(\sum_{i=0}^{17} \aleph_i)^{-1} - (\sum_{i=0}^{16} \aleph_i)^{-1}]P_{17} + \\ [(\sum_{i=0}^{18} \aleph_i)^{-1} - (\sum_{i=0}^{17} \aleph_i)^{-1}]P_{18} + [(\sum_{i=0}^{19} \aleph_i)^{-1} - (\sum_{i=0}^{18} \aleph_i)^{-1}]P_{19} + [(\sum_{i=0}^{20} \aleph_i)^{-1} - (\sum_{i=0}^{19} \aleph_i)^{-1}]P_{20} + \\ [(\sum_{i=0}^{21} \aleph_i)^{-1} - (\sum_{i=0}^{20} \aleph_i)^{-1}]P_{21} + [(\sum_{i=0}^{22} \aleph_i)^{-1} - (\sum_{i=0}^{21} \aleph_i)^{-1}]P_{22}.$

Proof

- 1). Let $\aleph = \aleph_0 + \sum_{i=1}^{22} \aleph_i P_i$, then $\aleph$ is invertible if and only if there exists $G = G_0 + \sum_{i=1}^{22} G_i P_i$ such that:
 $\aleph \times G = U_{n \times n}$, hence:

$$\left\{ \begin{array}{l}
 \aleph_0 G_0 = U_{n \times n} \\
 \sum_{i=0}^1 \aleph_i \sum_{i=0}^1 I_i - \aleph_0 G_0 = O_{n \times n} \\
 \sum_{i=0}^2 \aleph_i \sum_{i=0}^2 G_i - \sum_{i=0}^1 \aleph_i \sum_{i=0}^1 G_i = O_{n \times n} \\
 \sum_{i=0}^3 \aleph_i \sum_{i=0}^3 G_i - \sum_{i=0}^2 \aleph_i \sum_{i=0}^2 G_i = O_{n \times n} \\
 \sum_{i=0}^4 \aleph_i \sum_{i=0}^4 G_i - \sum_{i=0}^3 \aleph_i \sum_{i=0}^3 I_i = O_{n \times n} \\
 \sum_{i=0}^5 \aleph_i \sum_{i=0}^5 G_i - \sum_{i=0}^4 \aleph_i \sum_{i=0}^4 I_i = O_{n \times n} \\
 \sum_{i=0}^6 \aleph_i \sum_{i=0}^6 G_i - \sum_{i=0}^5 \aleph_i \sum_{i=0}^5 G_i = O_{n \times n} \\
 \sum_{i=0}^7 \tau_i \sum_{i=0}^7 G_i - \sum_{i=0}^6 \aleph_i \sum_{i=0}^6 G_i = O_{n \times n} \\
 \sum_{i=0}^8 \aleph_i \sum_{i=0}^8 G_i - \sum_{i=0}^7 \aleph_i \sum_{i=0}^7 G_i = O_{n \times n} \\
 \sum_{i=0}^9 \aleph_i \sum_{i=0}^9 G_i - \sum_{i=0}^8 \aleph_i \sum_{i=0}^8 G_i = O_{n \times n} \\
 \sum_{i=0}^{10} \aleph_i \sum_{i=0}^{10} G_i - \sum_{i=0}^9 \aleph_i \sum_{i=0}^9 G_i = O_{n \times n} \\
 \sum_{i=0}^{11} \aleph_i \sum_{i=0}^{11} G_i - \sum_{i=0}^{10} \aleph_i \sum_{i=0}^{10} G_i = O_{n \times n} \\
 \sum_{i=0}^{12} \aleph_i \sum_{i=0}^{12} G_i - \sum_{i=0}^{11} \aleph_i \sum_{i=0}^{11} G_i = O_{n \times n} \\
 \sum_{i=0}^{13} \aleph_i \sum_{i=0}^{13} G_i - \sum_{i=0}^{12} \aleph_i \sum_{i=0}^{12} G_i = O_{n \times n} \\
 \sum_{i=0}^{14} \aleph_i \sum_{i=0}^{14} G_i - \sum_{i=0}^{13} \aleph_i \sum_{i=0}^{13} G_i = O_{n \times n} \\
 \sum_{i=0}^{15} \aleph_i \sum_{i=0}^{15} G_i - \sum_{i=0}^{14} \aleph_i \sum_{i=0}^{14} G_i = O_{n \times n} \\
 \sum_{i=0}^{16} \aleph_i \sum_{i=0}^{16} G_i - \sum_{i=0}^{15} \aleph_i \sum_{i=0}^{15} G_i = O_{n \times n} \\
 \sum_{i=0}^{17} \aleph_i \sum_{i=0}^{17} G_i - \sum_{i=0}^{16} \aleph_i \sum_{i=0}^{16} G_i = O_{n \times n} \\
 \sum_{i=0}^{18} \aleph_i \sum_{i=0}^{18} G_i - \sum_{i=0}^{17} \aleph_i \sum_{i=0}^{17} G_i = O_{n \times n} \\
 \sum_{i=0}^{19} \aleph_i \sum_{i=0}^{19} G_i - \sum_{i=0}^{18} \aleph_i \sum_{i=0}^{18} G_i = O_{n \times n} \\
 \sum_{i=0}^{20} \aleph_i \sum_{i=0}^{20} G_i - \sum_{i=0}^{19} \aleph_i \sum_{i=0}^{19} G_i = O_{n \times n}
 \end{array} \right.$$

And $\sum_{i=0}^{21} \aleph_i \sum_{i=0}^{21} G_i - \sum_{i=0}^{20} \aleph_i \sum_{i=0}^{20} G_i = O_{n \times n}$, $\sum_{i=0}^{22} \aleph_i \sum_{i=0}^{22} G_i - \sum_{i=0}^{21} \aleph_i \sum_{i=0}^{21} G_i = O_{n \times n}$

$$\begin{cases} \aleph_0 G_0 = U_{n \times n} \\ \sum_{i=0}^q \aleph_i \sum_{i=0}^q G_i = U_{n \times n} ; 1 \leq q \leq 22 \end{cases}$$

Hence $\det(\sum_{i=0}^q \aleph_i) \neq 0$ for all $1 \leq q \leq 22$, so that $\det(\aleph)$ is invertible in $22 - SP_R$.

2). $\sum_{i=0}^q G_i = (\sum_{i=0}^q \aleph_i)^{-1}$ for $1 \leq q \leq 22$, therefor

$$\begin{aligned} \aleph^{-1} &= \aleph_0^{-1} + [(\sum_{i=0}^1 \aleph_i)^{-1} - \aleph_0^{-1}]P_1 + [(\sum_{i=0}^2 \aleph_i)^{-1} - (\sum_{i=0}^1 \aleph_i)^{-1}]P_2 + [(\sum_{i=0}^3 \aleph_i)^{-1} - \\ &(\sum_{i=0}^2 \aleph_i)^{-1}]P_3 + [(\sum_{i=0}^4 \aleph_i)^{-1} - (\sum_{i=0}^3 \aleph_i)^{-1}]P_4 + [(\sum_{i=0}^5 \aleph_i)^{-1} - (\sum_{i=0}^4 \aleph_i)^{-1}]P_5 + [(\sum_{i=0}^6 \aleph_i)^{-1} - \\ &(\sum_{i=0}^5 \aleph_i)^{-1}]P_6 + [(\sum_{i=0}^7 \aleph_i)^{-1} - (\sum_{i=0}^6 \aleph_i)^{-1}]P_7 + [(\sum_{i=0}^8 \aleph_i)^{-1} - (\sum_{i=0}^7 \aleph_i)^{-1}]P_8 + [(\sum_{i=0}^9 \aleph_i)^{-1} - \\ &(\sum_{i=0}^8 \aleph_i)^{-1}]P_9 + [(\sum_{i=0}^{10} \aleph_i)^{-1} - (\sum_{i=0}^9 \aleph_i)^{-1}]P_{10} + [(\sum_{i=0}^{11} \aleph_i)^{-1} - (\sum_{i=0}^{10} \aleph_i)^{-1}]P_{11} + [(\sum_{i=0}^{12} \aleph_i)^{-1} - \\ &(\sum_{i=0}^{11} \aleph_i)^{-1}]P_{12} + [(\sum_{i=0}^{13} \aleph_i)^{-1} - (\sum_{i=0}^{12} \aleph_i)^{-1}]P_{13} + [(\sum_{i=0}^{14} \aleph_i)^{-1} - (\sum_{i=0}^{13} \aleph_i)^{-1}]P_{14} + [(\sum_{i=0}^{15} \aleph_i)^{-1} - \\ &(\sum_{i=0}^{14} \aleph_i)^{-1}]P_{15} + [(\sum_{i=0}^{16} \aleph_i)^{-1} - (\sum_{i=0}^{15} \aleph_i)^{-1}]P_{16} + [(\sum_{i=0}^{17} \aleph_i)^{-1} - (\sum_{i=0}^{16} \aleph_i)^{-1}]P_{17} + [(\sum_{i=0}^{18} \aleph_i)^{-1} - \\ &(\sum_{i=0}^{17} \aleph_i)^{-1}]P_{18} + [(\sum_{i=0}^{19} \aleph_i)^{-1} - (\sum_{i=0}^{18} \aleph_i)^{-1}]P_{19} + [(\sum_{i=0}^{20} \aleph_i)^{-1} - (\sum_{i=0}^{19} \aleph_i)^{-1}]P_{20} + [(\sum_{i=0}^{21} \aleph_i)^{-1} - \\ &(\sum_{i=0}^{20} \aleph_i)^{-1}]P_{21} + [(\sum_{i=0}^{22} \aleph_i)^{-1} - (\sum_{i=0}^{21} \aleph_i)^{-1}]P_{22}. \end{aligned}$$

Theorem

Let $\aleph = \aleph_0 + \sum_{i=1}^{23} \aleph_i P_i$ be a 23-plithogenic matrix of size $n \times n$, hence:

1. $\aleph$ is invertible if and only if $\det \aleph$ is an invertible 23-plithogenic real number.
2. $\aleph^{-1} = \aleph_0^{-1} + [(\sum_{i=0}^1 \aleph_i)^{-1} - \aleph_0^{-1}]P_1 + [(\sum_{i=0}^2 \aleph_i)^{-1} - (\sum_{i=0}^1 \aleph_i)^{-1}]P_2 + [(\sum_{i=0}^3 \aleph_i)^{-1} - (\sum_{i=0}^2 \aleph_i)^{-1}]P_3 + [(\sum_{i=0}^4 \aleph_i)^{-1} - (\sum_{i=0}^3 \aleph_i)^{-1}]P_4 + [(\sum_{i=0}^5 \aleph_i)^{-1} - (\sum_{i=0}^4 \aleph_i)^{-1}]P_5 + [(\sum_{i=0}^6 \aleph_i)^{-1} - (\sum_{i=0}^5 \aleph_i)^{-1}]P_6 + [(\sum_{i=0}^7 \aleph_i)^{-1} - (\sum_{i=0}^6 \aleph_i)^{-1}]P_7 + [(\sum_{i=0}^8 \aleph_i)^{-1} - (\sum_{i=0}^7 \aleph_i)^{-1}]P_8 + [(\sum_{i=0}^9 \aleph_i)^{-1} - (\sum_{i=0}^8 \aleph_i)^{-1}]P_9 + [(\sum_{i=0}^{10} \aleph_i)^{-1} - (\sum_{i=0}^9 \aleph_i)^{-1}]P_{10} + [(\sum_{i=0}^{11} \aleph_i)^{-1} - (\sum_{i=0}^{10} \aleph_i)^{-1}]P_{11} + [(\sum_{i=0}^{12} \aleph_i)^{-1} - (\sum_{i=0}^{11} \aleph_i)^{-1}]P_{12} + [(\sum_{i=0}^{13} \aleph_i)^{-1} - (\sum_{i=0}^{12} \aleph_i)^{-1}]P_{13} + [(\sum_{i=0}^{14} \aleph_i)^{-1} - (\sum_{i=0}^{13} \aleph_i)^{-1}]P_{14} + [(\sum_{i=0}^{15} \aleph_i)^{-1} - (\sum_{i=0}^{14} \aleph_i)^{-1}]P_{15} + [(\sum_{i=0}^{16} \aleph_i)^{-1} - (\sum_{i=0}^{15} \aleph_i)^{-1}]P_{16} + [(\sum_{i=0}^{17} \aleph_i)^{-1} - (\sum_{i=0}^{16} \aleph_i)^{-1}]P_{17} + [(\sum_{i=0}^{18} \aleph_i)^{-1} - (\sum_{i=0}^{17} \aleph_i)^{-1}]P_{18} + [(\sum_{i=0}^{19} \aleph_i)^{-1} - (\sum_{i=0}^{18} \aleph_i)^{-1}]P_{19} + [(\sum_{i=0}^{20} \aleph_i)^{-1} - (\sum_{i=0}^{19} \aleph_i)^{-1}]P_{20} + [(\sum_{i=0}^{21} \aleph_i)^{-1} - (\sum_{i=0}^{20} \aleph_i)^{-1}]P_{21} + [(\sum_{i=0}^{22} \aleph_i)^{-1} - (\sum_{i=0}^{21} \aleph_i)^{-1}]P_{22} + [(\sum_{i=0}^{23} \aleph_i)^{-1} - (\sum_{i=0}^{22} \aleph_i)^{-1}]P_{23}.$

Proof

- 1). Let $\aleph = \aleph_0 + \sum_{i=1}^{23} \aleph_i P_i$, then $\aleph$ is invertible if and only if there exists $G = G_0 + \sum_{i=1}^{23} G_i P_i$ such that: $\aleph \times G = U_{n \times n}$, hence:

$$\left\{ \begin{array}{l}
\aleph_0 G_0 = U_{n \times n} \\
\sum_{i=0}^1 \aleph_i \sum_{i=0}^1 I_i - \aleph_0 G_0 = O_{n \times n} \\
\sum_{i=0}^2 \aleph_i \sum_{i=0}^2 G_i - \sum_{i=0}^1 \aleph_i \sum_{i=0}^1 G_i = O_{n \times n} \\
\sum_{i=0}^3 \aleph_i \sum_{i=0}^3 G_i - \sum_{i=0}^2 \aleph_i \sum_{i=0}^2 G_i = O_{n \times n} \\
\sum_{i=0}^4 \aleph_i \sum_{i=0}^4 G_i - \sum_{i=0}^3 \aleph_i \sum_{i=0}^3 I_i = O_{n \times n} \\
\sum_{i=0}^5 \aleph_i \sum_{i=0}^5 G_i - \sum_{i=0}^4 \aleph_i \sum_{i=0}^4 I_i = O_{n \times n} \\
\sum_{i=0}^6 \aleph_i \sum_{i=0}^6 G_i - \sum_{i=0}^5 \aleph_i \sum_{i=0}^5 G_i = O_{n \times n} \\
\sum_{i=0}^7 \tau_i \sum_{i=0}^7 G_i - \sum_{i=0}^6 \aleph_i \sum_{i=0}^6 G_i = O_{n \times n} \\
\sum_{i=0}^8 \aleph_i \sum_{i=0}^8 G_i - \sum_{i=0}^7 \aleph_i \sum_{i=0}^7 G_i = O_{n \times n} \\
\sum_{i=0}^9 \aleph_i \sum_{i=0}^9 G_i - \sum_{i=0}^8 \aleph_i \sum_{i=0}^8 G_i = O_{n \times n} \\
\sum_{i=0}^{10} \aleph_i \sum_{i=0}^{10} G_i - \sum_{i=0}^9 \aleph_i \sum_{i=0}^9 G_i = O_{n \times n} \\
\sum_{i=0}^{11} \aleph_i \sum_{i=0}^{11} G_i - \sum_{i=0}^{10} \aleph_i \sum_{i=0}^{10} G_i = O_{n \times n} \\
\sum_{i=0}^{12} \aleph_i \sum_{i=0}^{12} G_i - \sum_{i=0}^{11} \aleph_i \sum_{i=0}^{11} G_i = O_{n \times n} \\
\sum_{i=0}^{13} \aleph_i \sum_{i=0}^{13} G_i - \sum_{i=0}^{12} \aleph_i \sum_{i=0}^{12} G_i = O_{n \times n} \\
\sum_{i=0}^{14} \aleph_i \sum_{i=0}^{14} G_i - \sum_{i=0}^{13} \aleph_i \sum_{i=0}^{13} G_i = O_{n \times n} \\
\sum_{i=0}^{15} \aleph_i \sum_{i=0}^{15} G_i - \sum_{i=0}^{14} \aleph_i \sum_{i=0}^{14} G_i = O_{n \times n} \\
\sum_{i=0}^{16} \aleph_i \sum_{i=0}^{16} G_i - \sum_{i=0}^{15} \aleph_i \sum_{i=0}^{15} G_i = O_{n \times n} \\
\sum_{i=0}^{17} \aleph_i \sum_{i=0}^{17} G_i - \sum_{i=0}^{16} \aleph_i \sum_{i=0}^{16} G_i = O_{n \times n} \\
\sum_{i=0}^{18} \aleph_i \sum_{i=0}^{18} G_i - \sum_{i=0}^{17} \aleph_i \sum_{i=0}^{17} G_i = O_{n \times n} \\
\sum_{i=0}^{19} \aleph_i \sum_{i=0}^{19} G_i - \sum_{i=0}^{18} \aleph_i \sum_{i=0}^{18} G_i = O_{n \times n} \\
\sum_{i=0}^{20} \aleph_i \sum_{i=0}^{20} G_i - \sum_{i=0}^{19} \aleph_i \sum_{i=0}^{19} G_i = O_{n \times n}
\end{array} \right.$$

And $\sum_{i=0}^{21} \aleph_i \sum_{i=0}^{21} G_i - \sum_{i=0}^{20} \aleph_i \sum_{i=0}^{20} G_i = O_{n \times n}$, $\sum_{i=0}^{22} \aleph_i \sum_{i=0}^{22} G_i - \sum_{i=0}^{21} \aleph_i \sum_{i=0}^{21} G_i = O_{n \times n}$, $\sum_{i=0}^{23} \aleph_i \sum_{i=0}^{23} G_i - \sum_{i=0}^{22} \aleph_i \sum_{i=0}^{22} G_i = O_{n \times n}$

$$\left\{ \begin{array}{l} \aleph_0 G_0 = U_{n \times n} \\ \sum_{i=0}^q \aleph_i \sum_{i=0}^q G_i = U_{n \times n} ; 1 \leq q \leq 23 \end{array} \right.$$

Hence $\det(\sum_{i=0}^q \aleph_i) \neq 0$ for all $1 \leq q \leq 23$, so that $\det(\aleph)$ is invertible in $23 - SP_R$.

2). $\sum_{i=0}^q G_i = (\sum_{i=0}^q \aleph_i)^{-1}$ for $1 \leq q \leq 23$, therefor

$$\begin{aligned} \aleph^{-1} = & \aleph_0^{-1} + [(\sum_{i=0}^1 \aleph_i)^{-1} - \aleph_0^{-1}]P_1 + [(\sum_{i=0}^2 \aleph_i)^{-1} - (\sum_{i=0}^1 \aleph_i)^{-1}]P_2 + [(\sum_{i=0}^3 \aleph_i)^{-1} - \\ & (\sum_{i=0}^2 \aleph_i)^{-1}]P_3 + [(\sum_{i=0}^4 \aleph_i)^{-1} - (\sum_{i=0}^3 \aleph_i)^{-1}]P_4 + [(\sum_{i=0}^5 \aleph_i)^{-1} - (\sum_{i=0}^4 \aleph_i)^{-1}]P_5 + [(\sum_{i=0}^6 \aleph_i)^{-1} - \\ & (\sum_{i=0}^5 \aleph_i)^{-1}]P_6 + [(\sum_{i=0}^7 \aleph_i)^{-1} - (\sum_{i=0}^6 \aleph_i)^{-1}]P_7 + [(\sum_{i=0}^8 \aleph_i)^{-1} - (\sum_{i=0}^7 \aleph_i)^{-1}]P_8 + [(\sum_{i=0}^9 \aleph_i)^{-1} - \\ & (\sum_{i=0}^8 \aleph_i)^{-1}]P_9 + [(\sum_{i=0}^{10} \aleph_i)^{-1} - (\sum_{i=0}^9 \aleph_i)^{-1}]P_{10} + [(\sum_{i=0}^{11} \aleph_i)^{-1} - (\sum_{i=0}^{10} \aleph_i)^{-1}]P_{11} + [(\sum_{i=0}^{12} \aleph_i)^{-1} - \\ & (\sum_{i=0}^{11} \aleph_i)^{-1}]P_{12} + [(\sum_{i=0}^{13} \aleph_i)^{-1} - (\sum_{i=0}^{12} \aleph_i)^{-1}]P_{13} + [(\sum_{i=0}^{14} \aleph_i)^{-1} - (\sum_{i=0}^{13} \aleph_i)^{-1}]P_{14} + [(\sum_{i=0}^{15} \aleph_i)^{-1} - \\ & (\sum_{i=0}^{14} \aleph_i)^{-1}]P_{15} + [(\sum_{i=0}^{16} \aleph_i)^{-1} - (\sum_{i=0}^{15} \aleph_i)^{-1}]P_{16} + [(\sum_{i=0}^{17} \aleph_i)^{-1} - (\sum_{i=0}^{16} \aleph_i)^{-1}]P_{17} + [(\sum_{i=0}^{18} \aleph_i)^{-1} - \\ & (\sum_{i=0}^{17} \aleph_i)^{-1}]P_{18} + [(\sum_{i=0}^{19} \aleph_i)^{-1} - (\sum_{i=0}^{18} \aleph_i)^{-1}]P_{19} + [(\sum_{i=0}^{20} \aleph_i)^{-1} - (\sum_{i=0}^{19} \aleph_i)^{-1}]P_{20} + [(\sum_{i=0}^{21} \aleph_i)^{-1} - \\ & (\sum_{i=0}^{20} \aleph_i)^{-1}]P_{21} + [(\sum_{i=0}^{22} \aleph_i)^{-1} - (\sum_{i=0}^{21} \aleph_i)^{-1}]P_{22} + [(\sum_{i=0}^{23} \aleph_i)^{-1} - (\sum_{i=0}^{22} \aleph_i)^{-1}]P_{23}. \end{aligned}$$

Theorem

Let $\aleph = \aleph_0 + \sum_{i=1}^{24} \aleph_i P_i$ be a 24-plithogenic matrix of size $n \times n$, hence:

1. $\aleph$ is invertible if and only if $\det \aleph$ is an invertible 24-plithogenic real number.
2. $\aleph^{-1} = \aleph_0^{-1} + [(\sum_{i=0}^1 \aleph_i)^{-1} - \aleph_0^{-1}]P_1 + [(\sum_{i=0}^2 \aleph_i)^{-1} - (\sum_{i=0}^1 \aleph_i)^{-1}]P_2 + [(\sum_{i=0}^3 \aleph_i)^{-1} - (\sum_{i=0}^2 \aleph_i)^{-1}]P_3 + [(\sum_{i=0}^4 \aleph_i)^{-1} - (\sum_{i=0}^3 \aleph_i)^{-1}]P_4 + [(\sum_{i=0}^5 \aleph_i)^{-1} - (\sum_{i=0}^4 \aleph_i)^{-1}]P_5 + [(\sum_{i=0}^6 \aleph_i)^{-1} - (\sum_{i=0}^5 \aleph_i)^{-1}]P_6 + [(\sum_{i=0}^7 \aleph_i)^{-1} - (\sum_{i=0}^6 \aleph_i)^{-1}]P_7 + [(\sum_{i=0}^8 \aleph_i)^{-1} - (\sum_{i=0}^7 \aleph_i)^{-1}]P_8 + [(\sum_{i=0}^9 \aleph_i)^{-1} - (\sum_{i=0}^8 \aleph_i)^{-1}]P_9 + [(\sum_{i=0}^{10} \aleph_i)^{-1} - (\sum_{i=0}^9 \aleph_i)^{-1}]P_{10} + [(\sum_{i=0}^{11} \aleph_i)^{-1} - (\sum_{i=0}^{10} \aleph_i)^{-1}]P_{11} + [(\sum_{i=0}^{12} \aleph_i)^{-1} - (\sum_{i=0}^{11} \aleph_i)^{-1}]P_{12} + [(\sum_{i=0}^{13} \aleph_i)^{-1} - (\sum_{i=0}^{12} \aleph_i)^{-1}]P_{13} + [(\sum_{i=0}^{14} \aleph_i)^{-1} - (\sum_{i=0}^{13} \aleph_i)^{-1}]P_{14} + [(\sum_{i=0}^{15} \aleph_i)^{-1} - (\sum_{i=0}^{14} \aleph_i)^{-1}]P_{15} + [(\sum_{i=0}^{16} \aleph_i)^{-1} - (\sum_{i=0}^{15} \aleph_i)^{-1}]P_{16} + [(\sum_{i=0}^{17} \aleph_i)^{-1} - (\sum_{i=0}^{16} \aleph_i)^{-1}]P_{17} + [(\sum_{i=0}^{18} \aleph_i)^{-1} - (\sum_{i=0}^{17} \aleph_i)^{-1}]P_{18} + [(\sum_{i=0}^{19} \aleph_i)^{-1} - (\sum_{i=0}^{18} \aleph_i)^{-1}]P_{19} + [(\sum_{i=0}^{20} \aleph_i)^{-1} - (\sum_{i=0}^{19} \aleph_i)^{-1}]P_{20} + [(\sum_{i=0}^{21} \aleph_i)^{-1} - (\sum_{i=0}^{20} \aleph_i)^{-1}]P_{21} + [(\sum_{i=0}^{22} \aleph_i)^{-1} - (\sum_{i=0}^{21} \aleph_i)^{-1}]P_{22} + [(\sum_{i=0}^{23} \aleph_i)^{-1} - (\sum_{i=0}^{22} \aleph_i)^{-1}]P_{23} + [(\sum_{i=0}^{24} \aleph_i)^{-1} - (\sum_{i=0}^{23} \aleph_i)^{-1}]P_{24}.$

Proof

- 1). Let $\aleph = \aleph_0 + \sum_{i=1}^{24} \aleph_i P_i$, then $\aleph$ is invertible if and only if there exists $G = G_0 + \sum_{i=1}^{24} G_i P_i$ such that: $\aleph \times G = U_{n \times n}$, hence:

$$\left\{ \begin{array}{l}
\aleph_0 G_0 = U_{n \times n} \\
\sum_{i=0}^1 \aleph_i \sum_{i=0}^1 I_i - \aleph_0 G_0 = O_{n \times n} \\
\sum_{i=0}^2 \aleph_i \sum_{i=0}^2 G_i - \sum_{i=0}^1 \aleph_i \sum_{i=0}^1 G_i = O_{n \times n} \\
\sum_{i=0}^3 \aleph_i \sum_{i=0}^3 G_i - \sum_{i=0}^2 \aleph_i \sum_{i=0}^2 G_i = O_{n \times n} \\
\sum_{i=0}^4 \aleph_i \sum_{i=0}^4 G_i - \sum_{i=0}^3 \aleph_i \sum_{i=0}^3 I_i = O_{n \times n} \\
\sum_{i=0}^5 \aleph_i \sum_{i=0}^5 G_i - \sum_{i=0}^4 \aleph_i \sum_{i=0}^4 I_i = O_{n \times n} \\
\sum_{i=0}^6 \aleph_i \sum_{i=0}^6 G_i - \sum_{i=0}^5 \aleph_i \sum_{i=0}^5 G_i = O_{n \times n} \\
\sum_{i=0}^7 \tau_i \sum_{i=0}^7 G_i - \sum_{i=0}^6 \aleph_i \sum_{i=0}^6 G_i = O_{n \times n} \\
\sum_{i=0}^8 \aleph_i \sum_{i=0}^8 G_i - \sum_{i=0}^7 \aleph_i \sum_{i=0}^7 G_i = O_{n \times n} \\
\sum_{i=0}^9 \aleph_i \sum_{i=0}^9 G_i - \sum_{i=0}^8 \aleph_i \sum_{i=0}^8 G_i = O_{n \times n} \\
\sum_{i=0}^{10} \aleph_i \sum_{i=0}^{10} G_i - \sum_{i=0}^9 \aleph_i \sum_{i=0}^9 G_i = O_{n \times n} \\
\sum_{i=0}^{11} \aleph_i \sum_{i=0}^{11} G_i - \sum_{i=0}^{10} \aleph_i \sum_{i=0}^{10} G_i = O_{n \times n} \\
\sum_{i=0}^{12} \aleph_i \sum_{i=0}^{12} G_i - \sum_{i=0}^{11} \aleph_i \sum_{i=0}^{11} G_i = O_{n \times n} \\
\sum_{i=0}^{13} \aleph_i \sum_{i=0}^{13} G_i - \sum_{i=0}^{12} \aleph_i \sum_{i=0}^{12} G_i = O_{n \times n} \\
\sum_{i=0}^{14} \aleph_i \sum_{i=0}^{14} G_i - \sum_{i=0}^{13} \aleph_i \sum_{i=0}^{13} G_i = O_{n \times n} \\
\sum_{i=0}^{15} \aleph_i \sum_{i=0}^{15} G_i - \sum_{i=0}^{14} \aleph_i \sum_{i=0}^{14} G_i = O_{n \times n} \\
\sum_{i=0}^{16} \aleph_i \sum_{i=0}^{16} G_i - \sum_{i=0}^{15} \aleph_i \sum_{i=0}^{15} G_i = O_{n \times n} \\
\sum_{i=0}^{17} \aleph_i \sum_{i=0}^{17} G_i - \sum_{i=0}^{16} \aleph_i \sum_{i=0}^{16} G_i = O_{n \times n} \\
\sum_{i=0}^{18} \aleph_i \sum_{i=0}^{18} G_i - \sum_{i=0}^{17} \aleph_i \sum_{i=0}^{17} G_i = O_{n \times n} \\
\sum_{i=0}^{19} \aleph_i \sum_{i=0}^{19} G_i - \sum_{i=0}^{18} \aleph_i \sum_{i=0}^{18} G_i = O_{n \times n} \\
\sum_{i=0}^{20} \aleph_i \sum_{i=0}^{20} G_i - \sum_{i=0}^{19} \aleph_i \sum_{i=0}^{19} G_i = O_{n \times n}
\end{array} \right.$$

And $\sum_{i=0}^{21} \mathfrak{K}_i \sum_{i=0}^{21} G_i - \sum_{i=0}^{20} \mathfrak{K}_i \sum_{i=0}^{20} G_i = O_{n \times n}$, $\sum_{i=0}^{22} \mathfrak{K}_i \sum_{i=0}^{22} G_i - \sum_{i=0}^{21} \mathfrak{K}_i \sum_{i=0}^{21} G_i = O_{n \times n}$, $\sum_{i=0}^{23} \mathfrak{K}_i \sum_{i=0}^{23} G_i - \sum_{i=0}^{22} \mathfrak{K}_i \sum_{i=0}^{22} G_i = O_{n \times n}$, $\sum_{i=0}^{24} \mathfrak{K}_i \sum_{i=0}^{24} G_i - \sum_{i=0}^{23} \mathfrak{K}_i \sum_{i=0}^{23} G_i = O_{n \times n}$

$$\left\{ \begin{array}{l} \mathfrak{K}_0 G_0 = U_{n \times n} \\ \sum_{i=0}^q \mathfrak{K}_i \sum_{i=0}^q G_i = U_{n \times n} ; 1 \leq q \leq 24 \end{array} \right.$$

Hence $\det(\sum_{i=0}^q \mathfrak{K}_i) \neq 0$ for all $1 \leq q \leq 24$, so that $\det(\mathfrak{K})$ is invertible in $24 - SP_R$.

2). $\sum_{i=0}^q G_i = (\sum_{i=0}^q \mathfrak{K}_i)^{-1}$ for $1 \leq q \leq 24$, therefor

$$\begin{aligned} \mathfrak{K}^{-1} = & \mathfrak{K}_0^{-1} + [(\sum_{i=0}^1 \mathfrak{K}_i)^{-1} - \mathfrak{K}_0^{-1}]P_1 + [(\sum_{i=0}^2 \mathfrak{K}_i)^{-1} - (\sum_{i=0}^1 \mathfrak{K}_i)^{-1}]P_2 + [(\sum_{i=0}^3 \mathfrak{K}_i)^{-1} - \\ & (\sum_{i=0}^2 \mathfrak{K}_i)^{-1}]P_3 + [(\sum_{i=0}^4 \mathfrak{K}_i)^{-1} - (\sum_{i=0}^3 \mathfrak{K}_i)^{-1}]P_4 + [(\sum_{i=0}^5 \mathfrak{K}_i)^{-1} - (\sum_{i=0}^4 \mathfrak{K}_i)^{-1}]P_5 + [(\sum_{i=0}^6 \mathfrak{K}_i)^{-1} - \\ & (\sum_{i=0}^5 \mathfrak{K}_i)^{-1}]P_6 + [(\sum_{i=0}^7 \mathfrak{K}_i)^{-1} - (\sum_{i=0}^6 \mathfrak{K}_i)^{-1}]P_7 + [(\sum_{i=0}^8 \mathfrak{K}_i)^{-1} - (\sum_{i=0}^7 \mathfrak{K}_i)^{-1}]P_8 + [(\sum_{i=0}^9 \mathfrak{K}_i)^{-1} - \\ & (\sum_{i=0}^8 \mathfrak{K}_i)^{-1}]P_9 + [(\sum_{i=0}^{10} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^9 \mathfrak{K}_i)^{-1}]P_{10} + [(\sum_{i=0}^{11} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{10} \mathfrak{K}_i)^{-1}]P_{11} + [(\sum_{i=0}^{12} \mathfrak{K}_i)^{-1} - \\ & (\sum_{i=0}^{11} \mathfrak{K}_i)^{-1}]P_{12} + [(\sum_{i=0}^{13} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{12} \mathfrak{K}_i)^{-1}]P_{13} + [(\sum_{i=0}^{14} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{13} \mathfrak{K}_i)^{-1}]P_{14} + [(\sum_{i=0}^{15} \mathfrak{K}_i)^{-1} - \\ & (\sum_{i=0}^{14} \mathfrak{K}_i)^{-1}]P_{15} + [(\sum_{i=0}^{16} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{15} \mathfrak{K}_i)^{-1}]P_{16} + [(\sum_{i=0}^{17} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{16} \mathfrak{K}_i)^{-1}]P_{17} + [(\sum_{i=0}^{18} \mathfrak{K}_i)^{-1} - \\ & (\sum_{i=0}^{17} \mathfrak{K}_i)^{-1}]P_{18} + [(\sum_{i=0}^{19} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{18} \mathfrak{K}_i)^{-1}]P_{19} + [(\sum_{i=0}^{20} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{19} \mathfrak{K}_i)^{-1}]P_{20} + [(\sum_{i=0}^{21} \mathfrak{K}_i)^{-1} - \\ & (\sum_{i=0}^{20} \mathfrak{K}_i)^{-1}]P_{21} + [(\sum_{i=0}^{22} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{21} \mathfrak{K}_i)^{-1}]P_{22} + [(\sum_{i=0}^{23} \mathfrak{K}_i)^{-1} - (\sum_{i=0}^{22} \mathfrak{K}_i)^{-1}]P_{23} + [(\sum_{i=0}^{24} \mathfrak{K}_i)^{-1} - \\ & (\sum_{i=0}^{23} \mathfrak{K}_i)^{-1}]P_{24}. \end{aligned}$$

3. Conclusion

In this scientific work, we studied the Invertibility properties of all plithogenic square matrices of high orders between 20 and 24, where we presented the mathematical conditions and formulas for computing the inverses of all plithogenic square matrices with real plithogenic entries. This goal will be completed by proving many theorems that describe the Invertibility of all classic matrix parts of the corresponding symbolic m-plithogenic matrix.'

Acknowledgement:

The authors extend their appreciation to the Arab Open University for supporting this work.

References

- [1] Merkepçi, H., and Abobala, M., "On The Symbolic 2-Plithogenic Rings", International Journal of Neutrosophic Science, 2023.
- [2] Smarandache, F., "Introduction to the Symbolic Plithogenic Algebraic Structures (revisited)", Neutrosophic Sets and Systems, vol. 53, 2023.
- [3] Taffach, N., and Ben Othman, K., "An Introduction to Symbolic 2-Plithogenic Modules Over Symbolic 2-Plithogenic Rings", Neutrosophic Sets and Systems, Vol 54, 2023.
- [4] Albasheer, O., Hajjari, A., and Dalla, R., "On The Symbolic 3-Plithogenic Rings and Their Algebraic Properties", Neutrosophic Sets and Systems, Vol 54, 2023.
- [5] Ben Othman, K., "On Some Algorithms For Solving Symbolic 3-Plithogenic Equations", Neoma Journal Of Mathematics and Computer Science, 2023.
- [6] Rawashdeh, A., "An Introduction To The Symbolic 3-plithogenic Number Theory", Neoma Journal Of Mathematics and Computer Science, 2023.
- [7] Hazaymeh, Ayman. "Time Effective Fuzzy Soft Set and Its Some Applications with and Without a Neutrosophic." International Journal of Neutrosophic Science 23.2 (2023): 129-29.
- [8] Merkepçi, H., "On Novel Results about the Algebraic Properties of Symbolic 3-Plithogenic and 4-Plithogenic Real Square Matrices", Symmetry, MDPI, 2023.
- [9] Soueycatt, M., Charchekhandra, B., Abu Hakmeh, R., "On The Algebraic Properties Of Symbolic 6-Plithogenic Integers", Neutrosophic Sets and Systems, vol. 59, 2023.
- [10] Ben Othman, K., Von Shtawzen, O., Khaldi, A., and Ali, R., "On The Concept Of Symbolic 7-Plithogenic Real Matrices", Pure Mathematics For Theoretical Computer Science, Vol.1, 2023.
- [11] Ben Othman, K., Von Shtawzen, O., Khaldi, A., and Ali, R., "On The Symbolic 8-Plithogenic Matrices", Pure Mathematics For Theoretical Computer Science, Vol.1, 2023.
- [12] Merkepçi, M., and Abobala, M., "On Some Novel Results About Split-Complex Numbers, The Diagonalization Problem And Applications To Public Key Asymmetric Cryptography", Journal of Mathematics, Hindawi, 2023.
- [13] Abobala, M., Hatip, A., and Bal, M., "A Review On Recent Advantages In Algebraic Theory Of Neutrosophic Matrices", International Journal of Neutrosophic Science, Vol.17, 2021.

- [14] Abobala, M., Ziena, M.B., Doewes, R.I., Hussein, Z. The Representation of Refined Neutrosophic Matrices By Refined Neutrosophic Linear Functions, *International Journal of Neutrosophic Science*, 2022, 19(1), pp. 342–349
- [15] Abobala, M., On Refined Neutrosophic Matrices and Their Applications In Refined Neutrosophic Algebraic Equations, *Journal Of Mathematics*, Hindawi, 2021
- [16] Olgun, N., Hatip, A., Bal, M., and Abobala, M., " A Novel Approach To Necessary and Sufficient Conditions For The Diagonalization of Refined Neutrosophic Matrices", *International Journal of neutrosophic Science*, Vol. 16, pp. 72-79, 2021.
- [17] Soueycatt, M., Charchekhandra, B., Abu Hakmeh, R., " On The Foundations Of Symbolic 5-Plithogenic Number Theory", *Neutrosophic Sets and Systems*, vol. 59, 2023.
- [18] Merkepçi, H., and Rawashdeh, A., " On The Symbolic 2-Plithogenic Number Theory and Integers ", *Neutrosophic Sets and Systems*, Vol 54, 2023.
- [19] avier Gamboa-Cruzado, Renatto Oyague Guerra, Enrique Condor Tinoco, Guillermo Paucar-Carlos, Juan Gamarra Moreno, Warshine Barry. (2024). A Study of the 16-Plithogenic and 17-Plithogenic Square Real Matrices and Their Properties. *Journal of International Journal of Neutrosophic Science*, 23 (3), 63-76 (Doi : <https://doi.org/10.54216/IJNS.230306>).
- [20] Smarandache F., and Abobala, M., "n-Refined Neutrosophic Vector Spaces", *International Journal of Neutrosophic Science*, Vol. 7, pp. 47-54. 2020.
- [21] Abobala, M.,. "A Study of AH-Substructures in n-Refined Neutrosophic Vector Spaces", *International Journal of Neutrosophic Science*", Vol. 9, pp.74-85. 2020.
- [22] Abobala, M., "On The Characterization of Maximal and Minimal Ideals In Several Neutrosophic Rings", *Neutrosophic Sets and Systems*, Vol. 45, 2021.
- [23] Nader Mahmoud Taffach , Ahmed Hatip., "A Review on Symbolic 2-Plithogenic Algebraic Structures " *Galoitica Journal Of Mathematical Structures and Applications*, Vol.5, 2023.
- [24] Hatamleh, R., Zolotarev, V. A. (2014). On Two-Dimensional Model Representations of One Class of Commuting Operators. *Ukrainian Mathematical Journal*, 66(1), 122–144.
<https://doi.org/10.1007/s11253-014-0916-9>
- [25] Abobala, M., "Neutrosophic Real Inner Product Spaces", *Neutrosophic Sets and Systems*, Vol. 43, 2021.
- [26] P. Prabakaran, Bilal Abdallah, Turayeva Dinara Tulkunovna. (2024). On Certain Algebraic Properties of Symbolic 3-Plithogenic Real Square Matrices. *International Journal of Neutrosophic Science*, 23 (4), 350-357 (Doi : <https://doi.org/10.54216/IJNS.230427>).
- [27] Necati Olgun. (2023). On The 15-Plithogenic Square Real Matrices. *Galoitica: Journal of Mathematical Structures and Applications*, 9 (2), 41-48 (Doi : <https://doi.org/10.54216/GJMSA.090205>).
- [28] Noor Edin Rabeh, Othman Al-Basheer, Sara Sawalmeh, Rozina Ali. (2023). An Algebraic Approach to n-Plithogenic Square Matrices For $18 \leq n \leq 19$. *Journal of Neutrosophic and Fuzzy Systems*, 7 (2), 08-23 (Doi : <https://doi.org/10.54216/JNFS.070201>).
- [29] Isra Al-Shbeil, Wael Mahmoud M. Salameh, Hossam Almahasneh, Khalid Kaabneh, Mutaz Shatnawi, Khaled Al-Husban. (2024). Algorithms for Computing M-Plithogenic Eigen Values and Vectors for Some Special Values of M. *International journal of neutrosophic science* , 24 (1), 128-135 (Doi: <https://doi.org/10.54216/IJNS.240112>
- [30] Mayada Abualhomos, Ahmed Atallah Alsaraireh, Malik Bataineh, Nabeela Abu-Alkishik, Isra Al-Shbeil, Abdallah Al-Husban. (2024). On The Real Inner Products and Orthogonality for Plithogenic Vector Spaces of Orders 4 and 5. *International journal of neutrosophic science* , 24 (1), 94-103 (Doi : <https://doi.org/10.54216/IJNS.240109>)