

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/359296228>

# Some Results in Classes Of Neutrosophic Graphs

**Preprint** · March 2022

DOI: 10.20944/preprints202203.0248.v1

---

CITATIONS

0

---

READ

1

**1 author:**



Henry Garrett

**174** PUBLICATIONS **238** CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Number Graphs And Numbers [View project](#)



Neutrosophic SuperHyperGraphs and SuperHyperGraphs [View project](#)

# Some Results in Classes Of Neutrosophic Graphs

Henry Garrett

Independent Researcher

DrHenryGarrett@gmail.com

Twitter's ID: @DrHenryGarrett | ©DrHenryGarrett.wordpress.com

## Abstract

New setting is introduced to study co-neighborhood, neutrosophic t-neighborhood, neutrosophic quasi-vertex set, neutrosophic quasi-order, neutrosophic neighborhood, neutrosophic co-t-neighborhood, neutrosophic quasi-edge set, neutrosophic quasi-size, Neutrosophic number, neutrosophic co-neighborhood, co-neutrosophic number, quasi-number and quasi-co-number. Some classes of neutrosophic graphs are investigated.

**Keywords:** Neutrosophic Quasi-Order, Neutrosophic Quasi-Size, Neutrosophic Quasi-Number, Neutrosophic Quasi-Co-Number, Neutrosophic Co-t-Neighborhood  
**AMS Subject Classification:** 05C17, 05C22, 05E45

## 1 Background

Fuzzy set in **Ref.** [16], neutrosophic set in **Ref.** [3], related definitions of other sets in **Refs.** [3, 13, 15], graphs and new notions on them in **Refs.** [1, 4, 8–11, 14, 17], neutrosophic graphs in **Ref.** [5], studies on neutrosophic graphs in **Ref.** [2], relevant definitions of other graphs based on fuzzy graphs in **Ref.** [12], related definitions of other graphs based on neutrosophic graphs in **Ref.** [6], are proposed. Also, some studies and researches about neutrosophic graphs, are proposed as a book in **Ref.** [7].

## 2 Preliminaries

In this subsection, basic material which is used in this article, is presented. Also, new ideas and their clarifications are elicited.

Basic idea is about the model which is used. First definition introduces basic model.

**Definition 2.1.** (Graph).

$G = (V, E)$  is called a **graph** if  $V$  is a set of objects and  $E$  is a subset of  $V \times V$  ( $E$  is a set of 2-subsets of  $V$ ) where  $V$  is called **vertex set** and  $E$  is called **edge set**. Every two vertices have been corresponded to at most one edge.

Neutrosophic graph is the foundation of results in this paper which is defined as follows. Also, some related notions are demonstrated.

**Definition 2.2.** (Neutrosophic Graph And Its Special Case).

$NTG = (V, E, \sigma = (\sigma_1, \sigma_2, \sigma_3), \mu = (\mu_1, \mu_2, \mu_3))$  is called a **neutrosophic graph** if it's graph,  $\sigma_i : V \rightarrow [0, 1]$ , and  $\mu_i : E \rightarrow [0, 1]$ . We add one condition on it and we use **special case** of neutrosophic graph but with same name. The added condition is as follows, for every  $v_i v_j \in E$ ,

$$\mu(v_i v_j) \leq \sigma(v_i) \wedge \sigma(v_j).$$

(i) :  $\sigma$  is called **neutrosophic vertex set**.

(ii) :  $\mu$  is called **neutrosophic edge set**.

(iii) :  $|V|$  is called **order** of NTG and it's denoted by  $\mathcal{O}(NTG)$ .

(iv) :  $\sum_{v \in V} \sigma(v)$  is called **neutrosophic order** of NTG and it's denoted by  $\mathcal{O}_n(NTG)$ .

(v) :  $|E|$  is called **size** of NTG and it's denoted by  $\mathcal{S}(NTG)$ .

(vi) :  $\sum_{e \in E} \sum_{i=1}^3 \mu_i(e)$  is called **neutrosophic size** of NTG and it's denoted by  $\mathcal{S}_n(NTG)$ .

Some classes of well-known neutrosophic graphs are defined. These classes of neutrosophic graphs are used to form this study and the most results are about them.

**Definition 2.3.** Let  $NTG : (V, E, \sigma, \mu)$  be a neutrosophic graph. Then

(i) : a sequence of vertices  $P : x_0, x_1, \dots, x_n$  is called **path** where  $x_i x_{i+1} \in E$ ,  $i = 0, 1, \dots, n-1$ ;

(ii) : **strength** of path  $P : x_0, x_1, \dots, x_n$  is  $\bigwedge_{i=0, \dots, n-1} \mu(x_i x_{i+1})$ ;

(iii) : **connectedness** amid vertices  $x_0$  and  $x_n$  is

$$\mu^\infty(x, y) = \bigwedge_{P: x_0, x_1, \dots, x_n} \bigwedge_{i=0, \dots, n-1} \mu(x_i x_{i+1});$$

(iv) : a sequence of vertices  $P : x_0, x_1, \dots, x_n$  is called **cycle** where  $x_i x_{i+1} \in E$ ,  $i = 0, 1, \dots, n-1$  and there are two edges  $xy$  and  $uv$  such that  $\mu(xy) = \mu(uv) = \bigwedge_{i=0, 1, \dots, n-1} \mu(v_i v_{i+1})$ ;

(v) : it's **t-partite** where  $V$  is partitioned to  $t$  parts,  $V_1, V_2, \dots, V_t$  and the edge  $xy$  implies  $x \in V_i$  and  $y \in V_j$  where  $i \neq j$ . If it's complete, then it's denoted by  $K_{\sigma_1, \sigma_2, \dots, \sigma_t}$  where  $\sigma_i$  is  $\sigma$  on  $V_i$  instead  $V$  which mean  $x \notin V_i$  induces  $\sigma_i(x) = 0$ ;

(vi) : t-partite is **complete bipartite** if  $t = 2$ , and it's denoted by  $K_{\sigma_1, \sigma_2}$ ;

(vii) : complete bipartite is **star** if  $|V_1| = 1$ , and it's denoted by  $S_{1, \sigma_2}$ ;

(viii) : a vertex in  $V$  is **center** if the vertex joins to all vertices of a cycle. Then it's **wheel** and it's denoted by  $W_{1, \sigma_2}$ ;

(ix) : it's **complete** where  $\forall uv \in V$ ,  $\mu(uv) = \sigma(u) \wedge \sigma(v)$ ;

(x) : it's **strong** where  $\forall uv \in E$ ,  $\mu(uv) = \sigma(u) \wedge \sigma(v)$ .

The notions of neighbor and neighborhood are about some vertices which have one edge with a fixed vertex. These notions present vertices which are close to a fixed vertex as possible. Based on strong edge, it's possible to define different neighborhood as follows.

**Definition 2.4.** (Neighborhood).

Let  $NTG : (V, E, \sigma, \mu)$  be a neutrosophic graph. Suppose  $x \in V$ . Then

$$N(x) = \{y \in V \mid xy \in E\}.$$

**Definition 2.5.** (Co-Neighborhood).

Let  $NTG : (V, E, \sigma, \mu)$  be a neutrosophic graph and  $t \in \mathbb{N}$ . Suppose  $x_i \in V$ ,  $i = 1, 2, \dots, t$ . Then

(i)

$$N_t(x_1, x_2, \dots, x_t) = \{y \in V \mid x_i y \in E, i = 1, 2, \dots, t\}.$$

If  $\min_{x_1, x_2, \dots, x_t \in V} N_t[x_1, x_2, \dots, x_t] = V$ , then **neutrosophic t-neighborhood** is called **neutrosophic quasi-vertex set** and  $t$  is called **neutrosophic quasi-order**. Generally, when  $t$  isn't fixed, it's called **neutrosophic neighborhood**. **Neutrosophic number** is

$$\sum_{\min_{x_1, x_2, \dots, x_t \in V} N_t[x_1, x_2, \dots, x_t] = V} \sum_{j=1}^3 \sigma_j(x_i).$$

(ii)

$$N_t(x_1, x_2, \dots, x_t) = \{e \in E \mid e = x_i y \in E, i = 1, 2, \dots, t\}.$$

If  $\min_{x_1, x_2, \dots, x_t \in V} N_t[x_1, x_2, \dots, x_t] = E$ , then **neutrosophic co-t-neighborhood** is called **neutrosophic quasi-edge set** and  $t$  is called **neutrosophic quasi-size**. Generally, when  $t$  isn't fixed, it's called **neutrosophic co-neighborhood**. **Co-neutrosophic number** is

$$\sum_{\min_{x_1, x_2, \dots, x_t \in V} N_t[x_1, x_2, \dots, x_t] = E} \sum_{j=1}^3 \sigma_j(x_i).$$

### 3 Neutrosophic Quasi-Order

**Proposition 3.1.** Let  $NTG : (V, E, \sigma, \mu)$  be an odd path. Then  $\{x_2, x_4, \dots, x_{n-1}\}$  is related to quasi-order  $\lfloor \frac{n}{2} \rfloor$ .

*Proof.* Suppose  $NTG : (V, E, \sigma, \mu)$  be an odd path. Thus  $NTG : (V, E, \sigma, \mu)$  is  $P : x_1, x_2, \dots, x_n$  where  $n$  and 1 has same parity. There are two sets.  $\{x_1, x_3, \dots, x_n\}$  has quasi-order  $\lfloor \frac{n}{2} \rfloor + 1$  but  $\{x_2, x_4, \dots, x_{n-1}\}$  has quasi-order  $\lfloor \frac{n}{2} \rfloor$  which is minimum number amid these two sets. So  $\{x_2, x_4, \dots, x_{n-1}\}$  is related to quasi-order  $\lfloor \frac{n}{2} \rfloor$ .  $\{x_2, x_4, \dots, x_{n-1}\}$  is quasi-vertex set which is optimal.  $\square$

**Proposition 3.2.** Let  $NTG : (V, E, \sigma, \mu)$  be an even path. Then  $\{x_2, x_4, \dots, x_n\}$  is related to quasi-order  $\frac{n}{2}$ .

*Proof.* Suppose  $NTG : (V, E, \sigma, \mu)$  be an even path. Thus  $NTG : (V, E, \sigma, \mu)$  is  $P : x_1, x_2, \dots, x_n$  where  $n$  and 1 has different parity. There are two sets.  $\{x_1, x_3, \dots, x_{n-1}\}$  has quasi-order  $\lfloor \frac{n}{2} \rfloor$  and  $\{x_2, x_4, \dots, x_n\}$  has quasi-order  $\lfloor \frac{n}{2} \rfloor$  which is minimum number like first set. So  $\{x_2, x_4, \dots, x_n\}$  and  $\{x_1, x_3, \dots, x_{n-1}\}$  are related to quasi-order  $\lfloor \frac{n}{2} \rfloor$ .  $\{x_2, x_4, \dots, x_n\}$  and  $\{x_1, x_3, \dots, x_{n-1}\}$  are quasi-vertex sets which are optimal.  $\square$

**Proposition 3.3.** Let  $NTG : (V, E, \sigma, \mu)$  be an odd cycle. Then  $\{x_2, x_4, \dots, x_{n-1}\}$  is related to quasi-order  $\lfloor \frac{n}{2} \rfloor$ .

*Proof.* Suppose  $NTG : (V, E, \sigma, \mu)$  be an odd cycle. Thus  $NTG : (V, E, \sigma, \mu)$  is  $P : x_1, x_2, \dots, x_n, x_1$  where  $n$  and 1 has same parity. There are two sets.  $\{x_1, x_3, \dots, x_n\}$  has quasi-order  $\lfloor \frac{n}{2} \rfloor + 1$  but  $\{x_2, x_4, \dots, x_{n-1}\}$  has quasi-order  $\lfloor \frac{n}{2} \rfloor$  which is minimum number amid these two sets. So  $\{x_2, x_4, \dots, x_{n-1}\}$  is related to quasi-order  $\lfloor \frac{n}{2} \rfloor$ .  $\{x_2, x_4, \dots, x_{n-1}\}$  is quasi-vertex set which is optimal.  $\square$

**Proposition 3.4.** Let  $NTG : (V, E, \sigma, \mu)$  be an even cycle. Then  $\{x_2, x_4, \dots, x_n\}$  and  $\{x_1, x_3, \dots, x_{n-1}\}$  are related to quasi-order  $\frac{n}{2}$ .

*Proof.* Suppose  $NTG : (V, E, \sigma, \mu)$  be an even cycle. Thus  $NTG : (V, E, \sigma, \mu)$  is  $P : x_1, x_2, \dots, x_n, x_1$  where  $n$  and 1 has different parity. There are two sets.  $\{x_1, x_3, \dots, x_{n-1}\}$  has quasi-order  $\lfloor \frac{n}{2} \rfloor$  and  $\{x_2, x_4, \dots, x_n\}$  has quasi-order  $\lfloor \frac{n}{2} \rfloor$  which is minimum number like first set. So  $\{x_2, x_4, \dots, x_n\}$  and  $\{x_1, x_3, \dots, x_{n-1}\}$  are related to quasi-order  $\lfloor \frac{n}{2} \rfloor$ .  $\{x_2, x_4, \dots, x_n\}$  and  $\{x_1, x_3, \dots, x_{n-1}\}$  are quasi-vertex sets which are optimal.  $\square$

**Proposition 3.5.** Let  $NTG : (V, E, \sigma, \mu)$  be complete. Then  $\{x\}$  is related to quasi-order 1.

*Proof.* Let  $NTG : (V, E, \sigma, \mu)$  be complete. Suppose  $x$  is a given vertex. Thus

$$N[x] = \min_{x_1, x_2, \dots, x_t \in V} N_t[x_1, x_2, \dots, x_t] = V.$$

It implies the set  $\{x\}$  which is 1-neighborhood, is quasi-vertex set. Then  $\{x\}$  is related to quasi-order 1.  $\square$

**Proposition 3.6.** Let  $NTG : (V, E, \sigma, \mu)$  be star. Then  $\{x\}$  is related to quasi-order 1.

*Proof.* Let  $NTG : (V, E, \sigma, \mu)$  be star. Suppose  $c$  is the center. Thus

$$N[c] = \min_{x_1, x_2, \dots, x_t \in V} N_t[x_1, x_2, \dots, x_t] = V.$$

It implies the set  $\{c\}$  which is 1-neighborhood, is quasi-vertex set. Then  $\{c\}$  is related to quasi-order 1.  $\square$

## 4 Neutrosophic Quasi-Size

**Proposition 4.1.** Let  $NTG : (V, E, \sigma, \mu)$  be an odd path. Then either  $\{x_1, x_3, \dots, x_n\}$  or  $\{x_1, x_3, \dots, x_{n-1}\}$  are related to quasi-size  $\lfloor \frac{n}{3} \rfloor$ .

*Proof.* Suppose  $NTG : (V, E, \sigma, \mu)$  be an odd path. Thus  $NTG : (V, E, \sigma, \mu)$  is  $P : x_1, x_2, \dots, x_n$  where  $n$  and 1 has same parity. There is two sets. Either  $\{x_1, x_3, \dots, x_n\}$  or  $\{x_1, x_3, \dots, x_{n-1}\}$  has quasi-size  $\lfloor \frac{n}{3} \rfloor$  which are minimum number amid these all sets. So either  $\{x_1, x_3, \dots, x_n\}$  or  $\{x_1, x_3, \dots, x_{n-1}\}$  is related to quasi-size  $\lfloor \frac{n}{3} \rfloor$ . Either  $\{x_1, x_3, \dots, x_n\}$  or  $\{x_1, x_3, \dots, x_{n-1}\}$  are quasi-vertex set which are optimal.  $\square$

**Proposition 4.2.** Let  $NTG : (V, E, \sigma, \mu)$  be an even path. Then  $\{x_2, x_4, \dots, x_n\}$  is related to quasi-size  $\lfloor \frac{n}{3} \rfloor$ .

*Proof.* Suppose  $NTG : (V, E, \sigma, \mu)$  be an even path. Thus  $NTG : (V, E, \sigma, \mu)$  is  $P : x_1, x_2, \dots, x_n$  where  $n$  and 1 has different parity. There are two sets. Either  $\{x_1, x_3, \dots, x_n\}$  or  $\{x_1, x_3, \dots, x_{n-1}\}$ .  $\{x_1, x_3, \dots, x_n\}$  has quasi-size  $\lfloor \frac{n}{3} \rfloor$  and  $\{x_1, x_3, \dots, x_{n-1}\}$  has quasi-size  $\lfloor \frac{n}{3} \rfloor$  which is minimum number like first set. So either  $\{x_1, x_3, \dots, x_n\}$  or  $\{x_1, x_3, \dots, x_{n-1}\}$  is related to quasi-size  $\lfloor \frac{n}{3} \rfloor$ . Either  $\{x_1, x_3, \dots, x_n\}$  or  $\{x_1, x_3, \dots, x_{n-1}\}$  is quasi-vertex set which is optimal.  $\square$

**Proposition 4.3.** Let  $NTG : (V, E, \sigma, \mu)$  be an odd cycle. Then  $\{x_1, x_4, \dots, x_{n-3}\}$  is related to quasi-size  $\lfloor \frac{n}{3} \rfloor$ .

*Proof.* Suppose  $NTG : (V, E, \sigma, \mu)$  be an odd cycle. Thus  $NTG : (V, E, \sigma, \mu)$  is  $P : x_1, x_2, \dots, x_n, x_1$  where  $n$  and 1 has same parity. There are two sets.  $\{x_1, x_3, \dots, x_n\}$  has quasi-size  $\lfloor \frac{n}{3} \rfloor$  and  $\{x_1, x_3, \dots, x_{n-1}\}$  has quasi-size  $\lfloor \frac{n}{3} \rfloor$  which are minimum numbers amid all sets. So either  $\{x_1, x_3, \dots, x_{n-1}\}$  or  $\{x_1, x_3, \dots, x_n\}$  is related to quasi-size  $\lfloor \frac{n}{3} \rfloor$ . Either  $\{x_1, x_3, \dots, x_{n-1}\}$  or  $\{x_1, x_3, \dots, x_n\}$  is quasi-edge set which is optimal.  $\square$

**Proposition 4.4.** Let  $NTG : (V, E, \sigma, \mu)$  be an even cycle. Then  $\{x_1, x_4, \dots, x_{n-3}\}$  is related to quasi-size  $\lfloor \frac{n}{3} \rfloor$ .

*Proof.* Suppose  $NTG : (V, E, \sigma, \mu)$  be an even cycle. Thus  $NTG : (V, E, \sigma, \mu)$  is  $P : x_1, x_2, \dots, x_n, x_1$  where  $n$  and 1 has different parity. There are two sets.  $\{x_1, x_3, \dots, x_{n-1}\}$  has quasi-size  $\lfloor \frac{n}{3} \rfloor$  and  $\{x_1, x_3, \dots, x_{n-1}\}$  has quasi-size  $\lfloor \frac{n}{3} \rfloor$  which is minimum number like first set. So either  $\{x_1, x_3, \dots, x_n\}$  or  $\{x_1, x_3, \dots, x_{n-1}\}$  are related to quasi-size  $\lfloor \frac{n}{3} \rfloor$  which this number is optimal. Either  $\{x_1, x_3, \dots, x_n\}$  or  $\{x_1, x_3, \dots, x_{n-1}\}$  is quasi-edge set which are optimal.  $\square$

**Proposition 4.5.** Let  $NTG : (V, E, \sigma, \mu)$  be complete. Then  $\{x_1, x_2, \dots, x_{n-1}\}$  is related to quasi-size  $n - 1$ .

*Proof.* Let  $NTG : (V, E, \sigma, \mu)$  be complete. Suppose  $x$  is a given vertex. Thus

$$N[x_1, x_2, \dots, x_{n-1}] = \min_{x_1, x_2, \dots, x_{n-1} \in V} N_t[x_1, x_2, \dots, x_{n-1}] = E.$$

It implies the set  $\{x_1, x_2, \dots, x_{n-1}\}$  which is co- $(n - 1)$ -neighborhood, is only quasi-vertex set as optimal set. Then  $\{x_1, x_2, \dots, x_{n-1}\}$  is related to quasi-size  $n - 1$ .  $\square$

**Proposition 4.6.** Let  $NTG : (V, E, \sigma, \mu)$  be star. Then  $\{x\}$  is related to quasi-size 1.

*Proof.* Let  $NTG : (V, E, \sigma, \mu)$  be star. Suppose  $c$  is the center. Thus

$$N[c] = \min_{x_1, x_2, \dots, x_t \in V} N_t[x_1, x_2, \dots, x_t] = E.$$

It implies the set  $\{c\}$  which is 1-neighborhood, is quasi-vertex set. Then  $\{c\}$  is related to quasi-order 1.  $\square$

## 5 Setting of Neutrosophic Quasi-Number and Quasi-Co-Number

**Definition 5.1.** (Quasi-Number & Quasi-Co-Number).

Let  $NTG : (V, E, \sigma, \mu)$  be a neutrosophic graph. Then

(i) **quasi-number** for  $x_0$  is

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1});$$

(ii) **quasi-co-number** for  $x_0$  is

$$\mathcal{C}_c(x_0) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\}.$$

**Proposition 5.2.** Let  $NTG : (V, E, \sigma, \mu)$  be a complete-neutrosophic graph. Then

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1}^{\mathcal{O}} \mu(x_0 x_i).$$

*Proof.* Assume  $NTG : (V, E, \sigma, \mu)$  is a complete-neutrosophic graph and  $x_0 x'_0 \in E$ . Then

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1}^{\mathcal{O}} \mu(x_0 x_i).$$

Since

$$\bigvee_{P: x_0, x_1, \dots, x'_0} \bigwedge_{x_i x_{i+1} \in V(P)} \mu(x_i x_{i+1}) = \mu(x_0 x'_0).$$

and

$$\bigvee_{P: x_0, x_1, \dots, x'_0} \bigwedge_{x_i x_{i+1} \in V(P)} \mu(x_i x_{i+1}) \leq \mu(x_0 x'_0)$$

hold if  $NTG : (V, E, \sigma, \mu)$  is a complete-neutrosophic graph. Also,

$$\bigvee_{P: x_0, x_1, \dots, x'_0} \bigwedge_{x_i x_{i+1} \in V(P)} \mu(x_i x_{i+1}) \geq \mu(x_0 x'_0)$$

holds since  $x_0 x'_0$  is a path from  $x_0$  to  $x'_0$ . □ 135

**Proposition 5.3.** Let  $NTG : (V, E, \sigma, \mu)$  be a path-neutrosophic graph. Suppose  $x_0$  is a leaf and  $x_0 z \in E$ . Then □ 136

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z).$$

*Proof.* Assume  $NTG : (V, E, \sigma, \mu)$  is a path-neutrosophic graph. Suppose  $x_0$  is a leaf and  $x_0 z \in E$ . An arbitrary path  $P$  has consecutive vertices  $x_0, z$  as their ends from a given vertex to leaf  $z$ . Thus

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z).$$

It implies for every given vertex  $x_t \in V$ ,

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z).$$

Therefore,

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z). □ 137$$

**Proposition 5.4.** Let  $NTG : (V, E, \sigma, \mu)$  be a path-neutrosophic graph. Suppose  $x_0$  is a leaf and  $x_0 z \in E$  is weakest edge. Then

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 z).$$

*Proof.* Assume  $NTG : (V, E, \sigma, \mu)$  is a path-neutrosophic graph. Suppose  $x_0$  is a leaf and  $x_0z \in E$  is a weakest edge. An arbitrary path  $P$  has consecutive vertices  $x_0, z$  as their ends from a given vertex to leaf  $x_0$ . Thus

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z).$$

By  $x_0z \in E$  is a weakest edge,

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 z).$$

It implies for every given vertex  $x_t \in V$ ,

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 z).$$

Therefore,

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 z).$$

□ 138

**Proposition 5.5.** Let  $NTG : (V, E, \sigma, \mu)$  be a cycle-neutrosophic graph. Suppose  $x_0$  has a weakest edge  $x_0z \in E$ . Then

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

*Proof.* Assume  $NTG : (V, E, \sigma, \mu)$  is a cycle-neutrosophic graph. Suppose  $x_0$  has a weakest edge  $x_0z \in E$ . For every vertex, there are only two ways to form a path. By  $x_0z \in E$  is a weakest edge,

$$\mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

Thus

$$\bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

$P : x_0, z$  is a path from  $x_0$  to  $z$ . It implies

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

Therefore,

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

It means that

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

□ 139

**Proposition 5.6.** Let  $NTG : (V, E, \sigma, \mu)$  be a cycle-neutrosophic graph. Suppose  $x_0$  is incident to one edge  $x_0z \in E$  which isn't a weakest edge. Then

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$



*Proof.* Assume  $NTG : (V, E, \sigma, \mu)$  is a cycle-neutrosophic graph. Suppose  $x_0$  has an edge  $x_0z \in E$  which isn't a weakest edge. For every vertex, there are only two ways to form a path. By  $x_0z \in E$  isn't a weakest edge and  $P : x_0, z$  is a path from  $x_0$  to  $z$ . It implies

$$\bigvee_{P:x_0,x_1,\dots,x_t} \bigwedge_{i=0,\dots,t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

Therefore,

$$\bigvee_{x_t \in V} \bigvee_{P:x_0,x_1,\dots,x_t} \bigwedge_{i=0,\dots,t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

It means that

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P:x_0,x_1,\dots,x_t} \bigwedge_{i=0,\dots,t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

□ 140

**Proposition 5.7.** Let  $NTG : (V, E, \sigma, \mu)$  be a cycle-neutrosophic graph. Suppose  $x_0$  is incident to two edges  $x_0z, x_0z' \in E$  which aren't weakest edges. Then

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P:x_0,x_1,\dots,x_t} \bigwedge_{i=0,\dots,t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z) \vee \mu(x_0 z').$$

Furthermore, if  $\mu(x_0 z) < \mu(x_0 z')$ , then

$$\mathcal{C}(x_0) \leq \mu(x_0 z');$$

if  $\mu(x_0 z') < \mu(x_0 z)$ , then

$$\mathcal{C}(x_0) \leq \mu(x_0 z);$$

if  $\mu(x_0 z') = \mu(x_0 z)$ , then

$$\mathcal{C}(x_0) \leq \mu(x_0 z') = \mu(x_0 z).$$

*Proof.* Assume  $NTG : (V, E, \sigma, \mu)$  is a cycle-neutrosophic graph. Suppose  $x_0$  is incident to two edges  $x_0z, x_0z' \in E$  which aren't weakest edges. For every vertex, there are only two ways to form a path which are with edges either  $x_0z$  or  $x_0z'$ .  $P : x_0, z$  and  $P' : x_0, z'$  are paths from  $x_0$  to  $z$  and  $z'$ . It implies

$$\bigvee_{P:x_0,x_1,\dots,x_t} \bigwedge_{i=0,\dots,t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z) \vee \mu(x_0 z').$$

Therefore,

$$\bigvee_{x_t \in V} \bigvee_{P:x_0,x_1,\dots,x_t} \bigwedge_{i=0,\dots,t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z) \vee \mu(x_0 z').$$

It means that

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P:x_0,x_1,\dots,x_t} \bigwedge_{i=0,\dots,t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z) \vee \mu(x_0 z').$$

Furthermore, if  $\mu(x_0 z) < \mu(x_0 z')$ , then

$$\mathcal{C}(x_0) \leq \mu(x_0 z');$$

if  $\mu(x_0 z') < \mu(x_0 z)$ , then

$$\mathcal{C}(x_0) \leq \mu(x_0 z);$$

if  $\mu(x_0 z') = \mu(x_0 z)$ , then

$$\mathcal{C}(x_0) \leq \mu(x_0 z') = \mu(x_0 z).$$

□ 141

**Proposition 5.8.** Let  $NTG : (V, E, \sigma, \mu)$  be a star-neutrosophic graph. Suppose  $x_0$  is center. Then

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i).$$

*Proof.* Consider  $NTG : (V, E, \sigma, \mu)$  is a star-neutrosophic graph. Suppose  $x_0$  is center. All paths from  $x_0$  has a form  $x_0 x_i$ . Thus

$$\bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 x_i).$$

Hence

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i).$$

So

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i).$$

It implies

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i).$$

□ 142

**Proposition 5.9.** Let  $NTG : (V, E, \sigma, \mu)$  be a star-neutrosophic graph. Suppose  $x_0$  isn't center and  $c$  is a center. Then

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 c).$$

*Proof.* Consider  $NTG : (V, E, \sigma, \mu)$  is a star-neutrosophic graph. Suppose  $x_0$  isn't center and  $c$  is a center. There's only one path from  $x_0$  has a form  $x_0 c$ . Thus

$$\bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 c).$$

Hence

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 c).$$

So

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 c).$$

It implies

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 c).$$

□ 143

**Proposition 5.10.** Let  $NTG : (V, E, \sigma, \mu)$  be a star-strong-neutrosophic graph. Suppose  $x_0$  is center such that  $\bigwedge_{i=0}^{\mathcal{O}-1} = \sigma(x_0)$ . Then

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \sigma(x_0).$$

*Proof.* Consider  $NTG : (V, E, \sigma, \mu)$  is a star-neutrosophic graph. Suppose  $x_0$  is center such that  $\bigwedge_{i=0}^{\mathcal{O}-1} \mu(x_0 x_i) = \sigma(x_0)$ . All paths from  $x_0$  has a form  $x_0 x_i$ . Thus

$$\bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 x_i) = \sigma(x_0).$$

Hence

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i) = \sigma(x_0).$$

So

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i) = \sigma(x_0).$$

It implies

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \sigma(x_0).$$

□ 144

**Proposition 5.11.** Let  $NTG : (V, E, \sigma, \mu)$  be a complete-neutrosophic graph. Then

145

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{i=1}^{\mathcal{O}} \mu(x_0 x_i)\}. \end{aligned}$$

*Proof.* Assume  $NTG : (V, E, \sigma, \mu)$  is a complete-neutrosophic graph and  $x_0 x'_0 \in E$ . Then

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1}^{\mathcal{O}} \mu(x_0 x_i).$$

Since

$$\bigvee_{P: x_0, x_1, \dots, x'_0} \bigwedge_{x_i x_{i+1} \in V(P)} \mu(x_i x_{i+1}) = \mu(x_0 x'_0).$$

and

$$\bigvee_{P: x_0, x_1, \dots, x'_0} \bigwedge_{x_i x_{i+1} \in V(P)} \mu(x_i x_{i+1}) \leq \mu(x_0 x'_0)$$

hold if  $NTG : (V, E, \sigma, \mu)$  is a complete-neutrosophic graph. Also,

$$\bigvee_{P: x_0, x_1, \dots, x'_0} \bigwedge_{x_i x_{i+1} \in V(P)} \mu(x_i x_{i+1}) \geq \mu(x_0 x'_0)$$

holds since  $x_0 x'_0$  is a path from  $x_0$  to  $x'_0$ . So

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{i=1}^{\mathcal{O}} \mu(x_0 x_i)\}. \end{aligned}$$

□ 146

**Proposition 5.12.** Let  $NTG : (V, E, \sigma, \mu)$  be a path-neutrosophic graph. Suppose  $x_0$  is a leaf and  $x_0z \in E$ . Then

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) \leq \mu(x_0 z)\}. \end{aligned}$$

*Proof.* Assume  $NTG : (V, E, \sigma, \mu)$  is a path-neutrosophic graph. Suppose  $x_0$  is a leaf and  $x_0z \in E$ . An arbitrary path  $P$  has consecutive vertices  $x_0, z$  as their ends from a given vertex to leaf  $z$ . Thus

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z).$$

It implies for every given vertex  $x_t \in V$ ,

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z).$$

Therefore,

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z).$$

Thus

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) \leq \mu(x_0 z)\}. \end{aligned}$$

□ 147

**Proposition 5.13.** Let  $NTG : (V, E, \sigma, \mu)$  be a path-neutrosophic graph. Suppose  $x_0$  is a leaf and  $x_0z \in E$  is a weakest edge. Then

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \mu(x_0 z)\}. \end{aligned}$$

*Proof.* Assume  $NTG : (V, E, \sigma, \mu)$  is a path-neutrosophic graph. Suppose  $x_0$  is a leaf and  $x_0z \in E$  is a weakest edge. An arbitrary path  $P$  has consecutive vertices  $x_0, z$  as their ends from a given vertex to leaf  $x_0$ . Thus

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z).$$

By  $x_0z \in E$  is a weakest edge,

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 z).$$

It implies for every given vertex  $x_t \in V$ ,

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 z).$$

Therefore,

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 z).$$

Thus

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \mu(x_0 z)\}. \end{aligned}$$

□ 148

**Proposition 5.14.** Let  $NTG : (V, E, \sigma, \mu)$  be a cycle-neutrosophic graph. Suppose  $x_0$  has two weakest edges  $x_0 z, x_0 z' \in E$ . Then

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \mu(x_0 z) \vee \mu(x_0 z') = \mu(x_0 z) = \mu(x_0 z'), x_z x_{z'} \in E\} = \{x_0 z, x_0 z'\}. \end{aligned}$$

*Proof.* Assume  $NTG : (V, E, \sigma, \mu)$  is a cycle-neutrosophic graph. Suppose  $x_0$  has a weakest edge  $x_0 z \in E$ . For every vertex, there are only two ways to form a path. By  $x_0 z \in E$  is a weakest edge,

$$\mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

Thus

$$\bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

$P : x_0, z$  is a path from  $x_0$  to  $z$ . It implies

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

Therefore,

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

It means that

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

By  $x_0 z' \in E$  is a weakest edge,

$$\mu(x_i x_{i+1}) \geq \mu(x_0 z').$$

Thus

$$\bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z').$$

$P : x_0, z'$  is a path from  $x_0$  to  $z'$ . It implies

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z').$$

Therefore,

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z').$$

It means that

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z').$$

By for every vertex, there are only two ways to form a path,

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \mu(x_0 z) \vee \mu(x_0 z') = \mu(x_0 z) = \mu(x_0 z'), x_z x_{z'} \in E\} = \{x_0 z, x_0 z'\}. \end{aligned}$$

□ 149

**Proposition 5.15.** Let  $NTG : (V, E, \sigma, \mu)$  be a cycle-neutrosophic graph. Suppose  $x_0$  is incident to one edge  $x_0 z \in E$  which isn't a weakest edge. Then

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) \geq \mu(x_0 z), x_z x_{z'} \in E\}. \end{aligned}$$

*Proof.* Assume  $NTG : (V, E, \sigma, \mu)$  is a cycle-neutrosophic graph. Suppose  $x_0$  has an edge  $x_0 z \in E$  which isn't a weakest edge. For every vertex, there are only two ways to form a path. By  $x_0 z \in E$  isn't a weakest edge and  $P : x_0, z$  is a path from  $x_0$  to  $z$ . It implies

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

Therefore,

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

It means that

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \geq \mu(x_0 z).$$

Thus

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) \geq \mu(x_0 z), x_z x_{z'} \in E\}. \end{aligned}$$

□ 150

**Proposition 5.16.** Let  $NTG : (V, E, \sigma, \mu)$  be a cycle-neutrosophic graph. Suppose  $x_0$  is incident to two edges  $x_0 z, x_0 z' \in E$  which aren't weakest edges. Then

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) \leq \mu(x_0 z) \vee \mu(x_0 z'), x_z x_{z'} \in E\}. \end{aligned}$$

Furthermore, if  $\mu(x_0 z) < \mu(x_0 z')$ , then

$$\mathcal{C}_c(x_0) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) \leq \mu(x_0 z'), x_z x_{z'} \in E\};$$

if  $\mu(x_0 z') < \mu(x_0 z)$ , then

$$\mathcal{C}_c(x_0) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) \leq \mu(x_0 z), x_z x_{z'} \in E\};$$

if  $\mu(x_0 z') = \mu(x_0 z)$ , then

$$\mathcal{C}_c(x_0) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) \leq \mu(x_0 z) = \mu(x_0 z'), x_z x_{z'} \in E\}.$$

*Proof.* Assume  $NTG : (V, E, \sigma, \mu)$  is a cycle-neutrosophic graph. Suppose  $x_0$  is incident to two edges  $x_0z, x_0z' \in E$  which aren't weakest edges. For every vertex, there are only two ways to form a path which are with edges either  $x_0z$  or  $x_0z'$ .  $P : x_0, z$  and  $P' : x_0, z'$  are paths from  $x_0$  to  $z$  and  $z'$ . It implies

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z) \vee \mu(x_0 z').$$

Therefore,

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z) \vee \mu(x_0 z').$$

It means that

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) \leq \mu(x_0 z) \vee \mu(x_0 z').$$

Furthermore, if  $\mu(x_0 z) < \mu(x_0 z')$ , then

$$\mathcal{C}(x_0) \leq \mu(x_0 z');$$

if  $\mu(x_0 z') < \mu(x_0 z)$ , then

$$\mathcal{C}(x_0) \leq \mu(x_0 z);$$

if  $\mu(x_0 z') = \mu(x_0 z)$ , then

$$\mathcal{C}(x_0) \leq \mu(x_0 z') = \mu(x_0 z).$$

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) \leq \mu(x_0 z) \vee \mu(x_0 z'), x_z x_{z'} \in E\}. \end{aligned}$$

Furthermore, if  $\mu(x_0 z) < \mu(x_0 z')$ , then

$$\mathcal{C}_c(x_0) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) \leq \mu(x_0 z'), x_z x_{z'} \in E\};$$

if  $\mu(x_0 z') < \mu(x_0 z)$ , then

$$\mathcal{C}_c(x_0) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) \leq \mu(x_0 z), x_z x_{z'} \in E\};$$

if  $\mu(x_0 z') = \mu(x_0 z)$ , then

$$\mathcal{C}_c(x_0) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) \leq \mu(x_0 z) = \mu(x_0 z'), x_z x_{z'} \in E\}.$$

□ 151

**Proposition 5.17.** Let  $NTG : (V, E, \sigma, \mu)$  be a star-neutrosophic graph. Suppose  $x_0$  is center. Then

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i)\}. \end{aligned}$$

152

153

*Proof.* Consider  $NTG : (V, E, \sigma, \mu)$  is a star-neutrosophic graph. Suppose  $x_0$  is center. All paths from  $x_0$  has a form  $x_0x_i$ . Thus

$$\bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 x_i).$$

Hence

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i).$$

So

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i).$$

It implies

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i).$$

Thus

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i)\}. \end{aligned}$$

□ 154

**Proposition 5.18.** Let  $NTG : (V, E, \sigma, \mu)$  be a star-neutrosophic graph. Suppose  $x_0$  isn't center and  $c$  is a center. Then

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \mu(c x_0)\} = \{c x_0\}. \end{aligned}$$

*Proof.* Consider  $NTG : (V, E, \sigma, \mu)$  is a star-neutrosophic graph. Suppose  $x_0$  isn't center and  $c$  is a center. There's only one path from  $x_0$  has a form  $x_0 c$ . Thus

$$\bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 c).$$

Hence

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 c).$$

So

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 c).$$

It implies

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 c).$$

Thus

$$\begin{aligned} \mathcal{C}_c(x_0) &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ &= \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \mu(c x_0)\} = \{c x_0\}. \end{aligned}$$

□ 155



**Proposition 5.19.** Let  $NTG : (V, E, \sigma, \mu)$  be a star-strong-neutrosophic graph. Suppose  $x_0$  is center such that  $\bigwedge_{i=0}^{\mathcal{O}-1} = \sigma(x_0)$ . And  $x$  isn't center. Then

$$\mathcal{C}_c(x_0) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \sigma(x_0)\} = \{x_0 x_i\}_{i=1}^{\mathcal{O}-1}.$$

And

$$\mathcal{C}_c(x) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \sigma(x_0)\} = \{x_0 x_i\}_{i=1}^{\mathcal{O}-1}.$$

*Proof.* Consider  $NTG : (V, E, \sigma, \mu)$  is a star-neutrosophic graph. Suppose  $x_0$  is center such that  $\bigwedge_{i=0}^{\mathcal{O}-1} = \sigma(x_0)$ . All paths from  $x_0$  has a form  $x_0 x_i$ . Thus

$$\bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \mu(x_0 x_i) = \sigma(x_0).$$

Hence

$$\bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i) = \sigma(x_0).$$

So

$$\bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \bigvee_{i=1, \dots, \mathcal{O}} \mu(x_0 x_i) = \sigma(x_0).$$

It implies

$$\mathcal{C}(x_0) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1}) = \sigma(x_0).$$

Thus

$$\mathcal{C}_c(x_0) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \sigma(x_0)\} = \{x_0 x_i\}_{i=1}^{\mathcal{O}-1}.$$

And

$$\mathcal{C}_c(x) = \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \bigvee_{x_t \in V} \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1})\} = \\ \{x_z x_{z'} \mid \mu(x_z x_{z'}) = \sigma(x_0)\} = \{x_0 x_i\}_{i=1}^{\mathcal{O}-1}.$$

□ 156

## References 157

1. N.H. Abughazalah, and M. Khan, “Selection of Optimum Nonlinear Confusion Component of Information Confidentiality Mechanism Using Grey Theory Based Decision-Making Technique.” Preprints 2021, 2021120310 (doi: 10.20944/preprints202112.0310.v1). 158 159 160 161
2. M. Akram, and G. Shahzadi, “Operations on Single-Valued Neutrosophic Graphs”, Journal of uncertain systems 11 (1) (2017) 1-26. 162 163
3. K. Atanassov, “Intuitionistic fuzzy sets”, Fuzzy Sets Syst. 20 (1986) 87-96. 164

4. S. Boumova, P. Boyvalenkov, and M. Stoyanova, “*Bounds for the Minimum Distance and Covering Radius of Orthogonal Arrays via Their Distance Distributions.*” Preprints 2021, 2021120293 (doi: 10.20944/preprints202112.0293.v1).

165  
166  
167  
168

5. S. Broumi, M. Talea, A. Bakali and F. Smarandache, “*Single-valued neutrosophic graphs*”, Journal of New Theory 10 (2016) 86-101.

169  
170

6. N. Shah, and A. Hussain, “*Neutrosophic soft graphs*”, Neutrosophic Set and Systems 11 (2016) 31-44.

171  
172

7. Henry Garrett, (2022). “*Beyond Neutrosophic Graphs*”, Ohio: E-publishing: Educational Publisher 1091 West 1st Ave Grandview Heights, Ohio 43212 United States. ISBN: 978-1-59973-725-6 (http://fs.unm.edu/BeyondNeutrosophicGraphs.pdf).

173  
174  
175  
176

8. Henry Garrett, “*Co-degree and Degree of classes of Neutrosophic Hypergraphs*”, Preprints 2022, 2022010027 (doi: 10.20944/preprints202201.0027.v1).

177  
178

9. D. Jean, and S. Fault-Tolerant Seo, “*Detection Systems on the King’s Grid.*” Preprints 2022, 2022010249 (doi: 10.20944/preprints202201.0249.v1).

179  
180

10. H. Oliveira, “*Synthesis of Strategic Games With Multiple Pre-set Nash Equilibria - An Artificial Inference Approach Using Fuzzy ASA.*” Preprints 2019, 2019080128 (doi: 10.20944/preprints201908.0128.v2).

181  
182  
183

11. R.E. Perez, F.N. Gonzalez, M.C. Molina, and J.M. Moreno “*Methodological Analysis of Damage Estimation in Hydraulic Infrastructures.*” Preprints 2022, 2022010376 (doi: 10.20944/preprints202201.0376.v1).

184  
185  
186

12. A. Shannon and K.T. Atanassov, “*A first step to a theory of the intuitionistic fuzzy graphs*”, Proceeding of FUBEST (Lakov, D., Ed.) Sofia (1994) 59-61.

187  
188

13. F. Smarandache, “*A Unifying field in logics neutrosophy: Neutrosophic probability, set and logic, Rehoboth:* ” American Research Press (1998).

189  
190

14. F. Tang, X. Zhu, J. Hu, J. Tie, J. Zhou, and Y. Fu, “*Generative Adversarial Unsupervised Image Restoration in Hybrid Degradation Scenes.*” Preprints 2022, 2022020159 (doi: 10.20944/preprints202202.0159.v1).

191  
192  
193

15. H. Wang, F. Smarandache, Y. Zhang, and R. Sunderraman, “*Single-valued neutrosophic sets*”, Multispace and Multistructure 4 (2010) 410-413.

194  
195

16. L. A. Zadeh, “*Fuzzy sets*”, Information and Control 8 (1965) 338-354.

196

17. S. Zuev, Z. Hussain, and P. Kabalyants, “*Nanoparticle Based Water Treatment Model in Squeezing Channel under Magnetic Field.*” Preprints 2021, 2021120295 (doi: 10.20944/preprints202112.0295.v1).

197  
198  
199