

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/359028355>

Failed Independent Number in Neutrosophic Graphs

Preprint · March 2022

DOI: 10.20944/preprints202202.0334.v2

CITATIONS

0

1 author:



Henry Garrett

140 PUBLICATIONS 207 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



On Combinatorics [View project](#)



Neutrosophic Graphs [View project](#)

Failed Independent Number in Neutrosophic Graphs v2

Henry Garrett

Independent Researcher

DrHenryGarrett@gmail.com

Twitter's ID: @DrHenryGarrett | ©DrHenryGarrett.wordpress.com

Abstract

New setting is introduced to study neutrosophic failed-independent number and failed independent neutrosophic-number arising neighborhood of different vertices. Neighbor is a key term to have these notions. Having all possible edges amid vertices in a set is a key type of approach to have these notions namely neutrosophic failed-independent number and failed independent neutrosophic-number. Two numbers are obtained but now both settings leads to approach is on demand which is finding biggest set which have all vertices which are neighbors. Let $NTG : (V, E, \sigma, \mu)$ be a neutrosophic graph. Then failed independent number $\mathcal{I}(NTG)$ for a neutrosophic graph $NTG : (V, E, \sigma, \mu)$ is maximum cardinality of a set S of vertices such that every two vertices of S are endpoints for an edge, simultaneously; failed independent neutrosophic-number $\mathcal{I}_n(NTG)$ for a neutrosophic graph $NTG : (V, E, \sigma, \mu)$ is maximum neutrosophic cardinality of a set S of vertices such that every two vertices of S are endpoints for an edge, simultaneously. As concluding results, there are some statements, remarks, examples and clarifications about some classes of neutrosophic graphs namely path-neutrosophic graphs, cycle-neutrosophic graphs, complete-neutrosophic graphs, star-neutrosophic graphs, complete-bipartite-neutrosophic graphs and complete-t-partite-neutrosophic graphs. The clarifications are also presented in both sections "Setting of Neutrosophic Failed-Independent Number," and "Setting of Failed Independent Neutrosophic-Number," for introduced results and used classes. Neutrosophic number is reused in this way. It's applied to use the type of neutrosophic number in the way that, three values of a vertex are used and they've same share to construct this number to compare with other vertices. Summation of three values of vertex makes one number and applying it to a comparison. This approach facilitates identifying vertices which form neutrosophic failed-independent number and failed independent neutrosophic-number arising neighborhoods of vertices. In path-neutrosophic graphs, two neighbors, form maximal set but with slightly differences, in cycle-neutrosophic graphs, two neighbors forms maximal set. Other classes have same approaches. In complete-neutrosophic graphs, a set of all vertices leads us to neutrosophic failed-independent number and failed independent neutrosophic-number. In star-neutrosophic graphs, a set of vertices containing only center and one other vertex, makes maximal set. In complete-bipartite-neutrosophic graphs, a set of vertices including two vertices from different parts makes intended set but with slightly differences, in complete-t-partite-neutrosophic graphs, a set of t vertices from different parts makes intended set. In both settings, some classes of well-known neutrosophic graphs are studied. Some clarifications for each result and each definition are provided.

Using basic set to extend this set to set of all vertices has key role to have these notions in the form of neutrosophic failed-independent number and failed independent neutrosophic-number arising neighborhood of vertices. The cardinality of a set has eligibility to neutrosophic failed-independent number but the neutrosophic cardinality of a set has eligibility to call failed independent neutrosophic-number. Some results get more frameworks and perspective about these definitions. The way in that, two vertices have connections amid each other, opens the way to do some approaches. A vertex could affect on other vertex but there's no usage of edges. These notions are applied into neutrosophic graphs as individuals but not family of them as drawbacks for these notions. Finding special neutrosophic graphs which are well-known, is an open way to pursue this study. Some problems are proposed to pursue this study. Basic familiarities with graph theory and neutrosophic graph theory are proposed for this article.

Keywords: Neutrosophic Failed-independent Number, Failed independent Neutrosophic-Number, Minimal Set

AMS Subject Classification: 05C17, 05C22, 05E45

1 Background

Fuzzy set in Ref. [16], neutrosophic set in Ref. [3], related definitions of other sets in Refs. [3, 13, 15], graphs and new notions on them in Refs. [1, 4, 8–11, 14, 17], neutrosophic graphs in Ref. [5], studies on neutrosophic graphs in Ref. [2], relevant definitions of other graphs based on fuzzy graphs in Ref. [12], related definitions of other graphs based on neutrosophic graphs in Ref. [6], are proposed. Also, some studies and researches about neutrosophic graphs, are proposed as a book in Ref. [7].

In this section, I use two subsections to illustrate a perspective about the background of this study.

1.1 Motivation and Contributions

In this study, there's an idea which could be considered as a motivation.

Question 1.1. *Is it possible to use mixed versions of ideas concerning “Neutrosophic Failed-Independent Number”, “Failed Independent Neutrosophic-Number” and “Neutrosophic Graph” to define some notions which are applied to neutrosophic graphs?*

It's motivation to find notions to use in any classes of neutrosophic graphs. Real-world applications about time table and scheduling are another thoughts which lead to be considered as motivation. One connection amid two vertices have key roles to assign neutrosophic failed independent number, failed independent neutrosophic-number arising neighborhood of vertices. Thus they're used to define new ideas which conclude to the structure neutrosophic failed independent number and failed independent neutrosophic-number arising neighborhood of vertices. The concept of having edge and extra condition inspire us to study the behavior of vertices in the way that, some types of numbers, neutrosophic failed-independent number, failed independent neutrosophic-number arising neighborhood of vertices are the cases of study in the setting of individuals. In both settings, a corresponded number concludes the discussion. Also, there are some avenues to extend these notions.

The framework of this study is as follows. In the beginning, I introduce basic definitions to clarify about preliminaries. In subsection “Preliminaries”, new notions of neutrosophic failed-independent number, failed independent neutrosophic-number are highlighted, are introduced and are clarified as individuals. In section “Preliminaries”, sets of vertices have the key role in this way. General results are obtained and also, the

results about the basic notions of neutrosophic failed-independent number, failed independent neutrosophic-number are elicited. Some classes of neutrosophic graphs are studied in the terms of neutrosophic failed-independent number, in section “Setting of Neutrosophic Failed-Independent Number,” as individuals. In section “Setting of Failed Independent Neutrosophic-Number,” failed independent neutrosophic-number is applied into individuals. As concluding results, there are some statements, remarks, examples and clarifications about some classes of neutrosophic graphs namely path-neutrosophic graphs, cycle-neutrosophic graphs, complete-neutrosophic graphs, star-neutrosophic graphs, complete-bipartite-neutrosophic graphs and complete-t-partite-neutrosophic graphs. The clarifications are also presented in both sections “Setting of Neutrosophic failed-Independent Number,” and “Setting of Failed Independent Neutrosophic-Number,” for introduced results and used classes. In section “Applications in Time Table and Scheduling”, two applications are posed for quasi-complete and complete notions, namely complete-t-neutrosophic graphs and complete-neutrosophic graphs concerning time table and scheduling when the suspicions are about choosing some subjects and the mentioned models are considered as individual. In section “Open Problems”, some problems and questions for further studies are proposed. In section “Conclusion and Closing Remarks”, gentle discussion about results and applications is featured. In section “Conclusion and Closing Remarks”, a brief overview concerning advantages and limitations of this study alongside conclusions is formed.

1.2 Preliminaries

In this subsection, basic material which is used in this article, is presented. Also, new ideas and their clarifications are elicited.

Basic idea is about the model which is used. First definition introduces basic model.

Definition 1.2. (Graph).

$G = (V, E)$ is called a **graph** if V is a set of objects and E is a subset of $V \times V$ (E is a set of 2-subsets of V) where V is called **vertex set** and E is called **edge set**. Every two vertices have been corresponded to at most one edge.

Neutrosophic graph is the foundation of results in this paper which is defined as follows. Also, some related notions are demonstrated.

Definition 1.3. (Neutrosophic Graph And Its Special Case).

$NTG = (V, E, \sigma = (\sigma_1, \sigma_2, \sigma_3), \mu = (\mu_1, \mu_2, \mu_3))$ is called a **neutrosophic graph** if it's graph, $\sigma_i : V \rightarrow [0, 1]$, and $\mu_i : E \rightarrow [0, 1]$. We add one condition on it and we use **special case** of neutrosophic graph but with same name. The added condition is as follows, for every $v_i v_j \in E$,

$$\mu(v_i v_j) \leq \sigma(v_i) \wedge \sigma(v_j).$$

(i) : σ is called **neutrosophic vertex set**.

(ii) : μ is called **neutrosophic edge set**.

(iii) : $|V|$ is called **order** of NTG and it's denoted by $\mathcal{O}(NTG)$.

(iv) : $\sum_{v \in V} \sigma(v)$ is called **neutrosophic order** of NTG and it's denoted by $\mathcal{O}_n(NTG)$.

(v) : $|E|$ is called **size** of NTG and it's denoted by $\mathcal{S}(NTG)$.

(vi) : $\sum_{e \in E} \sum_{i=1}^3 \mu_i(e)$ is called **neutrosophic size** of NTG and it's denoted by $\mathcal{S}_n(NTG)$.

Some classes of well-known neutrosophic graphs are defined. These classes of neutrosophic graphs are used to form this study and the most results are about them.

Definition 1.4. Let $NTG : (V, E, \sigma, \mu)$ be a neutrosophic graph. Then

(i) : a sequence of vertices $P : x_0, x_1, \dots, x_{\mathcal{O}}$ is called **path** where
 $x_i x_{i+1} \in E, i = 0, 1, \dots, n-1$;

(ii) : **strength** of path $P : x_0, x_1, \dots, x_{\mathcal{O}}$ is $\bigwedge_{i=0, \dots, n-1} \mu(x_i x_{i+1})$;

(iii) : **connectedness** amid vertices x_0 and x_t is

$$\mu^{\infty}(x_0, x_t) = \bigvee_{P: x_0, x_1, \dots, x_t} \bigwedge_{i=0, \dots, t-1} \mu(x_i x_{i+1});$$

(iv) : a sequence of vertices $P : x_0, x_1, \dots, x_{\mathcal{O}}$ is called **cycle** where
 $x_i x_{i+1} \in E, i = 0, 1, \dots, n-1$ and there are two edges xy and uv such that
 $\mu(xy) = \mu(uv) = \bigwedge_{i=0, 1, \dots, n-1} \mu(v_i v_{i+1})$;

(v) : it's **t-partite** where V is partitioned to t parts, $V_1^{s_1}, V_2^{s_2}, \dots, V_t^{s_t}$ and the edge
 xy implies $x \in V_i^{s_i}$ and $y \in V_j^{s_j}$ where $i \neq j$. If it's complete, then it's denoted by
 $K_{\sigma_1, \sigma_2, \dots, \sigma_t}$ where σ_i is σ on $V_i^{s_i}$ instead V which mean $x \notin V_i$ induces $\sigma_i(x) = 0$.
Also, $|V_j^{s_j}| = s_j$;

(vi) : t-partite is **complete bipartite** if $t = 2$, and it's denoted by K_{σ_1, σ_2} ;

(vii) : complete bipartite is **star** if $|V_1| = 1$, and it's denoted by S_{1, σ_2} ;

(viii) : a vertex in V is **center** if the vertex joins to all vertices of a cycle. Then it's
wheel and it's denoted by W_{1, σ_2} ;

(ix) : it's **complete** where $\forall uv \in V, \mu(uv) = \sigma(u) \wedge \sigma(v)$;

(x) : it's **strong** where $\forall uv \in E, \mu(uv) = \sigma(u) \wedge \sigma(v)$.

The natural way proposes us to use the restriction "minimum" instead of
"maximum."

Definition 1.5. (Failed independent Number).

Let $NTG : (V, E, \sigma, \mu)$ be a neutrosophic graph. Then

(i) **failed independent number** $\mathcal{I}(NTG)$ for a neutrosophic graph
 $NTG : (V, E, \sigma, \mu)$ is minimum cardinality of a set S of vertices such that every
two vertices of S are endpoints for an edge, simultaneously;

(ii) **failed independent neutrosophic-number** $\mathcal{I}_n(NTG)$ for a neutrosophic
graph $NTG : (V, E, \sigma, \mu)$ is minimum neutrosophic cardinality of a set S of
vertices such that every two vertices of S are endpoints for an edge,
simultaneously.

For convenient usages, the word neutrosophic which is used in previous definition,
won't be used, usually.

In next part, clarifications about main definition are given. To avoid confusion and
for convenient usages, examples are usually used after every part and names are used in
the way that, abbreviation, simplicity, and summarization are the matters of mind.

Example 1.6. In Figure (2), a complete neutrosophic graph is illustrated. Some points
are represented in follow-up items as follows.




Figure 1. A Neutrosophic Graph in the Viewpoint of its Failed independent Number and its Failed Independent Neutrosophic-Number.

- (i) If $S = \{n_1, n_2\}$ is a set of vertices, then there's no vertex in S but n_1 and n_2 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There's one edge to have exclusive endpoints from S . It implies that $S = \{n_1, n_2\}$ is corresponded to failed independent number $\mathcal{I}(NTG)$ but not failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (ii) if $S = \{n_2, n_4\}$ is a set of vertices, then there's no vertex in S but n_2 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There's one edge to have exclusive endpoints from S . It implies that $S = \{n_2, n_4\}$ is corresponded to failed independent number $\mathcal{I}(NTG)$ but not failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (iii) if $S = \{n_1\}$ is a set of vertices, then there's no vertex in S but n_1 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. Furthermore, There's no edge to have exclusive endpoints from S . But it implies that $S = \{n_1\}$ isn't corresponded to both failed independent number $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (iv) if $S = \{n_3, n_4\}$ is a set of vertices, then there's no vertex in S but n_3 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There's one edge to have exclusive endpoints from S . It implies that $S = \{n_3, n_4\}$ is corresponded to both failed independent number $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (v) 2 is failed independent number and its corresponded sets are $\{n_1, n_2\}$, $\{n_1, n_3\}$, $\{n_1, n_4\}$, $\{n_2, n_3\}$, $\{n_2, n_4\}$, and $\{n_3, n_4\}$;
- (vi) 2.3 is failed independent neutrosophic-number and its corresponded set is $\{n_3, n_4\}$.

But the results are always about the number two where connected model is used. For example,

Proposition 1.7. Let $NTG : (V, E, \sigma, \mu)$ be a complete-neutrosophic graph. Then

$$\mathcal{I}(NTG) = 2.$$

Proof. Suppose $NTG : (V, E, \sigma, \mu)$ is a complete-neutrosophic graph. Every vertex is a neighbor for every given vertex. Assume $|S| = 2$. Then there are x and y in S such that they're endpoints of an edge, simultaneously. If $S = \{n_1, n_2\}$ is a set of vertices, then

there's no vertex in S but n_1 and n_2 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. Furthermore, There's one edge to have exclusive endpoints from S . It implies that $S = \{n_1, n_2\}$ is corresponded to failed independent number $\mathcal{I}(NTG)$. It induces by using the members of S , it's possible to have endpoints of an edge. There's one edge to have exclusive endpoints from S . It implies that $S = \{n_i\}_{|S|=2}$ is corresponded to failed independent number. Thus

$$\mathcal{I}(NTG) = 2.$$

□ 137

Thus we replace the term “minimum” by the term “maximum.” Hence, 138

Definition 1.8. (Failed independent Number). 139

Let $NTG : (V, E, \sigma, \mu)$ be a neutrosophic graph. Then 140

- (i) **failed independent number** $\mathcal{I}(NTG)$ for a neutrosophic graph 141
 $NTG : (V, E, \sigma, \mu)$ is maximum cardinality of a set S of vertices such that every 142
two vertices of S are endpoints for an edge, simultaneously; 143
- (ii) **failed independent neutrosophic-number** $\mathcal{I}_n(NTG)$ for a neutrosophic 144
graph $NTG : (V, E, \sigma, \mu)$ is maximum neutrosophic cardinality of a set S of 145
vertices such that every two vertices of S are endpoints for an edge, 146
simultaneously. 147

For convenient usages, the word neutrosophic which is used in previous definition, 148
won't be used, usually. 149

In next part, clarifications about main definition are given. To avoid confusion and 150
for convenient usages, examples are usually used after every part and names are used in 151
the way that, abbreviation, simplicity, and summarization are the matters of mind. 152

Example 1.9. In Figure (2), a complete neutrosophic graph is illustrated. Some points 153
are represented in follow-up items as follows. 154

- (i) If $S = \{n_1, n_2\}$ is a set of vertices, then there's no vertex in S but n_1 and n_2 . In 155
other side, for having an edge, there's a need to have two vertices. So by using the 156
members of S , it's possible to have endpoints of an edge. There's one edge to have 157
exclusive endpoints from S . But it implies that $S = \{n_1, n_2\}$ isn't corresponded to 158
both of failed independent number $\mathcal{I}(NTG)$ and failed independent 159
neutrosophic-number $\mathcal{I}_n(NTG)$; 160
- (ii) if $S = \{n_2, n_4\}$ is a set of vertices, then there's no vertex in S but n_2 and n_4 . In 161
other side, for having an edge, there's a need to have two vertices. So by using the 162
members of S , it's possible to have endpoints of an edge. There's one edge to have 163
exclusive endpoints from S . But it implies that $S = \{n_2, n_4\}$ isn't corresponded to 164
both of failed independent number $\mathcal{I}(NTG)$ and failed independent 165
neutrosophic-number $\mathcal{I}_n(NTG)$; 166
- (iii) if $S = \{n_1\}$ is a set of vertices, then there's no vertex in S but n_1 . In other side, 167
for having an edge, there's a need to have two vertices. So by using the members 168
of S , it's impossible to have endpoints of an edge. Furthermore, There's no edge 169
to have exclusive endpoints from S . But it implies that $S = \{n_1\}$ isn't 170
corresponded to both of failed independent number $\mathcal{I}(NTG)$ and failed 171
independent neutrosophic-number $\mathcal{I}_n(NTG)$; 172




Figure 2. A Neutrosophic Graph in the Viewpoint of its Failed independent Number and its Failed Independent Neutrosophic-Number.

- (iv) if $S = \{n_1, n_2, n_3, n_4\}$ is a set of vertices, then there's no vertex in S but n_1, n_2, n_3 , and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There are twelve edges to have exclusive endpoints from S . It implies that $S = \{n_1, n_2, n_3, n_4\}$ is corresponded to both failed independent number $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (v) 4 is failed independent number and its corresponded sets is $\{n_1, n_2, n_3, n_4\}$;
- (vi) $\mathcal{O}_n(NTG) = 5.9$ is failed independent neutrosophic-number and its corresponded set is $\{n_3, n_4\}$.

2 Setting of Neutrosophic Failed-Independent Number

In this section, I provide some results in the setting of neutrosophic failed-independent number. Some classes of neutrosophic graphs are chosen. Complete-neutrosophic graph, path-neutrosophic graph, cycle-neutrosophic graph, and star-neutrosophic graph, bipartite-neutrosophic graph, and t-partite-neutrosophic graph, are both of cases of study and classes which the results are about them.

Proposition 2.1. Let $NTG : (V, E, \sigma, \mu)$ be a complete-neutrosophic graph. Then

$$\mathcal{I}(NTG) = \mathcal{O}(NTG).$$

Proof. Suppose $NTG : (V, E, \sigma, \mu)$ is a complete-neutrosophic graph. Every vertex is a neighbor for every given vertex. Assume $|S| > 2$. Then there are x, y and z in S such that they're endpoints of an edge, simultaneously, and they form a triangle. In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There are all possible edges to have exclusive endpoints from S . It implies that $S = \{n_i\}_{|S|=\mathcal{O}(NTG)}$ is corresponded to failed independent number. Thus

$$\mathcal{I}(NTG) = \mathcal{O}(NTG).$$

□

The clarifications about results are in progress as follows. A complete-neutrosophic graph is related to previous result and it's studied to apply the definitions on it. To make it more clear, next part gives one special case to apply definitions and results on




Figure 3. A Neutrosophic Graph in the Viewpoint of its Failed Independent Number.

it. Some items are devised to make more sense about new notions. A complete-neutrosophic graph is related to previous result and it's studied to apply the definitions on it, too.

Example 2.2. In Figure (3), a complete neutrosophic graph is illustrated. Some points are represented in follow-up items as follows.

- (i) If $S = \{n_1, n_2\}$ is a set of vertices, then there's no vertex in S but n_1 and n_2 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There's one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S| \neq \mathcal{O}(NTG)}$. Thus it implies that $S = \{n_1, n_2\}$ isn't corresponded to both of failed independent number $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (ii) if $S = \{n_2, n_4\}$ is a set of vertices, then there's no vertex in S but n_2 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There's one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S| \neq \mathcal{O}(NTG)}$. Thus it implies that $S = \{n_2, n_4\}$ is corresponded to neither failed independent number $\mathcal{I}(NTG)$ nor failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (iii) if $S = \{n_1\}$ is a set of vertices, then there's no vertex in S but n_1 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. Furthermore, There's no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S| \neq \mathcal{O}(NTG)}$. Thus it implies that $S = \{n_1\}$ is corresponded to neither failed independent number $\mathcal{I}(NTG)$ nor failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (iv) if $S = \{n_1, n_2, n_3, n_4\}$ is a set of vertices, then there's no vertex in S but n_1, n_2, n_3 , and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. $S = \{n_i\}_{|S| = \mathcal{O}(NTG)}$. Thus there are twelve edges to have exclusive endpoints from S . It implies that $S = \{n_1, n_2, n_3, n_4\}$ is corresponded to both failed independent number $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (v) 4 is failed independent number and its corresponded sets is $\{n_1, n_2, n_3, n_4\}$;
- (vi) $\mathcal{O}_n(NTG) = 5.9$ is failed independent neutrosophic-number and its corresponded set is $\{n_3, n_4\}$.

Proposition 2.3. Let $NTG : (V, E, \sigma, \mu)$ be a path-neutrosophic graph. Then

$$\mathcal{I}(NTG) = 2.$$

Proof. Suppose $NTG : (V, E, \sigma, \mu)$ is a path-neutrosophic graph. Every vertex isn't a neighbor for every given vertex. If $|S| > 2$, then there are at least three vertices x, y and z such that if x is a neighbor for y and z , then y and z aren't neighbors. Thus there is no triangle but there's one edge. One edge has two endpoints. These endpoints are corresponded to failed independent number $\mathcal{I}(NTG)$. So

$$\mathcal{I}(NTG) = 2.$$

□ 226

Example 2.4. There are two sections for clarifications. 227

(a) In Figure (4), an odd-path-neutrosophic graph is illustrated. Some points are represented in follow-up items as follows. 228 229

- (i) If $S = \{n_2, n_4\}$ is a set of vertices, then there's no vertex in S but n_2 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. There's no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2}$ but it doesn't imply that $S = \{n_2, n_4\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$; 230 231 232 233 234 235
- (ii) if $S = \{n_1, n_3\}$ is a set of vertices, then there's no vertex in S but n_1 and n_3 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. There's no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2}$ but it doesn't imply that $S = \{n_1, n_3\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$; 236 237 238 239 240 241
- (iii) if $S = \{n_1, n_3, n_4, n_5\}$ is a set of vertices, then there's no vertex in S but n_1, n_3, n_4 and n_5 . In other side, for having an edge, there's a need to have two vertices which are consecutive. So by using the members either n_3, n_4 or n_4, n_5 of S , it's possible to have endpoints of an edge either n_3n_4 or n_4n_5 . There are two edges to have exclusive endpoints from S . $S = \{n_i\}_{|S|\neq 2}$ thus it implies that $S = \{n_1, n_3, n_4, n_5\}$ is corresponded to neither failed independent number $\mathcal{I}(NTG)$ nor failed independent neutrosophic-number $\mathcal{I}_n(NTG)$; 242 243 244 245 246 247 248 249
- (iv) if $S = \{n_5, n_6\}$ is a set of vertices, then there's no vertex in S but n_5 and n_6 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There's one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|\neq 2}$ thus it implies that $S = \{n_5, n_6\}$ is corresponded to both failed independent number $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$; 250 251 252 253 254 255
- (v) 2 is failed independent number and its corresponded set is $\{n_1, n_2\}, \{n_1, n_3\}, \{n_1, n_4\}, \{n_1, n_5\}, \{n_2, n_3\}, \{n_2, n_4\}, \{n_2, n_5\}, \{n_3, n_4\}, \{n_3, n_5\}$, and $\{n_4, n_5\}$; 256 257 258
- (vi) 3.2 is failed independent neutrosophic-number and its corresponded set is $\{n_1, n_2\}$. 259 260

(b) In Figure (5), an even-path-neutrosophic graph is illustrated. Some points are represented in follow-up items as follows. 261 262

- (i) If $S = \{n_2, n_4\}$ is a set of vertices, then there's no vertex in S but n_2 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. There's 263 264 265




Figure 4. A Neutrosophic Graph in the Viewpoint of its Failed Independent Number.

no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2}$ but it doesn't imply that $S = \{n_2, n_4\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;

(ii) if $S = \{n_1, n_3\}$ is a set of vertices, then there's no vertex in S but n_1 and n_3 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. There's no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2}$ but it doesn't imply that $S = \{n_1, n_3\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;

(iii) if $S = \{n_1, n_3, n_4, n_5\}$ is a set of vertices, then there's no vertex in S but n_1, n_3, n_4 and n_5 . In other side, for having an edge, there's a need to have two vertices which are consecutive. So by using the members either n_3, n_4 or n_4, n_5 of S , it's possible to have endpoints of an edge either n_3n_4 or n_4n_5 . There are two edges to have exclusive endpoints from S . $S = \{n_i\}_{|S|\neq 2}$ thus it implies that $S = \{n_1, n_3, n_4, n_5\}$ is corresponded to neither failed independent number $\mathcal{I}(NTG)$ nor failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;

(iv) if $S = \{n_5, n_6\}$ is a set of vertices, then there's no vertex in S but n_5 and n_6 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There's one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|\neq 2}$ thus it implies that $S = \{n_5, n_6\}$ is corresponded to both failed independent number $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;

(v) 2 is failed independent number and its corresponded set is $\{n_1, n_2\}, \{n_1, n_3\}, \{n_1, n_4\}, \{n_1, n_5\}, \{n_1, n_6\}, \{n_2, n_3\}, \{n_2, n_4\}, \{n_2, n_5\}, \{n_2, n_6\}, \{n_3, n_4\}, \{n_3, n_5\}, \{n_3, n_6\}, \{n_4, n_5\}, \{n_4, n_6\}$, and $\{n_5, n_6\}$;

(vi) 4.6 is failed independent neutrosophic-number and its corresponded set is $\{n_5, n_6\}$.

Proposition 2.5. Let $NTG : (V, E, \sigma, \mu)$ be a cycle-neutrosophic graph. Then

$$\mathcal{I}(NTG) = 2.$$

Proof. Suppose $NTG : (V, E, \sigma, \mu)$ is a cycle-neutrosophic graph. Every vertex isn't a neighbor for every given vertex. If $|S| > 2$, then there are at least three vertices x, y and z such that if x is a neighbor for y and z , then y and z aren't neighbors. Thus there is




Figure 5. A Neutrosophic Graph in the Viewpoint of its Failed Independent Number.

no triangle but there's one edge. One edge has two endpoints. These endpoints are corresponded to failed independent number $\mathcal{I}(NTG)$. So

$$\mathcal{I}(NTG) = 2.$$

□ 295

The clarifications about results are in progress as follows. An odd-cycle-neutrosophic graph is related to previous result and it's studied to apply the definitions on it. To make it more clear, next part gives one special case to apply definitions and results on it. Some items are devised to make more sense about new notions. An even-cycle-neutrosophic graph is related to previous result and it's studied to apply the definitions on it, too.

Example 2.6. There are two sections for clarifications.

- (a) In Figure (6), an even-cycle-neutrosophic graph is illustrated. Some points are represented in follow-up items as follows.
- (i) If $S = \{n_2, n_4\}$ is a set of vertices, then there's no vertex in S but n_2 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. There's no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2}$ but it doesn't imply that $S = \{n_2, n_4\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
 - (ii) if $S = \{n_1, n_3\}$ is a set of vertices, then there's no vertex in S but n_1 and n_3 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. There's no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2}$ but it doesn't imply that $S = \{n_1, n_3\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
 - (iii) if $S = \{n_1, n_3, n_4, n_5\}$ is a set of vertices, then there's no vertex in S but n_1, n_3, n_4 and n_5 . In other side, for having an edge, there's a need to have two vertices which are consecutive. So by using the members either n_3, n_4 or n_4, n_5 of S , it's possible to have endpoints of an edge either n_3n_4 or n_4n_5 . There are two edges to have exclusive endpoints from S . $S = \{n_i\}_{|S|\neq 2}$ thus it implies that $S = \{n_1, n_3, n_4, n_5\}$ is corresponded to neither failed independent number $\mathcal{I}(NTG)$ nor failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;

- (iv) if $S = \{n_2, n_3\}$ is a set of vertices, then there's no vertex in S but n_2 and n_3 .
In other side, for having an edge, there's a need to have two vertices. So by
using the members of S , it's possible to have endpoints of an edge. There's
one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S| \neq 2}$ thus it implies
that $S = \{n_2, n_3\}$ is corresponded to both failed independent number
 $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (v) 2 is failed independent number and its corresponded set is $\{n_1, n_2\}, \{n_1, n_3\},$
 $\{n_1, n_4\}, \{n_1, n_5\}, \{n_1, n_6\}, \{n_2, n_3\}, \{n_2, n_4\}, \{n_2, n_5\}, \{n_2, n_6\}, \{n_3, n_4\},$
 $\{n_3, n_5\}, \{n_3, n_6\}, \{n_4, n_5\}, \{n_4, n_6\}, \{n_5, n_6\}$, and $\{n_6, n_1\}$;
- (vi) 4 is failed independent neutrosophic-number and its corresponded set is
 $\{n_2, n_3\}$.
- (b) In Figure (7), an odd-cycle-neutrosophic graph is illustrated. Some points are
represented in follow-up items as follows.
- (i) If $S = \{n_2, n_4\}$ is a set of vertices, then there's no vertex in S but n_2 and n_4 .
In other side, for having an edge, there's a need to have two vertices. So by
using the members of S , it's impossible to have endpoints of an edge. There's
no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2}$ but it doesn't
imply that $S = \{n_2, n_4\}$ is corresponded to either failed independent number
 $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (ii) if $S = \{n_1, n_3\}$ is a set of vertices, then there's no vertex in S but n_1 and n_3 .
In other side, for having an edge, there's a need to have two vertices. So by
using the members of S , it's impossible to have endpoints of an edge. There's
no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2}$ but it doesn't
imply that $S = \{n_1, n_3\}$ is corresponded to either failed independent number
 $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (iii) if $S = \{n_1, n_3, n_4, n_5\}$ is a set of vertices, then there's no vertex in S but
 n_1, n_3, n_4 and n_5 . In other side, for having an edge, there's a need to have
two vertices which are consecutive. So by using the members either n_3, n_4 or
 n_4, n_5 of S , it's possible to have endpoints of an edge either n_3n_4 or n_4n_5 .
There are two edges to have exclusive endpoints from S . $S = \{n_i\}_{|S| \neq 2}$ thus
it implies that $S = \{n_1, n_3, n_4, n_5\}$ is corresponded to neither failed
independent number $\mathcal{I}(NTG)$ nor failed independent neutrosophic-number
 $\mathcal{I}_n(NTG)$;
- (iv) if $S = \{n_3, n_4\}$ is a set of vertices, then there's no vertex in S but n_3 and n_4 .
In other side, for having an edge, there's a need to have two vertices. So by
using the members of S , it's possible to have endpoints of an edge. There's
one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S| \neq 2}$ thus it implies
that $S = \{n_3, n_4\}$ is corresponded to both failed independent number
 $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (v) 2 is failed independent number and its corresponded set is $\{n_1, n_2\}, \{n_1, n_3\},$
 $\{n_1, n_4\}, \{n_1, n_5\}, \{n_2, n_3\}, \{n_2, n_4\}, \{n_2, n_5\}, \{n_3, n_4\}, \{n_3, n_5\}, \{n_4, n_5\},$
and $\{n_5, n_1\}$;
- (vi) 4.3 is failed independent neutrosophic-number and its corresponded set is
 $\{n_3, n_4\}$.

Proposition 2.7. Let $NTG : (V, E, \sigma, \mu)$ be a star-neutrosophic graph with center c .
Then

$$\mathcal{I}(NTG) = 2.$$




Figure 6. A Neutrosophic Graph in the Viewpoint of its Failed Independent Number.




Figure 7. A Neutrosophic Graph in the Viewpoint of its Failed Independent Number.

Proof. Suppose $NTG : (V, E, \sigma, \mu)$ is a star-neutrosophic graph. Every vertex isn't a neighbor for every given vertex. Every vertex is a neighbor for center. Furthermore, center is only neighbor for any given vertex. So center is only neighbor for all vertices. Hence all vertices including center and one other vertex are only members of S is a set which its cardinality is failed independent number $\mathcal{I}(NTG)$. In other words, if $|S| > 2$, then there are at least three vertices x, y and z such that if x is a neighbor for y and z , then y and z aren't neighbors and x is center. Thus there is no triangle but there's one edge. One edge has two endpoints which one of them is center. These endpoints are corresponded to failed independent number $\mathcal{I}(NTG)$. So

$$\mathcal{I}(NTG) = 2.$$

□ 369

The clarifications about results are in progress as follows. A star-neutrosophic graph is related to previous result and it's studied to apply the definitions on it. To make it more clear, next part gives one special case to apply definitions and results on it. Some items are devised to make more sense about new notions. A star-neutrosophic graph is related to previous result and it's studied to apply the definitions on it, too. 370 371 372 373 374

Example 2.8. There is one section for clarifications. In Figure (8), a star-neutrosophic graph is illustrated. Some points are represented in follow-up items as follows. 375 376

- (i) If $S = \{n_2, n_4\}$ is a set of vertices, then there's no vertex in S but n_2 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. There's no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2}$ but it doesn't imply that 377 378 379 380




Figure 8. A Neutrosophic Graph in the Viewpoint of its Failed Independent Number.

$S = \{n_2, n_4\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;

(ii) if $S = \{n_3, n_4, n_5\}$ is a set of vertices, then there's no vertex in S but n_3, n_4 and n_5 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. There's no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S| \neq 2}$ thus It doesn't imply that $S = \{n_3, n_5\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;

(iii) if $S = \{n_1, n_3, n_4, n_5\}$ is a set of vertices, then there's no vertex in S but n_1, n_3, n_4 and n_5 . In other side, for having an edge, there's a need to have two vertices which are consecutive. $S = \{n_i\}_{|S| = \mathcal{O}(NTG) - 1}$ so by using the members either n_1, n_3 or n_1, n_4 or n_1, n_5 of S , it's possible to have endpoints of an edge either n_1n_3 or n_1n_4 or n_1n_5 . There are three edges to have exclusive endpoints from S . $S = \{n_i\}_{|S| \neq 2}$ thus it doesn't imply that $S = \{n_1, n_3, n_4, n_5\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;

(iv) if $S = \{n_1, n_4\}$ is a set of vertices, then there's no vertex in S but n_1 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There's one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S| = 2}$ thus it implies that $S = \{n_1, n_4\}$ is corresponded to both of failed independent number $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;

(v) 2 is failed independent number and its corresponded sets are $\{n_1, n_2\}$, $\{n_1, n_3\}$, $\{n_1, n_4\}$, and $\{n_1, n_5\}$;

(vi) 3.7 is failed independent neutrosophic-number and its corresponded set is $\{n_2, n_3, n_4, n_5\}$.

Proposition 2.9. Let $NTG : (V, E, \sigma, \mu)$ be a complete-bipartite-neutrosophic graph. Then

$$\mathcal{I}(NTG) = 2.$$

Proof. Suppose $NTG : (V, E, \sigma, \mu)$ is a complete-bipartite-neutrosophic graph. Every vertex is a neighbor for all vertices in another part. Hence from any part, one vertex is chosen to be only members of S is a set which its cardinality is failed independent number $\mathcal{I}(NTG)$. There are two parts. Thus

$$\mathcal{I}(NTG) = 2.$$

□ 407

The clarifications about results are in progress as follows. A complete-bipartite-neutrosophic graph is related to previous result and it's studied to apply the definitions on it. To make it more clear, next part gives one special case to apply definitions and results on it. Some items are devised to make more senses about new notions. A complete-bipartite-neutrosophic graph is related to previous result and it's studied to apply the definitions on it, too.

Example 2.10. There is one section for clarifications. In Figure (9), a complete-bipartite-neutrosophic graph is illustrated. Some points are represented in follow-up items as follows.

- (i) If $S = \{n_2, n_4\}$ is a set of vertices, then there's no vertex in S but n_2 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints n_2 and n_4 of an edge n_2n_4 . There's one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2=\text{The number of parts.}}$ Thus it implies that $S = \{n_2, n_4\}$ is corresponded to failed independent number $\mathcal{I}(NTG)$ but not failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (ii) if $S = \{n_2, n_3, n_4\}$ is a set of vertices, then there's no vertex in S but n_2, n_3 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints n_2 and n_4 of an edge n_2n_4 . There are two edges to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2=\text{The number of parts.}}$ Thus it doesn't imply that $S = \{n_2, n_3, n_4\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (iii) if $S = \{n_1\}$ is a set of vertices, then there's no vertex in S but n_1 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. There is no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2=\text{The number of parts.}}$ Thus it doesn't imply that $S = \{n_1\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (iv) if $S = \{n_1, n_3\}$ is a set of vertices, then there's no vertex in S but n_1 and n_3 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There is one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2=\text{The number of parts.}}$ Thus it implies that $S = \{n_1, n_3\}$ is corresponded to both failed independent number $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (v) 2 is failed independent number and its corresponded sets are $\{n_1, n_2\}$, $\{n_1, n_3\}$, $\{n_2, n_4\}$, and $\{n_3, n_4\}$;
- (vi) 3.4 is failed independent neutrosophic-number and its corresponded set is $\{n_1, n_3\}$.

Proposition 2.11. Let $NTG : (V, E, \sigma, \mu)$ be a complete- t -partite-neutrosophic graph such that $t \neq 2$. Then

$$\mathcal{I}(NTG) = t.$$

Proof. Suppose $NTG : (V, E, \sigma, \mu)$ is a complete- t -partite-neutrosophic graph. Every vertex is a neighbor for all vertices in another part. Hence from any part, one vertex is chosen to be only members of S is a set which its cardinality is failed independent number $\mathcal{I}(NTG)$. There are t parts. Thus

$$\mathcal{I}(NTG) = t.$$




Figure 9. A Neutrosophic Graph in the Viewpoint of its Failed Independent Number.

The clarifications about results are in progress as follows. A complete-t-partite-neutrosophic graph is related to previous result and it's studied to apply the definitions on it. To make it more clear, next part gives one special case to apply definitions and results on it. Some items are devised to make more sense about new notions. A complete-t-partite-neutrosophic graph is related to previous result and it's studied to apply the definitions on it, too.

Example 2.12. There is one section for clarifications. In Figure (10), a complete-t-partite-neutrosophic graph is illustrated. Some points are represented in follow-up items as follows.

- (i) If $S = \{n_2, n_4\}$ is a set of vertices, then there's no vertex in S but n_2 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints n_2 and n_4 of an edge n_2n_4 . There's one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2=\text{The number of parts.}}$ Thus it implies that $S = \{n_2, n_4\}$ is corresponded to failed independent number $\mathcal{I}(NTG)$ but not failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (ii) if $S = \{n_2, n_3, n_4\}$ is a set of vertices, then there's no vertex in S but n_2, n_3 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints n_2 and n_4 of an edge n_2n_4 . There are two edges to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2=\text{The number of parts.}}$ Thus it doesn't imply that $S = \{n_2, n_3, n_4\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (iii) if $S = \{n_1\}$ is a set of vertices, then there's no vertex in S but n_1 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. There is no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2=\text{The number of parts.}}$ Thus it doesn't imply that $S = \{n_1\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (iv) if $S = \{n_1, n_3\}$ is a set of vertices, then there's no vertex in S but n_1 and n_3 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There is one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2=\text{The number of parts.}}$ Thus it implies




Figure 10. A Neutrosophic Graph in the Viewpoint of its Failed Independent Number.

that $S = \{n_1, n_3\}$ is corresponded to both failed independent number $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;

(v) 2 is failed independent number and its corresponded sets are $\{n_1, n_2\}$, $\{n_1, n_3\}$, $\{n_2, n_4\}$, and $\{n_3, n_4\}$;

(vi) 3.4 is failed independent neutrosophic-number and its corresponded set is $\{n_1, n_3\}$.

3 Setting of failed independent Neutrosophic-Number

In this section, I provide some results in the setting of failed-independent neutrosophic-number. Some classes of neutrosophic graphs are chosen. Complete-neutrosophic graph, path-neutrosophic graph, cycle-neutrosophic graph, and star-neutrosophic graph, bipartite-neutrosophic graph, and t-partite-neutrosophic graph, are both of cases of study and classes which the results are about them.

Proposition 3.1. Let $NTG : (V, E, \sigma, \mu)$ be a complete-neutrosophic graph. Then

$$\mathcal{I}_n(NTG) = \mathcal{O}_n(NTG).$$

Proof. Suppose $NTG : (V, E, \sigma, \mu)$ is a complete-neutrosophic graph. Every vertex is a neighbor for every given vertex. Assume $|S| > 2$. Then there are x, y and z in S such that they're endpoints of an edge, simultaneously, and they form a triangle. In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There are all possible edges to have exclusive endpoints from S . It implies that $S = \{n_i\}_{|S|=\mathcal{O}(NTG)}$ is corresponded to failed independent number. Thus

$$\mathcal{I}_n(NTG) = \mathcal{O}_n(NTG).$$

□ 491

The clarifications about results are in progress as follows. A complete-neutrosophic graph is related to previous result and it's studied to apply the definitions on it. To make it more clear, next part gives one special case to apply definitions and results on it. Some items are devised to make more sense about new notions. A complete-neutrosophic graph is related to previous result and it's studied to apply the definitions on it, too.




Figure 11. A Neutrosophic Graph in the Viewpoint of its Failed Independent Neutrosophic-Number.

Example 3.2. In Figure (11), a complete neutrosophic graph is illustrated. Some points are represented in follow-up items as follows.

- (i) If $S = \{n_1, n_2\}$ is a set of vertices, then there's no vertex in S but n_1 and n_2 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There's one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S| \neq \mathcal{O}(NTG)}$. Thus it implies that $S = \{n_1, n_2\}$ isn't corresponded to both of failed independent number $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (ii) if $S = \{n_2, n_4\}$ is a set of vertices, then there's no vertex in S but n_2 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There's one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S| \neq \mathcal{O}(NTG)}$. Thus it implies that $S = \{n_2, n_4\}$ is corresponded to neither failed independent number $\mathcal{I}(NTG)$ nor failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (iii) if $S = \{n_1\}$ is a set of vertices, then there's no vertex in S but n_1 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. Furthermore, There's no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S| \neq \mathcal{O}(NTG)}$. Thus it implies that $S = \{n_1\}$ is corresponded to neither failed independent number $\mathcal{I}(NTG)$ nor failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (iv) if $S = \{n_1, n_2, n_3, n_4\}$ is a set of vertices, then there's no vertex in S but n_1, n_2, n_3 , and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. $S = \{n_i\}_{|S| = \mathcal{O}(NTG)}$. Thus there are twelve edges to have exclusive endpoints from S . It implies that $S = \{n_1, n_2, n_3, n_4\}$ is corresponded to both failed independent number $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (v) 4 is failed independent number and its corresponded sets is $\{n_1, n_2, n_3, n_4\}$;
- (vi) $\mathcal{O}_n(NTG) = 5.9$ is failed independent neutrosophic-number and its corresponded set is $\{n_3, n_4\}$.

Proposition 3.3. Let $NTG : (V, E, \sigma, \mu)$ be a path-neutrosophic graph. Then

$$\mathcal{I}_n(NTG) = \max\left\{\sum_{i=1}^3 (\sigma_i(x_j) + \sigma_i(x_{j+1}))\right\}_{x_j x_{j+1} \in E}.$$

Proof. Suppose $NTG : (V, E, \sigma, \mu)$ is a path-neutrosophic graph. Every vertex isn't a neighbor for every given vertex. If $|S| > 2$, then there are at least three vertices x, y and z such that if x is a neighbor for y and z , then y and z aren't neighbors. Thus there is no triangle but there's one edge. One edge has two endpoints. These endpoints are corresponded to failed independent number $\mathcal{I}(NTG)$. So

$$\mathcal{I}_n(NTG) = \max\left\{\sum_{i=1}^3 (\sigma_i(x_j) + \sigma_i(x_{j+1}))\right\}_{x_j x_{j+1} \in E}.$$

□ 527

Example 3.4. There are two sections for clarifications. 528

(a) In Figure (12), an odd-path-neutrosophic graph is illustrated. Some points are 529
represented in follow-up items as follows. 530

(i) If $S = \{n_2, n_4\}$ is a set of vertices, then there's no vertex in S but n_2 and n_4 . 531
In other side, for having an edge, there's a need to have two vertices. So by 532
using the members of S , it's impossible to have endpoints of an edge. There's 533
no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2}$ but it doesn't 534
imply that $S = \{n_2, n_4\}$ is corresponded to either failed independent number 535
 $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$; 536

(ii) if $S = \{n_1, n_3\}$ is a set of vertices, then there's no vertex in S but n_1 and n_3 . 537
In other side, for having an edge, there's a need to have two vertices. So by 538
using the members of S , it's impossible to have endpoints of an edge. There's 539
no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2}$ but it doesn't 540
imply that $S = \{n_1, n_3\}$ is corresponded to either failed independent number 541
 $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$; 542

(iii) if $S = \{n_1, n_3, n_4, n_5\}$ is a set of vertices, then there's no vertex in S but 543
 n_1, n_3, n_4 and n_5 . In other side, for having an edge, there's a need to have 544
two vertices which are consecutive. So by using the members either n_3, n_4 or 545
 n_4, n_5 of S , it's possible to have endpoints of an edge either $n_3 n_4$ or $n_4 n_5$. 546
There are two edges to have exclusive endpoints from S . $S = \{n_i\}_{|S| \neq 2}$ thus 547
it implies that $S = \{n_1, n_3, n_4, n_5\}$ is corresponded to neither failed 548
independent number $\mathcal{I}(NTG)$ nor failed independent neutrosophic-number 549
 $\mathcal{I}_n(NTG)$; 550

(iv) if $S = \{n_5, n_6\}$ is a set of vertices, then there's no vertex in S but n_5 and n_6 . 551
In other side, for having an edge, there's a need to have two vertices. So by 552
using the members of S , it's possible to have endpoints of an edge. There's 553
one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S| \neq 2}$ thus it implies 554
that $S = \{n_5, n_6\}$ is corresponded to both failed independent number 555
 $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$; 556

(v) 2 is failed independent number and its corresponded set is $\{n_1, n_2\}, \{n_1, n_3\},$ 557
 $\{n_1, n_4\}, \{n_1, n_5\}, \{n_2, n_3\}, \{n_2, n_4\}, \{n_2, n_5\}, \{n_3, n_4\}, \{n_3, n_5\},$ and 558
 $\{n_4, n_5\}$; 559

(vi) 3.2 is failed independent neutrosophic-number and its corresponded set is 560
 $\{n_1, n_2\}$. 561

(b) In Figure (13), an even-path-neutrosophic graph is illustrated. Some points are 562
represented in follow-up items as follows. 563

(i) If $S = \{n_2, n_4\}$ is a set of vertices, then there's no vertex in S but n_2 and n_4 . 564
In other side, for having an edge, there's a need to have two vertices. So by 565




Figure 12. A Neutrosophic Graph in the Viewpoint of its Failed Independent Neutrosophic-Number.

using the members of S , it's impossible to have endpoints of an edge. There's no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2}$ but it doesn't imply that $S = \{n_2, n_4\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;

(ii) if $S = \{n_1, n_3\}$ is a set of vertices, then there's no vertex in S but n_1 and n_3 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. There's no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2}$ but it doesn't imply that $S = \{n_1, n_3\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;

(iii) if $S = \{n_1, n_3, n_4, n_5\}$ is a set of vertices, then there's no vertex in S but n_1, n_3, n_4 and n_5 . In other side, for having an edge, there's a need to have two vertices which are consecutive. So by using the members either n_3, n_4 or n_4, n_5 of S , it's possible to have endpoints of an edge either n_3n_4 or n_4n_5 . There are two edges to have exclusive endpoints from S . $S = \{n_i\}_{|S|\neq 2}$ thus it implies that $S = \{n_1, n_3, n_4, n_5\}$ is corresponded to neither failed independent number $\mathcal{I}(NTG)$ nor failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;

(iv) if $S = \{n_5, n_6\}$ is a set of vertices, then there's no vertex in S but n_5 and n_6 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There's one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|\neq 2}$ thus it implies that $S = \{n_5, n_6\}$ is corresponded to both failed independent number $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;

(v) 2 is failed independent number and its corresponded set is $\{n_1, n_2\}, \{n_1, n_3\}, \{n_1, n_4\}, \{n_1, n_5\}, \{n_1, n_6\}, \{n_2, n_3\}, \{n_2, n_4\}, \{n_2, n_5\}, \{n_2, n_6\}, \{n_3, n_4\}, \{n_3, n_5\}, \{n_3, n_6\}, \{n_4, n_5\}, \{n_4, n_6\}$, and $\{n_5, n_6\}$;

(vi) 4.6 is failed independent neutrosophic-number and its corresponded set is $\{n_5, n_6\}$.

Proposition 3.5. Let $NTG : (V, E, \sigma, \mu)$ be a cycle-neutrosophic graph. Then

$$\mathcal{I}_n(NTG) = \max\left\{\sum_{i=1}^3 (\sigma_i(x_j) + \sigma_i(x_{j+1}))\right\}_{x_j x_{j+1} \in E}.$$

Proof. Suppose $NTG : (V, E, \sigma, \mu)$ is a cycle-neutrosophic graph. Every vertex isn't a neighbor for every given vertex. If $|S| > 2$, then there are at least three vertices x, y and




Figure 13. A Neutrosophic Graph in the Viewpoint of its Failed Independent Neutrosophic-Number.

z such that if x is a neighbor for y and z , then y and z aren't neighbors. Thus there is no triangle but there's one edge. One edge has two endpoints. These endpoints are corresponded to failed independent number $\mathcal{I}(NTG)$. So

$$\mathcal{I}_n(NTG) = \max\left\{\sum_{i=1}^3 (\sigma_i(x_j) + \sigma_i(x_{j+1}))\right\}_{x_j x_{j+1} \in E}.$$

□ 596

The clarifications about results are in progress as follows. An odd-cycle-neutrosophic graph is related to previous result and it's studied to apply the definitions on it. To make it more clear, next part gives one special case to apply definitions and results on it. Some items are devised to make more sense about new notions. An even-cycle-neutrosophic graph is related to previous result and it's studied to apply the definitions on it, too.

Example 3.6. There are two sections for clarifications.

- (a) In Figure (14), an even-cycle-neutrosophic graph is illustrated. Some points are represented in follow-up items as follows.
- (i) If $S = \{n_2, n_4\}$ is a set of vertices, then there's no vertex in S but n_2 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. There's no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2}$ but it doesn't imply that $S = \{n_2, n_4\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
 - (ii) if $S = \{n_1, n_3\}$ is a set of vertices, then there's no vertex in S but n_1 and n_3 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. There's no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2}$ but it doesn't imply that $S = \{n_1, n_3\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
 - (iii) if $S = \{n_1, n_3, n_4, n_5\}$ is a set of vertices, then there's no vertex in S but n_1, n_3, n_4 and n_5 . In other side, for having an edge, there's a need to have two vertices which are consecutive. So by using the members either n_3, n_4 or n_4, n_5 of S , it's possible to have endpoints of an edge either $n_3 n_4$ or $n_4 n_5$. There are two edges to have exclusive endpoints from S . $S = \{n_i\}_{|S| \neq 2}$ thus

- it implies that $S = \{n_1, n_3, n_4, n_5\}$ is corresponded to neither failed
independent number $\mathcal{I}(NTG)$ nor failed independent neutrosophic-number
 $\mathcal{I}_n(NTG)$;
- (iv) if $S = \{n_2, n_3\}$ is a set of vertices, then there's no vertex in S but n_2 and n_3 .
In other side, for having an edge, there's a need to have two vertices. So by
using the members of S , it's possible to have endpoints of an edge. There's
one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S| \neq 2}$ thus it implies
that $S = \{n_2, n_3\}$ is corresponded to both failed independent number
 $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (v) 2 is failed independent number and its corresponded set is $\{n_1, n_2\}, \{n_1, n_3\},$
 $\{n_1, n_4\}, \{n_1, n_5\}, \{n_1, n_6\}, \{n_2, n_3\}, \{n_2, n_4\}, \{n_2, n_5\}, \{n_2, n_6\}, \{n_3, n_4\},$
 $\{n_3, n_5\}, \{n_3, n_6\}, \{n_4, n_5\}, \{n_4, n_6\}, \{n_5, n_6\},$ and $\{n_6, n_1\}$;
- (vi) 4 is failed independent neutrosophic-number and its corresponded set is
 $\{n_2, n_3\}$.
- (b) In Figure (15), an odd-cycle-neutrosophic graph is illustrated. Some points are
represented in follow-up items as follows.
- (i) If $S = \{n_2, n_4\}$ is a set of vertices, then there's no vertex in S but n_2 and n_4 .
In other side, for having an edge, there's a need to have two vertices. So by
using the members of S , it's impossible to have endpoints of an edge. There's
no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2}$ but it doesn't
imply that $S = \{n_2, n_4\}$ is corresponded to either failed independent number
 $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (ii) if $S = \{n_1, n_3\}$ is a set of vertices, then there's no vertex in S but n_1 and n_3 .
In other side, for having an edge, there's a need to have two vertices. So by
using the members of S , it's impossible to have endpoints of an edge. There's
no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2}$ but it doesn't
imply that $S = \{n_1, n_3\}$ is corresponded to either failed independent number
 $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (iii) if $S = \{n_1, n_3, n_4, n_5\}$ is a set of vertices, then there's no vertex in S but
 n_1, n_3, n_4 and n_5 . In other side, for having an edge, there's a need to have
two vertices which are consecutive. So by using the members either n_3, n_4 or
 n_4, n_5 of S , it's possible to have endpoints of an edge either n_3n_4 or n_4n_5 .
There are two edges to have exclusive endpoints from S . $S = \{n_i\}_{|S| \neq 2}$ thus
it implies that $S = \{n_1, n_3, n_4, n_5\}$ is corresponded to neither failed
independent number $\mathcal{I}(NTG)$ nor failed independent neutrosophic-number
 $\mathcal{I}_n(NTG)$;
- (iv) if $S = \{n_3, n_4\}$ is a set of vertices, then there's no vertex in S but n_3 and n_4 .
In other side, for having an edge, there's a need to have two vertices. So by
using the members of S , it's possible to have endpoints of an edge. There's
one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S| \neq 2}$ thus it implies
that $S = \{n_3, n_4\}$ is corresponded to both failed independent number
 $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (v) 2 is failed independent number and its corresponded set is $\{n_1, n_2\}, \{n_1, n_3\},$
 $\{n_1, n_4\}, \{n_1, n_5\}, \{n_2, n_3\}, \{n_2, n_4\}, \{n_2, n_5\}, \{n_3, n_4\}, \{n_3, n_5\}, \{n_4, n_5\},$
and $\{n_5, n_1\}$;
- (vi) 4.3 is failed independent neutrosophic-number and its corresponded set is
 $\{n_3, n_4\}$.




Figure 14. A Neutrosophic Graph in the Viewpoint of its Failed Independent Neutrosophic-Number.




Figure 15. A Neutrosophic Graph in the Viewpoint of its Failed Independent Neutrosophic-Number.

Proposition 3.7. Let $NTG : (V, E, \sigma, \mu)$ be a star-neutrosophic graph with center c . Then

$$\mathcal{I}_n(NTG) = \sum_{i=1}^3 \sigma_i(c) + \max\left\{\sum_{i=1}^3 \sigma_i(x_j)\right\}.$$

Proof. Suppose $NTG : (V, E, \sigma, \mu)$ is a star-neutrosophic graph. Every vertex isn't a neighbor for every given vertex. Every vertex is a neighbor for center. Furthermore, center is only neighbor for any given vertex. So center is only neighbor for all vertices. Hence all vertices including center and one other vertex are only members of S is a set which its cardinality is failed independent number $\mathcal{I}(NTG)$. In other words, if $|S| > 2$, then there are at least three vertices x, y and z such that if x is a neighbor for y and z , then y and z aren't neighbors and x is center. Thus there is no triangle but there's one edge. One edge has two endpoints which one of them is center. These endpoints are corresponded to failed independent number $\mathcal{I}(NTG)$. So

$$\mathcal{I}_n(NTG) = \sum_{i=1}^3 \sigma_i(c) + \max\left\{\sum_{i=1}^3 \sigma_i(x_j)\right\}.$$

□ 670

The clarifications about results are in progress as follows. A star-neutrosophic graph is related to previous result and it's studied to apply the definitions on it. To make it more clear, next part gives one special case to apply definitions and results on it. Some items are devised to make more sense about new notions. A star-neutrosophic graph is related to previous result and it's studied to apply the definitions on it, too. 671 672 673 674 675

Example 3.8. There is one section for clarifications. In Figure (16), a star-neutrosophic graph is illustrated. Some points are represented in follow-up items as follows. 676 677 678

- (i) If $S = \{n_2, n_4\}$ is a set of vertices, then there's no vertex in S but n_2 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. There's no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2}$ but it doesn't imply that $S = \{n_2, n_4\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$; 679 680 681 682 683 684
- (ii) if $S = \{n_3, n_4, n_5\}$ is a set of vertices, then there's no vertex in S but n_3, n_4 and n_5 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. There's no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|\neq 2}$ thus It doesn't imply that $S = \{n_3, n_5\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$; 685 686 687 688 689 690
- (iii) if $S = \{n_1, n_3, n_4, n_5\}$ is a set of vertices, then there's no vertex in S but n_1, n_3, n_4 and n_5 . In other side, for having an edge, there's a need to have two vertices which are consecutive. $S = \{n_i\}_{|S|=\mathcal{O}(NTG)-1}$ so by using the members either n_1, n_3 or n_1, n_4 or n_1, n_5 of S , it's possible to have endpoints of an edge either n_1n_3 or n_1n_4 or n_1n_5 . There are three edges to have exclusive endpoints from S . $S = \{n_i\}_{|S|\neq 2}$ thus it doesn't imply that $S = \{n_1, n_3, n_4, n_5\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$; 691 692 693 694 695 696 697 698
- (iv) if $S = \{n_1, n_4\}$ is a set of vertices, then there's no vertex in S but n_1 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the 699 700




Figure 16. A Neutrosophic Graph in the Viewpoint of its Failed Independent Neutrosophic-Number.

members of S , it's possible to have endpoints of an edge. There's one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2}$ thus it implies that $S = \{n_1, n_4\}$ is corresponded to both of failed independent number $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;

- (v) 2 is failed independent number and its corresponded sets are $\{n_1, n_2\}$, $\{n_1, n_3\}$, $\{n_1, n_4\}$, and $\{n_1, n_5\}$;
- (vi) 3.7 is failed independent neutrosophic-number and its corresponded set is $\{n_2, n_3, n_4, n_5\}$.

Proposition 3.9. Let $NTG : (V, E, \sigma, \mu)$ be a complete-bipartite-neutrosophic graph. Then

$$\mathcal{I}_n(NTG) = \max\left\{\sum_{i=1}^3 (\sigma_i(x_j) + \sigma_i(x_{j'}))\right\}_{x_j \in V_1, x_{j'} \in V_2}.$$

Proof. Suppose $NTG : (V, E, \sigma, \mu)$ is a complete-bipartite-neutrosophic graph. Every vertex is a neighbor for all vertices in another part. Hence from any part, one vertex is chosen to be only members of S is a set which its cardinality is failed independent number $\mathcal{I}(NTG)$. There are two parts. Thus

$$\mathcal{I}_n(NTG) = \max\left\{\sum_{i=1}^3 (\sigma_i(x_j) + \sigma_i(x_{j'}))\right\}_{x_j \in V_1, x_{j'} \in V_2}.$$

□ 709

The clarifications about results are in progress as follows. A complete-bipartite-neutrosophic graph is related to previous result and it's studied to apply the definitions on it. To make it more clear, next part gives one special case to apply definitions and results on it. Some items are devised to make more senses about new notions. A complete-bipartite-neutrosophic graph is related to previous result and it's studied to apply the definitions on it, too.

Example 3.10. There is one section for clarifications. In Figure (17), a complete-bipartite-neutrosophic graph is illustrated. Some points are represented in follow-up items as follows.

- (i) If $S = \{n_2, n_4\}$ is a set of vertices, then there's no vertex in S but n_2 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the




Figure 17. A Neutrosophic Graph in the Viewpoint of its Failed Independent Neutrosophic-Number.

members of S , it's possible to have endpoints n_2 and n_4 of an edge n_2n_4 . There's one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2} = \text{The number of parts.}$ Thus it implies that $S = \{n_2, n_4\}$ is corresponded to failed independent number $\mathcal{I}(NTG)$ but not failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;

(ii) if $S = \{n_2, n_3, n_4\}$ is a set of vertices, then there's no vertex in S but n_2, n_3 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints n_2 and n_4 of an edge n_2n_4 . There are two edges to have exclusive endpoints from S .

$S = \{n_i\}_{|S|=2} = \text{The number of parts.}$ Thus it doesn't imply that $S = \{n_2, n_3, n_4\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;

(iii) if $S = \{n_1\}$ is a set of vertices, then there's no vertex in S but n_1 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. There is no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2} = \text{The number of parts.}$ Thus it doesn't imply that $S = \{n_1\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;

(iv) if $S = \{n_1, n_3\}$ is a set of vertices, then there's no vertex in S but n_1 and n_3 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There is one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2} = \text{The number of parts.}$ Thus it implies that $S = \{n_1, n_3\}$ is corresponded to both failed independent number $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;

(v) 2 is failed independent number and its corresponded sets are $\{n_1, n_2\}$, $\{n_1, n_3\}$, $\{n_2, n_4\}$, and $\{n_3, n_4\}$;

(vi) 3.4 is failed independent neutrosophic-number and its corresponded set is $\{n_1, n_3\}$.

Proposition 3.11. Let $NTG : (V, E, \sigma, \mu)$ be a complete- t -partite-neutrosophic graph such that $t \neq 2$. Then

$$\mathcal{I}_n(NTG) = \max \left\{ \sum_{i=1}^3 (\sigma_i(x_{j_1}) + \sigma_i(x_{j_2}) + \cdots + \sigma_i(x_{j_t})) \right\}_{x_{j_1} \in V_1, x_{j_2} \in V_2, \dots, x_{j_t} \in V_t}.$$

Proof. Suppose $NTG : (V, E, \sigma, \mu)$ is a complete-t-partite-neutrosophic graph. Every vertex is a neighbor for all vertices in another part. Hence from any part, one vertex is chosen to be only members of S is a set which its cardinality is failed independent number $\mathcal{I}(NTG)$. There are t parts. Thus

$$\mathcal{I}_n(NTG) = \max \left\{ \sum_{i=1}^3 (\sigma_i(x_{j_1}) + \sigma_i(x_{j_2}) + \cdots + \sigma_i(x_{j_t})) \right\}_{x_{j_1} \in V_1, x_{j_2} \in V_2, \dots, x_{j_t} \in V_t}.$$

□ 747

The clarifications about results are in progress as follows. A complete-t-partite-neutrosophic graph is related to previous result and it's studied to apply the definitions on it. To make it more clear, next part gives one special case to apply definitions and results on it. Some items are devised to make more sense about new notions. A complete-t-partite-neutrosophic graph is related to previous result and it's studied to apply the definitions on it, too.

Example 3.12. There is one section for clarifications. In Figure (18), a complete-t-partite-neutrosophic graph is illustrated. Some points are represented in follow-up items as follows.

- (i) If $S = \{n_2, n_4\}$ is a set of vertices, then there's no vertex in S but n_2 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints n_2 and n_4 of an edge n_2n_4 . There's one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2=\text{The number of parts.}}$. Thus it implies that $S = \{n_2, n_4\}$ is corresponded to failed independent number $\mathcal{I}(NTG)$ but not failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (ii) if $S = \{n_2, n_3, n_4\}$ is a set of vertices, then there's no vertex in S but n_2, n_3 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints n_2 and n_4 of an edge n_2n_4 . There are two edges to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2=\text{The number of parts.}}$. Thus it doesn't imply that $S = \{n_2, n_3, n_4\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (iii) if $S = \{n_1\}$ is a set of vertices, then there's no vertex in S but n_1 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. There is no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2=\text{The number of parts.}}$. Thus it doesn't imply that $S = \{n_1\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (iv) if $S = \{n_1, n_3\}$ is a set of vertices, then there's no vertex in S but n_1 and n_3 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There is one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2=\text{The number of parts.}}$. Thus it implies that $S = \{n_1, n_3\}$ is corresponded to both failed independent number $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (v) 2 is failed independent number and its corresponded sets are $\{n_1, n_2\}$, $\{n_1, n_3\}$, $\{n_2, n_4\}$, and $\{n_3, n_4\}$;
- (vi) 3.4 is failed independent neutrosophic-number and its corresponded set is $\{n_1, n_3\}$.




Figure 18. A Neutrosophic Graph in the Viewpoint of its Failed Independent Neutrosophic-Number.

4 Applications in Time Table and Scheduling

In this section, two applications for time table and scheduling are provided where the models are either complete models which mean complete connections are formed as individual and family of complete models with common neutrosophic vertex set or quasi-complete models which mean quasi-complete connections are formed as individual and family of quasi-complete models with common neutrosophic vertex set.

Designing the programs to achieve some goals is general approach to apply on some issues to function properly. Separation has key role in the context of this style. Separating the duration of work which are consecutive, is the matter and it has importance to avoid mixing up.

Step 1. (Definition) Time table is an approach to get some attributes to do the work fast and proper. The style of scheduling implies special attention to the tasks which are consecutive.

Step 2. (Issue) Scheduling of program has faced with difficulties to differ amid consecutive sections. Beyond that, sometimes sections are not the same.

Step 3. (Model) The situation is designed as a model. The model uses data to assign every section and to assign to relation amid sections, three numbers belong unit interval to state indeterminacy, possibilities and determinacy. There's one restriction in that, the numbers amid two sections are at least the number of the relations amid them. Table (1), clarifies about the assigned numbers to these situations.

Table 1. Scheduling concerns its Subjects and its Connections as a neutrosophic graph in a Model.

Sections of <i>NTG</i>	n_1	$n_2 \cdots$	n_5
Values	$(0.7, 0.9, 0.3)$	$(0.4, 0.2, 0.8) \cdots$	$(0.4, 0.2, 0.8)$
Connections of <i>NTG</i>	E_1	$E_2 \cdots$	E_6
Values	$(0.4, 0.2, 0.3)$	$(0.5, 0.2, 0.3) \cdots$	$(0.3, 0.2, 0.3)$




Figure 19. A Neutrosophic Graph in the Viewpoint of its Failed Independent Number and its Failed Independent Neutrosophic-Number.

4.1 Case 1: Complete-t-partite Model alongside its failed independent Number and its failed independent Neutrosophic-Number

Step 4. (Solution) The neutrosophic graph alongside its failed independent number and its failed independent neutrosophic-number as model, propose to use specific number. Every subject has connection with some subjects. Thus the connection is applied as possible and the model demonstrates quasi-full connections as quasi-possible. Using the notion of strong on the connection amid subjects, causes the importance of subject goes in the highest level such that the value amid two consecutive subjects, is determined by those subjects. If the configuration is star, the number is different. Also, it holds for other types such that complete, wheel, path, and cycle. The collection of situations is another application of failed independent number and its failed independent neutrosophic-number when the notion of family is applied in the way that all members of family are from same classes of neutrosophic graphs. As follows, There are five subjects which are represented as Figure (19). This model is strong and even more it's quasi-complete. And the study proposes using specific number which is called failed independent number and failed independent neutrosophic-number. There are also some analyses on other numbers in the way that, the clarification is gained about being special number or not. Also, in the last part, there is one neutrosophic number to assign to this model and situation to compare them with same situations to get more precise. Consider Figure (19). In Figure (19), an complete-t-partite-neutrosophic graph is illustrated. Some points are represented in follow-up items as follows.

- (i) If $S = \{n_2, n_4\}$ is a set of vertices, then there's no vertex in S but n_2 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints n_2 and n_4 of an edge n_2n_4 . There's one edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2}$ = The number of parts. Thus it implies that $S = \{n_2, n_4\}$ is corresponded to failed independent number $\mathcal{I}(NTG)$ but not failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (ii) if $S = \{n_2, n_3, n_4\}$ is a set of vertices, then there's no vertex in S but n_2, n_3 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints n_2 and n_4 of




Figure 20. A Neutrosophic Graph in the Viewpoint of its Failed Independent Number and its Failed Independent Neutrosophic-Number.

- an edge n_2n_4 . There are two edges to have exclusive endpoints from S .
 $S = \{n_i\}_{|S|=2}$ = The number of parts. Thus it doesn't imply that
 $S = \{n_2, n_3, n_4\}$ is corresponded to either failed independent number
 $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (iii) if $S = \{n_1\}$ is a set of vertices, then there's no vertex in S but n_1 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. There is no edge to have exclusive endpoints from S . $S = \{n_i\}_{|S|=2}$ = The number of parts. Thus it doesn't imply that $S = \{n_1\}$ is corresponded to either failed independent number $\mathcal{I}(NTG)$ or failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (iv) if $S = \{n_1, n_3\}$ is a set of vertices, then there's no vertex in S but n_1 and n_3 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There is one edge to have exclusive endpoints from S .
 $S = \{n_i\}_{|S|=2}$ = The number of parts. Thus it implies that $S = \{n_1, n_3\}$ is corresponded to both failed independent number $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (v) 2 is failed independent number and its corresponded sets are $\{n_1, n_2\}$, $\{n_1, n_3\}$, $\{n_2, n_4\}$, and $\{n_3, n_4\}$;
- (vi) 3.4 is failed independent neutrosophic-number and its corresponded set is $\{n_1, n_3\}$.

4.2 Case 2: Complete Model alongside its A Neutrosophic Graph in the Viewpoint of its failed independent Number and its failed independent Neutrosophic-Number.

Step 4. (Solution) The neutrosophic graph alongside its failed independent number and its failed independent neutrosophic-number as model, propose to use specific number. Every subject has connection with every given subject in deemed way. Thus the connection applied as possible and the model demonstrates full connections as possible between parts but with different view where symmetry amid vertices and edges are the matters. Using the notion of strong on the connection amid subjects, causes the importance of subject goes in the highest level such that the value amid two consecutive subjects, is determined by those subjects. If the configuration is complete multipartite, the number is different. Also, it holds for other types such that star, wheel, path, and cycle. The collection of situations is another application of failed independent number and failed

independent neutrosophic-number when the notion of family is applied in the way that all members of family are from same classes of neutrosophic graphs. As follows, There are four subjects which are represented in the formation of one model as Figure (20). This model is neutrosophic strong as individual and even more it's complete. And the study proposes using specific number which is called failed independent number and failed independent neutrosophic-number for this model. There are also some analyses on other numbers in the way that, the clarification is gained about being special number or not. Also, in the last part, there is one neutrosophic number to assign to these models as individual. A model as a collection of situations to compare them with another model as a collection of situations to get more precise. Consider Figure (20). There is one section for clarifications.

- (i) If $S = \{n_1, n_2\}$ is a set of vertices, then there's no vertex in S but n_1 and n_2 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There's one edge to have exclusive endpoints from S . But it implies that $S = \{n_1, n_2\}$ isn't corresponded to both of failed independent number $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (ii) if $S = \{n_2, n_4\}$ is a set of vertices, then there's no vertex in S but n_2 and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There's one edge to have exclusive endpoints from S . But it implies that $S = \{n_2, n_4\}$ isn't corresponded to both of failed independent number $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (iii) if $S = \{n_1\}$ is a set of vertices, then there's no vertex in S but n_1 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's impossible to have endpoints of an edge. Furthermore, There's no edge to have exclusive endpoints from S . But it implies that $S = \{n_1\}$ isn't corresponded to both of failed independent number $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (iv) if $S = \{n_1, n_2, n_3, n_4\}$ is a set of vertices, then there's no vertex in S but n_1, n_2, n_3 , and n_4 . In other side, for having an edge, there's a need to have two vertices. So by using the members of S , it's possible to have endpoints of an edge. There are twelve edges to have exclusive endpoints from S . It implies that $S = \{n_1, n_2, n_3, n_4\}$ is corresponded to both failed independent number $\mathcal{I}(NTG)$ and failed independent neutrosophic-number $\mathcal{I}_n(NTG)$;
- (v) 4 is failed independent number and its corresponded sets is $\{n_1, n_2, n_3, n_4\}$;
- (vi) $\mathcal{O}_n(NTG) = 5.9$ is failed independent neutrosophic-number and its corresponded set is $\{n_3, n_4\}$.

5 Open Problems

In this section, some questions and problems are proposed to give some avenues to pursue this study. The structures of the definitions and results give some ideas to make new settings which are eligible to extend and to create new study.

Notion concerning failed independent number and failed independent neutrosophic-number are defined in neutrosophic graphs. Neutrosophic number is also reused. Thus,

Question 5.1. *Is it possible to use other types of failed independent number and failed independent neutrosophic-number?*

- Question 5.2.** Are existed some connections amid different types of failed independent number and failed independent neutrosophic-number in neutrosophic graphs?
- Question 5.3.** Is it possible to construct some classes of neutrosophic graphs which have “nice” behavior?
- Question 5.4.** Which mathematical notions do make an independent study to apply these types in neutrosophic graphs?
- Problem 5.5.** Which parameters are related to this parameter?
- Problem 5.6.** Which approaches do work to construct applications to create independent study?
- Problem 5.7.** Which approaches do work to construct definitions which use all definitions and the relations amid them instead of separate definitions to create independent study?

6 Conclusion and Closing Remarks

In this section, concluding remarks and closing remarks are represented. The drawbacks of this article are illustrated. Some benefits and advantages of this study are highlighted. This study uses two definitions concerning failed independent number and failed independent neutrosophic-number arising neighborhoods of vertices to study neutrosophic graphs. New neutrosophic number is reused which is too close to the notion of neutrosophic number but it’s different since it uses all values as type-summation on them. Comparisons amid number and edges are done by using neutrosophic tool. The connections of vertices which aren’t clarified by one edge differ them from each other and put them in different categories to represent a number which

Table 2. A Brief Overview about Advantages and Limitations of this study

Advantages	Limitations
1. Neutrosophic Failed Independent Number	1. Wheel-Neutrosophic Graphs
2. Failed Independent Neutrosophic-Number	
3. Neutrosophic Number	2. Study on Families
4. Study on Classes of Neutrosophic Graphs	
5. Using Neighborhood of Vertices	3. Same Models in Family

is called failed independent number and failed independent neutrosophic-number. Further studies could be about changes in the settings to compare these notions amid different settings of neutrosophic graphs theory. One way is finding some relations amid all definitions of notions to make sensible definitions. In Table (2), some limitations and advantages of this study are pointed out.

References

1. N.H. Abughazalah, and M. Khan, “Selection of Optimum Nonlinear Confusion Component of Information Confidentiality Mechanism Using Grey Theory Based Decision-Making Technique.” Preprints 2021, 2021120310 (doi: 10.20944/preprints202112.0310.v1).

2. M. Akram, and G. Shahzadi, “Operations on Single-Valued Neutrosophic Graphs”, Journal of uncertain systems 11 (1) (2017) 1-26.

955

956

3. K. Atanassov, “Intuitionistic fuzzy sets”, Fuzzy Sets Syst. 20 (1986) 87-96.

957

4. S. Boumova, P. Boyvalenkov, and M. Stoyanova, “Bounds for the Minimum Distance and Covering Radius of Orthogonal Arrays via Their Distance Distributions.” Preprints 2021, 2021120293 (doi: 10.20944/preprints202112.0293.v1).

958

959

960

961

5. S. Broumi, M. Talea, A. Bakali and F. Smarandache, “Single-valued neutrosophic graphs”, Journal of New Theory 10 (2016) 86-101.

962

963

6. N. Shah, and A. Hussain, “Neutrosophic soft graphs”, Neutrosophic Set and Systems 11 (2016) 31-44.

964

965

7. Henry Garrett, (2022). “Beyond Neutrosophic Graphs”, Ohio: E-publishing: Educational Publisher 1091 West 1st Ave Grandview Heights, Ohio 43212 United States. ISBN: 978-1-59973-725-6 (http://fs.unm.edu/BeyondNeutrosophicGraphs.pdf).

966

967

968

969

8. Henry Garrett, “Co-degree and Degree of classes of Neutrosophic Hypergraphs”, Preprints 2022, 2022010027 (doi: 10.20944/preprints202201.0027.v1).

970

971

9. D. Jean, and S. Fault-Tolerant Seo, “Detection Systems on the King’s Grid.” Preprints 2022, 2022010249 (doi: 10.20944/preprints202201.0249.v1).

972

973

10. H. Oliveira, “Synthesis of Strategic Games With Multiple Pre-set Nash Equilibria - An Artificial Inference Approach Using Fuzzy ASA.” Preprints 2019, 2019080128 (doi: 10.20944/preprints201908.0128.v2).

974

975

976

11. R.E. Perez, F.N. Gonzalez, M.C. Molina, and J.M. Moreno “Methodological Analysis of Damage Estimation in Hydraulic Infrastructures.” Preprints 2022, 2022010376 (doi: 10.20944/preprints202201.0376.v1).

977

978

979

12. A. Shannon and K.T. Atanassov, “A first step to a theory of the intuitionistic fuzzy graphs”, Proceeding of FUBEST (Lakov, D., Ed.) Sofia (1994) 59-61.

980

981

13. F. Smarandache, “A Unifying field in logics neutrosophy: Neutrosophic probability, set and logic, Rehoboth: ” American Research Press (1998).

982

983

14. F. Tang, X. Zhu, J. Hu, J. Tie, J. Zhou, and Y. Fu, “Generative Adversarial Unsupervised Image Restoration in Hybrid Degradation Scenes.” Preprints 2022, 2022020159 (doi: 10.20944/preprints202202.0159.v1).

984

985

986

15. H. Wang, F. Smarandache, Y. Zhang, and R. Sunderraman, “Single-valued neutrosophic sets”, Multispace and Multistructure 4 (2010) 410-413.

987

988

16. L. A. Zadeh, “Fuzzy sets”, Information and Control 8 (1965) 338-354.

989

17. S. Zuev, Z. Hussain, and P. Kabalyants, “Nanoparticle Based Water Treatment Model in Squeezing Channel under Magnetic Field.” Preprints 2021, 2021120295 (doi: 10.20944/preprints202112.0295.v1).

990

991

992

33/33

View publication stats