

## Binding Number of Some Special Classes of Trees

B.Chaluvaraju<sup>1</sup>, H.S.Boregowda<sup>2</sup> and S.Kumbinarsaiah<sup>3</sup>

<sup>1</sup>Department of Mathematics, Bangalore University, Janana Bharathi Campus, Bangalore-560 056, India

<sup>2</sup>Department of Studies and Research in Mathematics, Tumkur University, Tumkur-572 103, India

<sup>3</sup>Department of Mathematics, Karnatak University, Dharwad-580 003, India

E-mail: bchaluvraju@gmail.com, bgsamarasa@gmail.com, kumbinarsaiah@gmail.com

**Abstract:** The binding number of a graph  $G = (V, E)$  is defined to be the minimum of  $|N(X)|/|X|$  taken over all nonempty set  $X \subseteq V(G)$  such that  $N(X) \neq V(G)$ . In this article, we explore the properties and bounds on binding number of some special classes of trees.

**Key Words:** Graph, tree, realizing set, binding number, Smarandachely binding number.

**AMS(2010):** 05C05.

### §1. Introduction

In this article, we consider finite, undirected, simple and connected graphs  $G = (V, E)$  with vertex set  $V$  and edge set  $E$ . As such  $n = |V|$  and  $m = |E|$  denote the number of vertices and edges of a graph  $G$ , respectively. An edge - induced subgraph is a subset of the edges of a graph  $G$  together with any vertices that are their endpoints. In general, we use  $\langle X \rangle$  to denote the subgraph induced by the set of edges  $X \subseteq E$ . A graph  $G$  is connected if it has a  $u - v$  path whenever  $u, v \in V(G)$  (otherwise,  $G$  is disconnected). The open neighborhood of a vertex  $v \in V(G)$  is  $N(v) = \{u \in V : uv \in E(G)\}$  and the closed neighborhood  $N[v] = N(v) \cup \{v\}$ . The degree of  $v$ , denoted by  $\deg(v)$ , is the cardinality of its open neighborhood. A vertex with degree one in a graph  $G$  is called pendant or a leaf or an end-vertex, and its neighbor is called its support or cut vertex. An edge incident to a leaf in a graph  $G$  is called a pendant edge. A graph with no cycle is acyclic. A tree  $T$  is a connected acyclic graph. Unless mentioned otherwise, for terminology and notation the reader may refer Harary [3].

Woodall [7] defined the binding number of  $G$  as follows: If  $X \subseteq V(G)$ , then the open neighborhood of the set  $X$  is defined as  $N(X) = \bigcup_{x \in X} N(x)$ . The binding number of  $G$ , denoted  $b(G)$ , is given by

$$b(G) = \min_{X \in F} \frac{|N(X)|}{|X|},$$

where  $F = \{X \subseteq V(G) : X \neq \emptyset, N(X) \neq V(G)\}$ . We say that  $b(G)$  is realized on a set  $X$  if  $X \in F$  and  $b(G) = \frac{|N(X)|}{|X|}$ , and the set  $X$  is called a realizing set for  $b(G)$ . Generally, for a given graph  $H$ , a *Smarandachely binding number*  $b_H(G)$  is the minimum number  $b(G)$  on such  $F$  with

---

<sup>1</sup>Received July 23, 2015, Accepted February 18, 2016.

$\langle X \rangle_G \not\cong H$  for  $\forall X \in F$ . Clearly, if  $H$  is not a spanning subgraph of  $G$ , then  $b_H(G) = b(G)$ .

For complete review and the following existing results on the binding number and its related concepts, we follow [1], [2], [5] and [6].

**Theorem 1.1** *For any path  $P_n$  with  $n \geq 2$  vertices,*

$$b(P_n) = \begin{cases} 1 & \text{if } n \text{ is even;} \\ \frac{n-1}{n+1} & \text{if } n \text{ is odd.} \end{cases}$$

**Theorem 1.2** *For any spanning subgraph  $H$  of a graph  $G$ ,  $b(G) \leq b(H)$ .*

In [8], Wayne Goddard established several bounds including ones linking the binding number of a tree to the distribution of its end-vertices  $end(G) = \{v \in V(G) : deg(v) = 1\}$ . Also, let  $\varrho(v) = |N(v) \cap end(G)|$  and  $\varrho(G) = \max \{\varrho(v) : v \in V(G)\}$ . The following result is obviously true if  $\varrho(G) = 0$  and if  $\varrho(G) = 1$ , follows from taking  $X = \{N(v) \cap end(G)\}$ , where  $v$  is a vertex for which  $\varrho(v) = \varrho(G)$ .

**Theorem 1.3** *For any graph  $G$ ,  $\varrho(G).b(G) \leq 1$ .*

**Theorem 1.4** *For any nontrivial tree  $T$ ,*

- (1)  $b(T) \geq 1/\Delta(T)$ ;
- (2)  $b(T) \geq 1/\varrho(T) + 1$ .

## §2. Main Results

**Observation 2.1** Let  $T$  be a tree with  $n \geq 3$  vertices, having  $(n-1)$ -pendant vertices, which are connected to unique vertex. Then  $b(T)$  is the reciprocal of number of vertices connected to unique vertex.

**Observation 2.2** Let  $T$  be a nontrivial tree. Then  $b(T) > 0$ .

**Observation 2.3** Let  $T$  be a tree with  $b(T) < 1$ . Then every realizing set of  $T$  is independent.

**Theorem 2.4** *For any Star  $K_{1,n-1}$  with  $n \geq 2$  vertices,*

$$b(K_{1,n-1}) = \frac{1}{n-1}.$$

*Proof* Let  $K_{1,n-1}$  be a star with  $n \geq 2$  vertices. If  $K_{1,n-1}$  has  $\{v_1, v_2, \dots, v_n\}$  vertices with  $deg(v_1) = n-1$  and  $deg(v_2) = deg(v_3) = \dots = deg(v_n) = 1$ . We prove the result by induction on  $n$ . For  $n = 2$ , then  $|N(X)| = |X| = 1$  and  $b(K_{1,1}) = 1$ . For  $n = 3$ ,  $|N(X)| < |X| = 2$  and  $b(K_{1,2}) = \frac{1}{2}$ . Let us assume the result is true for  $n = k$  for some  $k$ , where  $k$  is a positive integer. Hence  $b(K_{1,k-1}) = \frac{1}{k-1}$ .

Now we shall show that the result is true for  $n > k$ . Since  $(k + 1)$ - pendant vertices in  $K_{1,k+1}$  are connected to the unique vertex  $v_1$ . Here newly added vertex  $v_{k+1}$  must be adjacent to  $v_1$  only. Otherwise  $K_{1,k+1}$  loses its star criteria and  $v_{k+1}$  is not adjacent to  $\{v_2, v_3, \dots, v_k\}$ , then  $K_{1,k+1}$  has  $k$  number of pendant vertices connected to vertex  $v_1$ . Therefore by Observation 2.1, the desired result follows.  $\square$

**Theorem 2.5** *Let  $T_1$  and  $T_2$  be two stars with order  $n_1$  and  $n_2$ , respectively. Then  $n_1 < n_2$  if and only if  $b(T_1) > b(T_2)$ .*

*Proof* By Observation 2.1 and Theorem 2.4, we have  $b(T_1) = \frac{1}{n_1}$  and  $b(T_2) = \frac{1}{n_2}$ . Due to the fact of  $n_1 < n_2$  if and only if  $\frac{1}{n_1} > \frac{1}{n_2}$ . Thus the result follows.  $\square$

**Definition 2.6** *The double star  $K_{r,s}^*$  is a tree with diameter 3 and central vertices of degree  $r$  and  $s$  respectively, where the diameter of graph is the length of the shortest path between the most distanced vertices.*

**Theorem 2.7** *For any double star  $K_{r,s}^*$  with  $1 \leq r \leq s$  vertices,*

$$b(K_{r,s}^*) = \frac{1}{\max\{r, s\} - 1}.$$

*Proof* Suppose  $K_{r,s}^*$  is a double star with  $1 \leq r \leq s$  vertices. Then there exist exactly two central vertices  $x$  and  $y$  for all  $x, y \in V(K_{r,s}^*)$  such that the degree of  $x$  and  $y$  are  $r$  and  $s$  respectively. By definition, the double star  $K_{r,s}^*$  is a tree with diameter 3 having only one edge between  $x$  and  $y$ . Therefore the vertex  $x$  is adjacent to  $(r - 1)$ -pendant vertices and the vertex  $y$  is adjacent to  $(s - 1)$ -pendant vertices.

Clearly  $\max\{r - 1, s - 1\}$  pendant vertices are adjacent to a unique vertex  $x$  or  $y$  as the case may be. Therefore  $b(K_{r,s}^*) = \frac{1}{\max\{r-1, s-1\}}$ . Hence the result follows.  $\square$

**Definition 2.8** *A subdivided star, denoted  $K_{1,n-1}^*$  is a star  $K_{1,n-1}$  whose edges are subdivided once, that is each edge is replaced by a path of length 2 by adding a vertex of degree 2.*

**Observation 2.9** Let  $K_{1,n-1}$  be a star with  $n \geq 2$  vertices. Then cardinality of the vertex set of  $K_{1,n-1}^*$  is  $p = 2n - 1$ .

**Theorem 2.10** *For any subdivided star  $K_{1,n-1}^*$  with  $n \geq 2$  vertices,*

$$b(K_{1,n-1}^*) = \begin{cases} \frac{1}{2} & \text{if } n = 2; \\ \frac{2}{3} & \text{if } n = 3; \\ 1 & \text{otherwise.} \end{cases}$$

*Proof* By Observation 2.9, the subdivided star  $K_{1,n-1}^*$  has  $p = 2n - 1$  vertices. Then the following cases arise:

**Case 1.** If  $n = 2$ , then by Theorem 1.1,  $b(K_{1,2-1}^*) = b(P_3) = \frac{1}{2}$ .

**Case 2.** If  $n = 3$ , then by Theorem 1.1,  $b(K_{1,3-1}^*) = b(P_5) = \frac{2}{3}$ .

**Case 3.** If a vertex  $v_1 \in V(K_{1,n-1})$  with  $\deg(v_1) = n - 1$  and  $\deg(N(v_1)) = 1$ , where  $N(v_1) = \{v_2, v_3, \dots, v_n\}$ . Clearly, each edge  $\{v_1v_2, v_1v_3, \dots, v_1v_n\}$  takes one vertex on each edge having degree 2, so that the resulting graph will be subdivided star  $K_{1,n-1}^*$ , in which  $\{v_1\}$  and  $\{v_2, v_3, \dots, v_n\}$  vertices do not lose their properties. But the maximum degree vertex  $v_1$  is a cut vertex of  $K_{1,n-1}^*$ . Therefore  $b(K_{1,n-1}) < b(K_{1,n-1}^*)$  for  $n \geq 4$  vertices. Since each newly added vertex  $\{u_i\}$  is adjacent to exactly one pendent vertex  $\{v_j\}$ , where  $i = j$  and  $2 \leq i, j \leq n$ , in  $K_{1,n-1}^*$ . By the definition of binding number  $|N(X)| = |X|$ . Hence the result follows.  $\square$

**Definition 2.11** A  $B_{t,k}$  graph is said to be a Banana tree if the graph is obtained by connecting one pendant vertex of each  $t$ -copies of an  $k$ -star graph with a single root vertex that is distinct from all the stars.

**Theorem 2.12** For any Banana tree  $B_{t,k}$  with  $t \geq 2$  copies and  $k \geq 3$  number of stars,

$$b(B_{t,k}) = \frac{1}{k-2}.$$

*Proof* Let  $t$  be the number of distinct  $k$ -stars. Then it has  $k - 1$ -pendant vertices and the binding number of each  $k$ -stars is  $\frac{1}{k-1}$ . But in  $B_{t,k}$ , each  $t$  copies of distinct  $k$ -stars are joined by single root vertex. Then the resulting graph is connected and each  $k$ -star has  $k - 2$  number of vertices having degree 1, which are connected to unique vertex. By Observation 2.1, the result follows.  $\square$

**Definition 2.13** A caterpillar tree  $C^*(T)$  is a tree in which removing all the pendant vertices and incident edges produces a path graph.

For example,  $b(C^*(K_1)) = 0$ ;  $b(C^*(P_2)) = b(C^*(P_4)) = 1$ ;  $b(C^*(P_3)) = \frac{1}{2}$ ;  $b(C^*(P_5)) = \frac{2}{3}$  and  $b(C^*(K_{1,n-1})) = \frac{1}{n-1}$ .

**Theorem 2.14** For any caterpillar tree  $C^*(T)$  with  $n \geq 3$  vertices,

$$b(K_{1,n-1}) \leq b(C^*(T)) \leq b(P_n).$$

*Proof* By mathematical induction, if  $n = 3$ , then by Theorem 1.1 and Observation 2.1, we have  $b(K_{1,2}) = b(C^*(T)) = b(P_3) = \frac{1}{2}$ . Thus the result follows. Assume that the result is true for  $n = k$ . Now we shall prove the result for  $n > k$ . Let  $C^*(T)$  be a Caterpillar tree with  $k + 1$ -vertices. Then the following cases arise:

**Case 1.** If  $k + 1$  is odd, then  $b(C^*(T)) \leq \frac{k}{k+1}$ .

**Case 2.** If  $k + 1$  is even, then  $b(C^*(T)) \leq 1$ .

By above cases, we have  $b(C^*(T)) \leq b(P_n)$ . Since,  $k$  vertices in  $C^*(T)$  exist  $k$ -stars, which

contributed at least  $\frac{1}{k-1}$ . Hence the lower bound follows.  $\square$

**Definition 2.15** *The binary tree  $B^*$  is a tree like structure that is rooted and in which each vertex has at least two children and child of a vertex is designated as its left or right child.*

To prove our next result we make use of the following conditions of Binary tree  $B^*$ .

$C_1$ : If  $B^*$  has at least one vertex having two children and that two children has no any child.

$C_2$ : If  $B^*$  has no vertex having two children which are not having any child.

**Theorem 2.16** *Let  $B^*$  be a Binary tree with  $n \geq 3$  vertices. Then*

$$b(B^*) = \begin{cases} \frac{1}{2} & \text{if } B^* \text{ satisfy } C_1; \\ b(P_n) & \text{if } B^* \text{ satisfy } C_2. \end{cases}$$

*Proof* Let  $B^*$  be a Binary tree with  $n \geq 3$  vertices. Then the following cases are arises:

**Case 1.** Suppose binary tree  $B^*$  has only one vertex, say  $v_1$  has two children and that two children has no any child. Then only vertex  $v_1$  has two pendant vertices and no other vertex has more than two pendant vertices. That is maximum at most two pendant vertices are connected to unique vertex. There fore  $b(B^*) = \frac{1}{2}$  follows.

**Case 2.** Suppose binary tree  $B^*$  has no vertex having two free child. That is each non-pendant vertex having only one child, then this binary tree gives path. This implies that  $b(B^*) = b(P_n)$  with  $n \geq 3$  vertices. Thus the result follows.  $\square$

**Definition 2.17** *The  $t$ -centipede  $C_t^*$  is the tree on  $2t$ -vertices obtained by joining the bottoms of  $t$  - copies of the path graph  $P_2$  laid in a row with edges.*

**Theorem 2.18** *For any  $t$ -centipede  $C_t^*$  with  $2t$ -vertices,*

$$b(C_t^*) = 1.$$

*Proof* If  $n = 1$ , then tree  $C_1^*$  is a 1-centipede with 2-vertices. Thus  $b(C_1^*) = 1$ . Suppose the result is true for  $n > 1$  vertices, say  $n = t$  for some  $t$ , that is  $b(C_t^*) = 1$ . Further, we prove  $n = t+1$ ,  $b(C_{t+1}^*) = 1$ . In a  $(t+1)$  - centipede exactly one vertex from each of the  $(k+1)$ - copies of  $P_2$  are laid on a row with edges. Hence the resulting graph must be connected and each such vertex is connected to exactly one pendant vertex. By the definition of binding number  $|N(X)| = |X|$ . Hence the result follows.  $\square$

**Definition 2.19** *The Fire-cracker graph  $F_{t,s}$  is a tree obtained by the concatenation of  $t$  - copies of  $s$  - stars by linking one pendant vertex from each.*

**Theorem 2.20** For any Fire-cracker graph  $F_{t,s}$  with  $t \geq 2$  and  $s \geq 3$ .

$$b(F_{t,s}) = \frac{1}{s-1}.$$

*Proof* If  $s = 2$ , then Fire-cracker graph  $F_{t,2}$  is a  $t$ -centipede and  $b(F_{t,2}) = 1$ . If  $t \geq 2$  and  $s \geq 3$ , then  $t$  - copies of  $s$  - stars are connected by adjoining one pendant vertex from each  $s$ -stars. This implies that the resulting graph is connected and a Fire-cracker graph  $F_{t,s}$ . Then this connected graph has  $(s-2)$ -vertices having degree 1, which are connected to unique vertex. Hence the result follows.  $\square$

**Theorem 2.21** For any nontrivial tree  $T$ ,

$$\frac{1}{n-1} \leq b(T) \leq 1.$$

Further, the lower bound attains if and only if  $T = K_{1,n-1}$  and the upper bound attains if the tree  $T$  has 1-factor or there exists a realizing set  $X$  such that  $X \cap N(X) = \phi$ .

*Proof* The upper bound is proved by Woodall in [7] with the fact of  $\delta(T) = 1$ . Let  $X \in F$  and  $\frac{|N(X)|}{|X|} = b(T)$ . Then  $|N(X)| \geq 1$ , since the set  $X$  is not empty. Suppose,  $|N(X)| \geq n - \delta(T) + 1$ . If  $\delta(T) = 1$ , then any vertex of  $T$  is adjacent to atleast one vertex in  $X$ . This implies that  $N(X) = V(T)$ , which is a contradiction. Therefore  $|X| \leq n - 1$  and  $b(T) = |N(X)|/|X| \geq 1/(n-1)$ . Thus the lower bound follows.  $\square$

**Acknowledgments** The authors wish to thank Prof.N.D.Soner for his help and valuable suggestions in the preparation of this paper.

## References

- [1] I.Anderson, Binding number of a graphs: A Survey, *Advances in Graph Theory*, ed. V. R. Kulli, Vishwa International Publications, Gulbarga (1991) 1-10.
- [2] W.H.Cunningham, Computing the binding number of a graph, *Discrete Applied Math.* 27(1990) 283-285.
- [3] F.Harary, *Graph Theory*, Addison Wesley, Reading Mass, (1969).
- [4] V.G.Kane, S.P.Mohanty and R.S.Hales, Product graphs and binding number, *Ars Combin.*, 11 (1981) 201-224.
- [5] P.Kwasnik and D.R.Woodall, Binding number of a graph and the existence of  $k$ - factors, *Quarterly J. Math.* 38 (1987) 221-228.
- [6] N.Tokushinge, Binding number and minimum degree for  $k$ -factors, *J. Graph Theory* 13(1989) 607-617.
- [7] D.R.Woodall, The binding number of a graph and its Anderson number, *J. Combinatorial Theory Ser. B*, 15 (1973) 225-255.
- [8] Wayne Goddard, The Binding Number of Trees and  $K_{(1,3)}$ -free Graphs, *J. Combin. Math. Combin. Comput*, 7 (1990)193-200.