

Smarandache Directionally n -Signed Graphs — A Survey

P.Siva Kota Reddy

(Department of Mathematics, Acharya Institute of Technology, Soladevanahalli, Bangalore-560 107, India)

E-mail: reddy_math@yahoo.com; pskreddy@acharya.ac.in

Abstract: Let $G = (V, E)$ be a graph. By *directional labeling* (or *d-labeling*) of an edge $x = uv$ of G by an ordered n -tuple $(a_1, a_2, \dots, a_n)$, we mean a labeling of the edge x such that we consider the label on uv as $(a_1, a_2, \dots, a_n)$ in the direction from u to v , and the label on x as $(a_n, a_{n-1}, \dots, a_1)$ in the direction from v to u . In this survey, we study graphs, called *(n, d) -sigraphs*, in which every edge is *d-labeled* by an n -tuple $(a_1, a_2, \dots, a_n)$, where $a_k \in \{+, -\}$, for $1 \leq k \leq n$. Several variations and characterizations of directionally n -signed graphs have been proposed and studied. These include the various notions of balance and others.

Key Words: Signed graphs, directional labeling, complementation, balance.

AMS(2010): 05C 22

§1. Introduction

For graph theory terminology and notation in this paper we follow the book [3]. All graphs considered here are finite and simple. There are two ways of labeling the edges of a graph by an ordered n -tuple $(a_1, a_2, \dots, a_n)$ (See [10]).

1. *Undirected labeling* or *labeling*. This is a labeling of each edge uv of G by an ordered n -tuple $(a_1, a_2, \dots, a_n)$ such that we consider the label on uv as $(a_1, a_2, \dots, a_n)$ irrespective of the direction from u to v or v to u .

2. *Directional labeling* or *d-labeling*. This is a labeling of each edge uv of G by an ordered n -tuple $(a_1, a_2, \dots, a_n)$ such that we consider the label on uv as $(a_1, a_2, \dots, a_n)$ in the direction from u to v , and $(a_n, a_{n-1}, \dots, a_1)$ in the direction from v to u .

Note that the *d-labeling* of edges of G by ordered n -tuples is equivalent to labeling the symmetric digraph $\vec{G} = (V, \vec{E})$, where uv is a symmetric arc in $\vec{G}$ if, and only if, uv is an edge in G , so that if $(a_1, a_2, \dots, a_n)$ is the *d-label* on uv in G , then the labels on the arcs $\vec{uv}$ and $\vec{vu}$ are $(a_1, a_2, \dots, a_n)$ and $(a_n, a_{n-1}, \dots, a_1)$ respectively.

Let H_n be the *n -fold sign group*, $H_n = \{+, -\}^n = \{(a_1, a_2, \dots, a_n) : a_1, a_2, \dots, a_n \in \{+, -\}\}$ with co-ordinate-wise multiplication. Thus, writing $a = (a_1, a_2, \dots, a_n)$ and $t =$

¹Reported at the First International Conference on Smarandache Multispaces and Multistructures, June 28-30, 2013, Beijing, P.R.China.

²Received May 16, 2013, Accepted June 10, 2013.

$(t_1, t_2, \dots, t_n)$ then $at := (a_1t_1, a_2t_2, \dots, a_nt_n)$. For any $t \in H_n$, the *action* of t on H_n is $a^t = at$, the co-ordinate-wise product.

Let $n \geq 1$ be a positive integer. An n -signed graph (n -signed digraph) is a graph $G = (V, E)$ in which each edge (arc) is labeled by an ordered n -tuple of signs, i.e., an element of H_n . A signed graph $G = (V, E)$ is a graph in which each edge is labeled by $+$ or $-$. Thus a 1-signed graph is a signed graph. Signed graphs are well studied in literature (See for example [1, 4-7, 13-21, 23, 24]).

In this survey, we study graphs in which each edge is labeled by an ordered n -tuple $a = (a_1, a_2, \dots, a_n)$ of signs (i.e., an element of H_n) in one direction but in the other direction its label is the reverse: $a^r = (a_n, a_{n-1}, \dots, a_1)$, called *directionally labeled n -signed graphs* (or (n, d) -signed graphs).

Note that an n -signed graph $G = (V, E)$ can be considered as a symmetric digraph $\vec{G} = (V, \vec{E})$, where both $\vec{uv}$ and $\vec{vu}$ are arcs if, and only if, uv is an edge in G . Further, if an edge uv in G is labeled by the n -tuple $(a_1, a_2, \dots, a_n)$, then in $\vec{G}$ both the arcs $\vec{uv}$ and $\vec{vu}$ are labeled by the n -tuple $(a_1, a_2, \dots, a_n)$.

In [1], the authors study voltage graph defined as follows: A *voltage graph* is an ordered triple $\vec{G} = (V, \vec{E}, M)$, where V and $\vec{E}$ are the vertex set and arc set respectively and M is a group. Further, each arc is labeled by an element of the group M so that if an arc $\vec{uv}$ is labeled by an element $a \in M$, then the arc $\vec{vu}$ is labeled by its inverse, a^{-1} .

Since each n -tuple $(a_1, a_2, \dots, a_n)$ is its own inverse in the group H_n , we can regard an n -signed graph $G = (V, E)$ as a voltage graph $\vec{G} = (V, \vec{E}, H_n)$ as defined above. Note that the d -labeling of edges in an (n, d) -signed graph considering the edges as symmetric directed arcs is different from the above labeling. For example, consider a $(4, d)$ -signed graph in Figure 1. As mentioned above, this can also be represented by a symmetric 4-signed digraph. Note that this is not a voltage graph as defined in [1], since for example; the label on $\vec{v_2v_1}$ is not the (group) inverse of the label on $\vec{v_1v_2}$.




Fig.1

In [8-9], the authors initiated a study of $(3, d)$ and $(4, d)$ -Signed graphs. Also, discussed some applications of $(3, d)$ and $(4, d)$ -Signed graphs in real life situations.

In [10], the authors introduced the notion of complementation and generalize the notion of balance in signed graphs to the directionally n -signed graphs. In this context, the authors look upon two kinds of complementation: complementing some or all of the signs, and reversing the order of the signs on each edge. Also given some motivation to study (n, d) -signed graphs in connection with relations among human beings in society.

In [10], the authors defined complementation and isomorphism for (n, d) -signed graphs as

follows: For any $t \in H_n$, the t -complement of $a = (a_1, a_2, \dots, a_n)$ is: $a^t = at$. The reversal of $a = (a_1, a_2, \dots, a_n)$ is: $a^r = (a_n, a_{n-1}, \dots, a_1)$. For any $T \subseteq H_n$, and $t \in H_n$, the t -complement of T is $T^t = \{a^t : a \in T\}$.

For any $t \in H_n$, the t -complement of an (n, d) -signed graph $G = (V, E)$, written G^t , is the same graph but with each edge label $a = (a_1, a_2, \dots, a_n)$ replaced by a^t . The reversal G^r is the same graph but with each edge label $a = (a_1, a_2, \dots, a_n)$ replaced by a^r .

Let $G = (V, E)$ and $G' = (V', E')$ be two (n, d) -signed graphs. Then G is said to be *isomorphic* to G' and we write $G \cong G'$, if there exists a bijection $\phi : V \rightarrow V'$ such that if uv is an edge in G which is d -labeled by $a = (a_1, a_2, \dots, a_n)$, then $\phi(u)\phi(v)$ is an edge in G' which is d -labeled by a , and conversely.

For each $t \in H_n$, an (n, d) -signed graph $G = (V, E)$ is t -self complementary, if $G \cong G^t$. Further, G is *self reverse*, if $G \cong G^r$.

Proposition 1.1(E. Sampathkumar et al. [10]) *For all $t \in H_n$, an (n, d) -signed graph $G = (V, E)$ is t -self complementary if, and only if, G^a is t -self complementary, for any $a \in H_n$.*

For any cycle C in G , let $\mathcal{P}(\vec{C})$ [10] denotes the product of the n -tuples on C given by $(a_{11}, a_{12}, \dots, a_{1n})(a_{21}, a_{22}, \dots, a_{2n}) \dots (a_{m1}, a_{m2}, \dots, a_{mn})$ and

$$\mathcal{P}(\overleftarrow{C}) = (a_{mn}, a_{m(n-1)}, \dots, a_{m1})(a_{(m-1)n}, a_{(m-1)(n-1)}, \dots, a_{(m-1)1}) \dots (a_{1n}, a_{1(n-1)}, \dots, a_{11}).$$

Similarly, for any path P in G , $\mathcal{P}(\vec{P})$ denotes the product of the n -tuples on P given by $(a_{11}, a_{12}, \dots, a_{1n})(a_{21}, a_{22}, \dots, a_{2n}) \dots (a_{m-1,1}, a_{m-1,2}, \dots, a_{m-1,n})$ and

$$\mathcal{P}(\overleftarrow{P}) = (a_{(m-1)n}, a_{(m-1)(n-1)}, \dots, a_{(m-1)1}) \dots (a_{1n}, a_{1(n-1)}, \dots, a_{11}).$$

An n -tuple $(a_1, a_2, \dots, a_n)$ is *identity n -tuple*, if each $a_k = +$, for $1 \leq k \leq n$, otherwise it is a *non-identity n -tuple*. Further an n -tuple $a = (a_1, a_2, \dots, a_n)$ is *symmetric*, if $a^r = a$, otherwise it is a *non-symmetric n -tuple*. In (n, d) -signed graph $G = (V, E)$ an edge labeled with the identity n -tuple is called an *identity edge*, otherwise it is a *non-identity edge*.

Note that the above products $\mathcal{P}(\vec{C})$ ($\mathcal{P}(\vec{P})$) as well as $\mathcal{P}(\overleftarrow{C})$ ($\mathcal{P}(\overleftarrow{P})$) are n -tuples. In general, these two products need not be equal.

§2. Balance in an (n, d) -Signed Graph

In [10], the authors defined two notions of balance in an (n, d) -signed graph $G = (V, E)$ as follows:

Definition 2.1 *Let $G = (V, E)$ be an (n, d) -sigraph. Then,*

- (i) G is *identity balanced* (or *i-balanced*), if $\mathcal{P}(\vec{C})$ on each cycle of G is the identity n -tuple, and
- (ii) G is *balanced*, if every cycle contains an even number of non-identity edges.

Note: An *i-balanced* (n, d) -sigraph need not be balanced and conversely. For example, consider the $(4, d)$ -sigraphs in Figure.2. In Figure.2(a) G is an *i-balanced* but not balanced, and in Figure.2(b) G is balanced but not *i-balanced*.




Fig.2

2.1 Criteria for balance

An (n, d) -signed graph $G = (V, E)$ is i -balanced if each non-identity n -tuple appears an even number of times in $P(\vec{C})$ on any cycle of G .

However, the converse is not true. For example see Figure.3(a). In Figure.3(b), the number of non-identity 4-tuples is even and hence it is balanced. But it is not i -balanced, since the 4-tuple $(+ + - -)$ (as well as $(- - + +)$) does not appear an even number of times in $P(\vec{C})$ of 4-tuples.




Fig.3

In [10], the authors obtained following characterizations of balanced and i -balanced (n, d) -sigraphs:

Proposition 2.2(E.Sampathkumar et al. [10]) *An (n, d) -signed graph $G = (V, E)$ is balanced if, and only if, there exists a partition $V_1 \cup V_2$ of V such that each identity edge joins two vertices in V_1 or V_2 , and each non-identity edge joins a vertex of V_1 and a vertex of V_2 .*

As earlier discussed, let $P(C)$ denote the product of the n -tuples in $P(\vec{C})$ on any cycle C in an (n, d) -sigraph $G = (V, E)$.

Theorem 2.3(E.Sampathkumar et al. [10]) *An (n, d) -signed graph $G = (V, E)$ is i -balanced if, and only if, for each k , $1 \leq k \leq n$, the number of n -tuples in $P(C)$ whose k^{th} co-ordinate is $-$ is even.*

In H_n , let S_1 denote the set of non-identity symmetric n -tuples and S_2 denote the set of non-symmetric n -tuples. The product of all n -tuples in each S_k , $1 \leq k \leq 2$ is the identity n -tuple.

Theorem 2.4(E.Sampathkumar et al. [10]) *An (n, d) -signed graph $G = (V, E)$ is i -balanced, if both of the following hold:*

- (i) *In $P(C)$, each n -tuple in S_1 occurs an even number of times, or each n -tuple in S_1 occurs odd number of times (the same parity, or equal mod 2).*
- (ii) *In $P(C)$, each n -tuple in S_2 occurs an even number of times, or each n -tuple in S_2 occurs an odd number of times.*

In [11], the authors obtained another characterization of i -balanced (n, d) -signed graphs as follows:

Theorem 2.5(E.Sampathkumar et al. [11]) *An (n, d) -signed graph $G = (V, E)$ is i -balanced if, and only if, any two vertices u and v have the property that for any two edge distinct $u - v$ paths $\vec{P}_1 = (u = u_0, u_1, \dots, u_m = v)$ and $\vec{P}_2 = (u = v_0, v_1, \dots, v_n = v)$ in G , $\mathcal{P}(\vec{P}_1) = (\mathcal{P}(\vec{P}_2))^r$ and $\mathcal{P}(\vec{P}_2) = (\mathcal{P}(\vec{P}_1))^r$.*

From the above result, the following are the easy consequences:

Corollary 2.6 *In an i -balanced (n, d) -signed graph G if two vertices are joined by at least 3 paths then the product of n tuples on any paths joining them must be symmetric.*

A graph $G = (V, E)$ is said to be k -connected for some positive integer k , if between any two vertices there exists at least k disjoint paths joining them.

Corollary 2.7 *If the underlying graph of an i -balanced (n, d) -signed graph is 3-connected, then all the edges in G must be labeled by a symmetric n -tuple.*

Corollary 2.8 *A complete (n, d) -signed graph on $p \geq 4$ is i -balanced then all the edges must be labeled by symmetric n -tuple.*

2.2 Complete (n, d) -Signed Graphs

In [11], the authors defined: an (n, d) -sigraph is *complete*, if its underlying graph is complete. Based on the complete (n, d) -signed graphs, the authors proved the following results: An (n, d) -signed graph is *complete*, if its underlying graph is complete.

Proposition 2.9(E.Sampathkumar et al. [11]) *The four triangles constructed on four vertices $\{a, b, c, d\}$ can be directed so that given any pair of vertices say (a, b) the product of the edges of these 4 directed triangles is the product of the n -tuples on the arcs $\vec{ab}$ and $\vec{ba}$.*

Corollary 2.10 *The product of the n -tuples of the four triangles constructed on four vertices $\{a, b, c, d\}$ is identity if at least one edge is labeled by a symmetric n -tuple.*

The i -balance base with axis a of a complete (n, d) -signed graph $G = (V, E)$ consists list of the product of the n -tuples on the triangles containing a [11].

Theorem 2.11(E.Sampathkumar et al. [11]) *If the i -balance base with axis a and n -tuple of an*

edge adjacent to a is known, the product of the n -tuples on all the triangles of G can be deduced from it.

In the statement of above result, it is not necessary to know the n -tuple of an edge incident at a . But it is sufficient that an edge incident at a is a symmetric n -tuple.

Theorem 2.12(E.Sampathkumar et al. [11]) *A complete (n, d) -sigraph $G = (V, E)$ is i -balanced if, and only if, all the triangles of a base are identity.*

Theorem 2.13(E.Sampathkumar et al. [11]) *The number of i -balanced complete (n, d) -sigraphs of m vertices is p^{m-1} , where $p = 2^{\lceil n/2 \rceil}$.*

§3. Path Balance in (n, d) -Signed Graphs

In [11], E.Sampathkumar et al. defined the path balance in an (n, d) -signed graphs as follows:

Let $G = (V, E)$ be an (n, d) -sigraph. Then G is

1. *Path i -balanced*, if any two vertices u and v satisfy the property that for any $u - v$ paths P_1 and P_2 from u to v , $\mathcal{P}(\vec{P}_1) = \mathcal{P}(\vec{P}_2)$.
2. *Path balanced* if any two vertices u and v satisfy the property that for any $u - v$ paths P_1 and P_2 from u to v have same number of non identity n -tuples.

Clearly, the notion of path balance and balance coincides. That is an (n, d) -signed graph is balanced if, and only if, G is path balanced.

If an (n, d) signed graph G is i -balanced then G need not be path i -balanced and conversely.

In [11], the authors obtained the characterization path i -balanced (n, d) -signed graphs as follows:

Theorem 3.1(Characterization of path i -balanced $(n; d)$ signed graphs) *An (n, d) -signed graph is path i -balanced if, and only if, any two vertices u and v satisfy the property that for any two vertex disjoint $u - v$ paths P_1 and P_2 from u to v , $\mathcal{P}(\vec{P}_1) = \mathcal{P}(\vec{P}_2)$.*

§4. Local Balance in (n, d) -Signed Graphs

The notion of local balance in signed graph was introduced by F. Harary [5]. A signed graph $S = (G, \sigma)$ is locally at a vertex v , or S is *balanced at v* , if all cycles containing v are balanced. A cut point in a connected graph G is a vertex whose removal results in a disconnected graph. The following result due to Harary [5] gives interdependence of local balance and cut vertex of a signed graph.

Theorem 4.1(F.Harary [5]) *If a connected signed graph $S = (G, \sigma)$ is balanced at a vertex u . Let v be a vertex on a cycle C passing through u which is not a cut point, then S is balanced at v .*

In [11], the authors extend the notion of local balance in signed graph to (n, d) -signed graphs as follows: Let $G = (V, E)$ be a (n, d) -signed graph. Then for any vertices $v \in V(G)$, G is *locally i -balanced at v* (*locally balanced at v*) if all cycles in G containing v is i -balanced (balanced).

Analogous to the above result, in [11] we have the following for an (n, d) signed graphs:

Theorem 4.2 *If a connected (n, d) -signed graph $G = (V, E)$ is locally i -balanced (locally balanced) at a vertex u and v be a vertex on a cycle C passing through u which is not a cut point, then S is locally i -balanced (locally balanced) at v .*

§5. Symmetric Balance in (n, d) -Signed Graphs

In [22], P.S.K.Reddy and U.K.Misra defined a new notion of balance called *symmetric balance* or *s-balanced* in (n, d) -signed graphs as follows:

Let $n \geq 1$ be an integer. An n -tuple $(a_1, a_2, \dots, a_n)$ is *symmetric*, if $a_k = a_{n-k+1}, 1 \leq k \leq n$. Let $H_n = \{(a_1, a_2, \dots, a_n) : a_k \in \{+, -\}, a_k = a_{n-k+1}, 1 \leq k \leq n\}$ be the set of all symmetric n -tuples. Note that H_n is a group under coordinate wise multiplication, and the order of H_n is 2^m , where $m = \lceil n/2 \rceil$. Let $G = (V, E)$ be an (n, d) -signed graph. Then G is *symmetric balanced* or *s-balanced* if $P(\vec{C})$ on each cycle C of G is symmetric n -tuple.

Note: If an (n, d) -signed graph $G = (V, E)$ is i -balanced then clearly G is s -balanced. But a s -balanced (n, d) -signed graph need not be i -balanced. For example, the $(4, d)$ -signed graphs in Figure 4. G is an s -balanced but not i -balanced.




Fig.4

In [22], the authors obtained the following results based on symmetric balance or s -balanced in (n, d) -signed graphs.

Theorem 5.1(P.S.K.Reddy and U.K.Mishra [22]) *A (n, d) -signed graph is s -balanced if and only if every cycle of G contains an even number of non-symmetric n -tuples.*

The following result gives a necessary and sufficient condition for a balanced (n, d) -signed graph to be s -balanced.

Theorem 5.2(P.S.K.Reddy and U.K.Mishra [22]) *A balanced (n, d) signed graph $G = (V, E)$ is s -balanced if and only if every cycle of G contains even number of non identity symmetric n*

tuples.

In [22], the authors obtained another characterization of s -balanced (n, d) -signed graphs, which is analogous to the partition criteria for balance in signed graphs due to Harary [4].

Theorem 5.3(Characterization of s -balanced (n, d) -sigraph) *An (n, d) -signed graph $G = (V, E)$ is s balanced if and only if the vertex set $V(G)$ of G can be partitioned into two sets V_1 and V_2 such that each symmetric edge joins the vertices in the same set and each non-symmetric edge joins a vertex of V_1 and a vertex of V_2 .*

An n -marking $\mu : V(G) \rightarrow H_n$ of an (n, d) -signed graph $G = (V, E)$ is an assignment n -tuples to the vertices of G . In [22], the authors given another characterization of s -balanced (n, d) -signed graphs which gives a relationship between the n -marking and s -balanced (n, d) -signed graphs.

Theorem 5.4(P.S.K.Reddy and U.K.Mishra [22]) *An (n, d) -signed graph $G = (V, E)$ is s -balanced if and only if there exists an n -marking μ of vertices of G such that if the n -tuple on any arc $\vec{uv}$ is symmetric or nonsymmetric according as the n -tuple $\mu(u)\mu(v)$ is.*

§6. Directionally 2-Signed Graphs

In [12], E.Sampathkumar et al. proved that the directionally 2-signed graphs are equivalent to bidirected graphs, where each end of an edge has a sign. A bidirected graph implies a signed graph, where each edge has a sign. Signed graphs are the special case $n = 1$, where directionality is trivial. Directionally 2-signed graphs (or $(2, d)$ -signed graphs) are also special, in a less obvious way. A bidirected graph $B = (G, \beta)$ is a graph $G = (V, E)$ in which each end (e, u) of an edge $e = uv$ has a sign $\beta(e, u) \in \{+, -\}$. G is the underlying graph and β is the bidirection. (The $+$ sign denotes an arrow on the u -end of e pointed into the vertex u ; a $-$ sign denotes an arrow directed out of u . Thus, in a bidirected graph each end of an edge has an independent direction. Bidirected graphs were defined by Edmonds [2].) In view of this, E.Sampathkumar et al. [12] proved the following result:

Theorem 6.1(E.Sampathkumar et al. [12]) *Directionally 2-signed graphs are equivalent to bidirected graphs.*

§7. Conclusion

In this brief survey, we have described directionally n -signed graphs (or (n, d) -signed graphs) and their characterizations. Many of the characterizations are more recent. This is an active area of research. We have included a set of references which have been cited in our description. These references are just a small part of the literature, but they should provide a good start for readers interested in this area.

References

- [1] B.D.Acharya and M. Acharya, New algebraic models of a Social System, *Indian J. of Pure and Appl. Math.*, 17(2) 1986, 152-168.
- [2] J.Edmonds and E.L.Johnson, Matching: a well-solved class of integral linear programs, in: Richard Guy et al., eds., *Combinatorial Structures and Their Applications* (Proc. Calgary Int. Conf., Calgary, 1969), pp. 89-92. Gordon and Breach, New York, 1970.
- [3] F.Harary, *Graph Theory*, Addison-Wesley Publishing Co., 1969.
- [4] F.Harary, On the notion of balance of a signed graph, *Michigan Math. J.*, 2(1953), 143-146.
- [5] F.Harary, On local balance and N -balance in signed graphs, *Michigan Math. J.*, 3(1955), 37-41.
- [6] F.Harary, R. Norman and D. Cartwright, *Structural models: An introduction to the theory of directed graphs*, J. Wiley, New York, 1965.
- [7] R.Rangarajan, M. S.Subramanya and P.Siva Kota Reddy, The H -Line Signed Graph of a Signed Graph, *International J. Math. Combin.*, 2 (2010), 37-43.
- [8] E.Sampathkumar, P.Siva Kota Reddy and M. S.Subramanya, $(3, d)$ -sigraph and its applications, *Advn. Stud. Contemp. Math.*, 17(1) (2008), 57-67.
- [9] E.Sampathkumar, P.Siva Kota Reddy and M. S.Subramanya, $(4, d)$ -sigraph and its applications, *Advn. Stud. Contemp. Math.*, 20(1) (2010), 115-124.
- [10] E.Sampathkumar, P.Siva Kota Reddy, and M.S.Subramanya, Directionally n -signed graphs, in: B.D. Acharya et al., eds., *Advances in Discrete Mathematics and Applications: Mysore, 2008* (Proc. Int. Conf. Discrete Math., ICDM-2008), pp. 153-160, Ramanujan Math. Soc. Lect. Notes Ser., 13, Ramanujan Mathematical Society, Mysore, India, 2010.
- [11] E.Sampathkumar, P.Siva Kota Reddy, and M.S.Subramanya, Directionally n -signed graphs-II, *Int. J. Math. Combin.*, 4 (2009), 89-98.
- [12] E.Sampathkumar, M.A.Sriraj and T.Zaslavsky, Directionally 2-signed and bidirected graphs.
- [13] P.Siva Kota Reddy and M.S.Subramanya, Signed graph equation $L^k(S) \sim \overline{S}$, *International J. Math. Combin.*, 4 (2009), 84-88.
- [14] P.Siva Kota Reddy, S.Vijay and V.Lokesha, n^{th} Power signed graphs, *Proceedings of the Jangjeon Math. Soc.*, 12(3) (2009), 307-313.
- [15] P.Siva Kota Reddy, S.Vijay and H.C.Savithri, A Note on Path Sidigraphs, *International J. Math. Combin.*, 1 (2010), 42-46.
- [16] P.Siva Kota Reddy, S.Vijay and V.Lokesha, n^{th} Power signed graphs-II, *International J. Math. Combin.*, 1 (2010), 74-79.
- [17] P.Siva Kota Reddy and S.Vijay, Total minimal dominating signed graph, *International J. Math. Combin.*, 3 (2010), 11-16.
- [18] P.Siva Kota Reddy and K.V.Madhusudhan, Negation switching equivalence in signed graphs, *International J. Math. Combin.*, 3 (2010), 85-90.
- [19] P.Siva Kota Reddy, t -Path Sigraphs, *Tamsui Oxford J. of Math. Sciences*, 26(4) (2010), 433-441.
- [20] P.Siva Kota Reddy, B.Prashanth and Kavita S. Permi, A Note on Antipodal Signed Graphs, *International J. Math. Combin.*, 1 (2011), 107-112.

- [21] P.Siva Kota Reddy, B.Prashanth, and T.R.Vasanth Kumar, Antipodal Signed Directed Graphs, *Advn. Stud. Contemp. Math.*, 21(4) (2011), 355-360.
- [22] P.Siva Kota Reddy and U.K.Misra, Directionally n -signed graphs-III: The notion of symmetric balance, Submitted.
- [23] T.Zaslavsky, Signed Graphs, *Discrete Appl. Math.*, 4(1)(1982), 47-74.
- [24] T.Zaslavsky, A mathematical bibliography of Signed and Gain graphs and its allied areas, *Electron. J. Combin.*, Dynamic Surveys in Combinatorics (1998), No. DS8. Eighth ed. (2012).