



6 Interval Valued Pythagorean Fuzzy Ideals in Semigroups

Veerappan Chinnadurai*, Arul Selvam

Department of Mathematics, Annamalai University, Chidambaram, Tamilnadu, India.

PAPER INFO	ABSTRACT
Chronicle: Received: 01 July 2020 Reviewed: 11 August 2020 Revised: 04 September 2020 Accepted: 24 November 2020	In this paper, we define the new notion of interval-valued Pythagorean fuzzy ideals in semigroups and established the properties of its with suitable examples. Also, we introduce the concept of interval valued Pythagorean fuzzy sub-semigroup, interval valued Pythagorean fuzzy left (resp. right) ideal, interval valued Pythagorean fuzzy bi-ideal, interval valued Pythagorean fuzzy interior ideal and homomorphism of an interval valued Pythagorean fuzzy ideal in semigroups with suitable illustration. We show that every interval valued Pythagorean fuzzy left (resp. right) ideal is an interval valued Pythagorean fuzzy bi-ideal.
Keywords: Pythagorean Fuzzy. Fuzzy Ideals. Homomorphism. Semigroups.	

1. Introduction

In 1965, Zadeh [18, 19] introduced the concept of a fuzzy set. He also developed the notion of interval-valued fuzzy set in 1975, which extends the fuzzy set. A semigroup is an algebraic structure comprising a non-empty set together with an associative binary operation. Atanassov [2] introduced the intuitionistic fuzzy set with some properties. Atanassov [3] developed the concept of interval-valued intuitionistic fuzzy set. Thillaigovindan and Chinnadurai [15, 16] discussed interval-valued fuzzy ideals in algebraic structures. In 2018, Chen [4, 5] introduced the concept of interval-valued Pythagorean fuzzy outranking of various methods in the application. Garg [8, 9] presented the notion of interval-valued Pythagorean fuzzy sets of multi-criteria decision-making methods. In 2013, Yager [17] started the notion of Pythagorean fuzzy set, the sum of the squares of membership and non-membership belongs to the unit interval $[0, 1]$. Peng [13]



* Corresponding author

E-mail address: chinnadurau@gmail.com

DOI: 10.22105/jfea.2020.252687.1023

developed the new operations for an interval-valued Pythagorean fuzzy set. Peng and Yang [14] presented the notion of interval-valued Pythagorean fuzzy set. In 2019, Hussain et al. [10] started the notions of rough Pythagorean fuzzy ideals in the semigroups. Akram[1] established the properties of fuzzy lie algebras. Kumar et al. [11] approached transportation decision making problems using Pythagorean fuzzy set. Das and Edalatpanah [6] studied the concept of fuzzy linear fractional progress with trapezoidal fuzzy numbers. Edalatpanah [7] used triangular intuitionistic fuzzy numbers to deal with data envelopment analysis model. Najafi and Edalatpanah [12] used iterative methods to study linear complementarily problems. In this paper, we discuss some of the properties of interval-valued Pythagorean fuzzy ideals in the semigroups.

2. Preliminaries

Definition 1. [12]. Let X be a universe of discourse, A Pythagorean fuzzy set (PFS) $P = \{w, \phi_p(w), \psi_p(w)/w \in X\}$ where $\phi: X \rightarrow [0,1]$ and $\psi: X \rightarrow [0,1]$ represent the degree of membership and non-membership of the object $w \in X$ to the set P subset to the condition $0 \leq (\phi_p(w))^2 + (\psi_p(w))^2 \leq 1$ for all $w \in X$. For the sake of simplicity a PFS is denoted as $P = (\phi_p(w), \psi_p(w))$.

3. Interval-Valued Pythagorean Fuzzy Ideals in Semigroups

Definition 2. An Interval-Valued Pythagorean Fuzzy Set (IVPFS) $\tilde{P} = [\tilde{\phi}_p, \tilde{\psi}_p]$ on S is known to be an interval-valued Pythagorean fuzzy sub-semigroup of S . If for all $w_1, w_2 \in S$, it holds.

$$\tilde{\phi}_p(w_1 w_2) \geq \min\{\tilde{\phi}_p(w_1), \tilde{\phi}_p(w_2)\},$$

$$\tilde{\psi}_p(w_1 w_2) \leq \max\{\tilde{\psi}_p(w_1), \tilde{\psi}_p(w_2)\}.$$

Example 1. Consider a semigroup $S = \{u, v, w, x, y\}$ with the Cayley Table.

Table 1. Cayley table.

•	u	v	w	x	y
u	u	u	u	u	u
v	u	v	u	x	u
w	u	y	w	w	y
x	u	v	x	x	v
y	u	y	u	w	u

Define an interval-valued Pythagorean fuzzy set(IVPFS) $\tilde{P} = [\tilde{\phi}_p, \tilde{\psi}_p]$ in S as follows.

S	$[\tilde{\phi}_p(w_1), \tilde{\psi}_p(w_1)]$
u	[0.7,0.8], [0.1,0.2]
v	[0.4,0.6], [0.4,0.5]
w	[0.3,0.5], [0.5,0.6]
x	[0.1,0.2], [0.3,0.5]
y	[0.3,0.5], [0.5,0.6]

$$\begin{aligned}\widetilde{\Phi}_p(uv) &\geq \min\{\widetilde{\Phi}_p(u), \widetilde{\Phi}_p(v)\} \\ ([0.7, 0.8], [0.1, 0.2]) &\geq [0.4, 0.6], [0.1, 0.2]. \\ \widetilde{\Psi}_p(uv) &\leq \max\{\widetilde{\Psi}_p(u), \widetilde{\Psi}_p(v)\}. \\ ([0.7, 0.8], [0.1, 0.2]) &\leq [0.7, 0.8], [0.4, 0.5].\end{aligned}$$

Thus $\tilde{P} = [\widetilde{\Phi}_p, \widetilde{\Psi}_p]$ is an Interval-Valued Pythagorean Fuzzy Sub-Semigroup (IVPFSS) of S .

Definition 3. An IVPFS $\tilde{P} = (\widetilde{\Phi}_p, \widetilde{\Psi}_p)$ on semigroup S , is said to be an interval-valued Pythagorean fuzzy left (\tilde{P}_{LI})(resp. right(\tilde{P}_{RI})) ideal of S . If for all $w_1, w_2 \in S$, it holds.

$$\begin{aligned}\widetilde{\Phi}_p(w_1 w_2) &\geq \widetilde{\Phi}_p(w_2); \\ \widetilde{\Psi}_p(w_1 w_2) &\leq \widetilde{\Psi}_p(w_2) \text{ (resp. right } (\tilde{P}_{RI}); \\ \widetilde{\Phi}_p(w_1 w_2) &\geq \widetilde{\Phi}_p(w_1); \\ \widetilde{\Psi}_p(w_1 w_2) &\leq \widetilde{\Psi}_p(w_1).\end{aligned}$$

Definition 4. An IVPFS $\tilde{P} = [\widetilde{\Phi}_p, \widetilde{\Psi}_p]$ on S is called IVPFI (\tilde{P}_I) of S . If for all $w_1, w_2 \in S$, it \tilde{P} is both a left and right IVPFI of S .

$$\begin{aligned}(w_1 w_2) &\geq \max\{\widetilde{\Phi}_p(w_1), \widetilde{\Phi}_p(w_2)\}; \\ \widetilde{\Psi}_p(w_1 w_2) &\leq \min\{\widetilde{\Psi}_p(w_1), \widetilde{\Psi}_p(w_2)\}.\end{aligned}$$

Definition 5. An IVPFS $\tilde{P} = [\widetilde{\Phi}_p, \widetilde{\Psi}_p]$ on S is known to be an interval-Valued Pythagorean Fuzzy Bi-Ideal (IVPFBI) (\tilde{P}_{BI}) of S . If for all $a, w_1, w_2 \in S$ and satisfy.

$$\begin{aligned}\widetilde{\Phi}_p(w_1 a w_2) &\geq \min\{\widetilde{\Phi}_p(w_1), \widetilde{\Phi}_p(w_2)\}; \\ \widetilde{\Psi}_p(w_1 a w_2) &\leq \max\{\widetilde{\Psi}_p(w_1), \widetilde{\Psi}_p(w_2)\}.\end{aligned}$$

Example 2. Consider a semigroup $S = \{u, v, w, x, y\}$ with the Cayley Table.

Define an interval-valued Pythagorean fuzzy set $\tilde{P} = [\widetilde{\Phi}_p, \widetilde{\Psi}_p]$ in S as follows.

S	$[\widetilde{\Phi}_p(w_1), \widetilde{\Psi}_p(w_1)]$
u	$[0.8, 0.9], [0.1, 0.3]$
v	$[0.3, 0.5], [0.7, 0.9]$
w	$[0.4, 0.6], [0.6, 0.7]$
x	$[0.3, 0.5], [0.7, 0.9]$
y	$[0.7, 0.8], [0.4, 0.5]$

Thus $\tilde{P} = [\widetilde{\Phi}_p, \widetilde{\Psi}_p]$ is an interval valued Pythagorean fuzzy bi-ideal of S .

Definition 7. An IVPFS $\tilde{P} = \langle [\widetilde{\Phi}_p, \widetilde{\Psi}_p] \rangle$ on S is known to be an interval-valued Pythagorean fuzzy interior ideal (IVPFII) (\tilde{P}_{II}) of S . If for all $a, w_1, w_2 \in S$ and satisfy.

$$\widetilde{\Phi}_p(w_1 a w_2) \geq \widetilde{\Phi}_p(a);$$

$$\widetilde{\psi}_p(w_1 a w_2) \leq \widetilde{\psi}_p(a).$$

Definition 8. For any non-empty subset N of a semigroup S is defined to be a structure $\chi_N = \{w_1, [\tilde{\phi}_{\chi_N}(w_1), \tilde{\psi}_{\chi_N}(w_1)] | w_1 \in S\}$ which is briefly denoted by $\chi_N = [\tilde{\phi}_{\chi_N}, \tilde{\psi}_{\chi_N}]$

$$\text{where, } \tilde{\phi}_{\chi_N}(w_1) = \begin{cases} \tilde{1} & \text{if } x \in N \\ \tilde{0} & \text{otherwise} \end{cases} \quad \tilde{\psi}_{\chi_N}(w_1) = \begin{cases} \tilde{0} & \text{if } x \in N \\ \tilde{1} & \text{otherwise} \end{cases}.$$

Theorem 1. Let S be a semigroup. Then the following are equivalent.

- The intersection of two interval-valued Pythagorean fuzzy sub-semigroup of S , is an interval-valued Pythagorean fuzzy sub-semigroup of S .
- The intersection of two interval-valued Pythagorean fuzzy left (resp. right) ideal of S , is IVPFLI (resp. IVPFRI) of S .

Proof. Let $\tilde{P}_1 = [\tilde{\phi}_{p_1}, \tilde{\psi}_{p_1}]$ and $\tilde{P}_2 = [\tilde{\phi}_{p_2}, \tilde{\psi}_{p_2}]$ be two interval-valued Pythagorean fuzzy sub-semigroup of S . Let $w_1, w_2 \in S$.

Then,

$$\begin{aligned} (\tilde{\phi}_{p_1} \cap \tilde{\phi}_{p_2})(w_1, w_2) &= \min\{\tilde{\phi}_{p_1}(w_1, w_2), \tilde{\phi}_{p_2}(w_1, w_2)\} \\ &\geq \min\{\min\{\tilde{\phi}_{p_1}(w_1), \tilde{\phi}_{p_1}(w_2)\}, \min\{\tilde{\phi}_{p_2}(w_1), \tilde{\phi}_{p_2}(w_2)\}\} \\ &= \min\{\min\{\tilde{\phi}_{p_1}(w_1), \tilde{\phi}_{p_2}(w_1)\}, \min\{\tilde{\phi}_{p_1}(w_2), \tilde{\phi}_{p_2}(w_2)\}\} \\ &= \min\{\tilde{\phi}_{p_1} \cap \tilde{\phi}_{p_2}(w_1), \tilde{\phi}_{p_1} \cap \tilde{\phi}_{p_2}(w_2)\}; \end{aligned}$$

$$\begin{aligned} (\tilde{\psi}_{p_1} \cup \tilde{\psi}_{p_2})(w_1, w_2) &= \max\{\tilde{\psi}_{p_1}(w_1, w_2), \tilde{\psi}_{p_2}(w_1, w_2)\} \\ &\leq \max\{\max\{\tilde{\psi}_{p_1}(w_1), \tilde{\psi}_{p_1}(w_2)\}, \max\{\tilde{\psi}_{p_2}(w_1), \tilde{\psi}_{p_2}(w_2)\}\} \\ &= \max\{\max\{\tilde{\psi}_{p_1}(w_1), \tilde{\psi}_{p_2}(w_1)\}, \max\{\tilde{\psi}_{p_1}(w_2), \tilde{\psi}_{p_2}(w_2)\}\} \\ &= \max\{\tilde{\psi}_{p_1} \cup \tilde{\psi}_{p_2}(w_1), \tilde{\psi}_{p_1} \cup \tilde{\psi}_{p_2}(w_2)\}. \end{aligned}$$

Therefore, $\tilde{P}_1 \cap \tilde{P}_2 = \{(\tilde{\phi}_{p_1} \cap \tilde{\phi}_{p_2}), (\tilde{\psi}_{p_1} \cup \tilde{\psi}_{p_2})\}$.

Interval-valued Pythagorean fuzzy sub-semigroup of S .

$$\begin{aligned} (\tilde{\phi}_{p_1} \cap \tilde{\phi}_{p_2})(w_1, w_2) &= \min\{\tilde{\phi}_{p_1}(w_1, w_2), \tilde{\phi}_{p_2}(w_1, w_2)\} \\ &\geq \min\{\tilde{\phi}_{p_1}(w_2), \tilde{\phi}_{p_2}(w_2)\} \\ &= (\tilde{\phi}_{p_1} \cap \tilde{\phi}_{p_2})(w_2); \end{aligned}$$

$$\begin{aligned} (\tilde{\psi}_{p_1} \cup \tilde{\psi}_{p_2})(w_1, w_2) &= \max\{\tilde{\psi}_{p_1}(w_1, w_2), \tilde{\psi}_{p_2}(w_1, w_2)\} \\ &\leq \max\{\tilde{\psi}_{p_1}(w_2), \tilde{\psi}_{p_2}(w_2)\} \\ &= (\tilde{\psi}_{p_1} \cup \tilde{\psi}_{p_2})(w_2). \end{aligned}$$

Therefore, $\tilde{P}_1 \cap \tilde{P}_2 = \{(\tilde{\phi}_{p_1} \cap \tilde{\phi}_{p_2}), (\tilde{\psi}_{p_1} \cup \tilde{\psi}_{p_2})\}$ is an interval-valued Pythagorean fuzzy left (resp. right) ideal of S .

Theorem 2. An IVPFS $\tilde{P} = [\tilde{\phi}_p, \tilde{\psi}_p]$ of a semigroup S is an IVPFBI of S , if and only if $\langle (\phi_p^L, \phi_p^U), (\psi_p^L, \psi_p^U) \rangle$ of S .

Proof. Let $\tilde{P} = [\tilde{\phi}_p, \tilde{\psi}_p]$ be an interval-valued Pythagorean fuzzy bi-ideal of S , for any $w_1, w_2 \in S$.

Then, we have membership

$$\begin{aligned} [\phi_p^L(w_1 w_2), \phi_p^U(w_1 w_2)] &= \tilde{\phi}_p(w_1 w_2) \\ &\geq \min\{\tilde{\phi}_p(w_1), \tilde{\phi}_p(w_2)\} \\ &= \min\{[\phi_p^L(w_1), \phi_p^U(w_1)], [\phi_p^L(w_2), \phi_p^U(w_2)]\} \\ &= \min\{[\phi_p^L(w_1), \phi_p^L(w_2)], [\phi_p^U(w_1), \phi_p^U(w_2)]\}. \end{aligned}$$

It follows that $\phi_p^L(w_1 w_2) \geq \min\{\phi_p^L(w_1), \phi_p^L(w_2)\}$ and $\phi_p^U(w_1 w_2) \geq \min\{\phi_p^U(w_1), \phi_p^U(w_2)\}$ and non-membership

$$\begin{aligned} [\psi_p^L(w_1 w_2), \psi_p^U(w_1 w_2)] &= \tilde{\psi}_p(w_1 w_2) \\ &\leq \max\{\tilde{\psi}_p(w_1), \tilde{\psi}_p(w_2)\} \\ &= \max\{[\psi_p^L(w_1), \psi_p^U(w_1)], [\psi_p^L(w_2), \psi_p^U(w_2)]\} \\ &= \max\{[\psi_p^L(w_1), \psi_p^L(w_2)], [\psi_p^U(w_1), \psi_p^U(w_2)]\}. \end{aligned}$$

It follows that $\psi_p^L(w_1 w_2) \leq \max\{\psi_p^L(w_1), \psi_p^L(w_2)\}$ and $\psi_p^U(w_1 w_2) \leq \max\{\psi_p^U(w_1), \psi_p^U(w_2)\}$

Therefore, $\tilde{P} = \langle (\phi_p^L, \phi_p^U), (\psi_p^L, \psi_p^U) \rangle$ are Pythagorean fuzzy ideal of S .

Conversely, suppose that $([\phi_p^L, \phi_p^U], [\psi_p^L, \psi_p^U])$ are Pythagorean fuzzy ideal of S , let $w_1, w_2 \in S$ t.

$$\begin{aligned} \tilde{\phi}_p(w_1 w_2) &= [\phi_p^L(w_1 w_2), \phi_p^U(w_1 w_2)] \\ &\geq [\min\{\phi_p^L(w_1), \phi_p^L(w_2)\}, \min\{\phi_p^U(w_1), \phi_p^U(w_2)\}] \\ &= \min\{[\phi_p^L(w_1), \phi_p^U(w_1)], [\phi_p^L(w_2), \phi_p^U(w_2)]\} \\ &= \min\{\tilde{\phi}_p(w_1), \tilde{\phi}_p(w_2)\}; \end{aligned}$$

$$\begin{aligned} \tilde{\psi}_p(w_1 w_2) &= [\psi_p^L(w_1 w_2), \psi_p^U(w_1 w_2)] \\ &\leq [\max\{\psi_p^L(w_1), \psi_p^L(w_2)\}, \max\{\psi_p^U(w_1), \psi_p^U(w_2)\}] \\ &= \max\{[\psi_p^L(w_1), \psi_p^U(w_1)], [\psi_p^L(w_2), \psi_p^U(w_2)]\} \\ &= \max\{\tilde{\psi}_p(w_1), \tilde{\psi}_p(w_2)\}. \end{aligned}$$

$\tilde{P} = [\tilde{\phi}_p, \tilde{\psi}_p]$ is an interval-valued Pythagorean fuzzy sub-semigroup of S .

$$\begin{aligned} \tilde{\phi}_p(w_1 a w_2) &= [\phi_p^L(w_1 a w_2), \phi_p^U(w_1 a w_2)] \\ &\geq [\min\{\phi_p^L(w_1), \phi_p^L(w_2)\}, \min\{\phi_p^U(w_1), \phi_p^U(w_2)\}] \\ &= \min\{[\phi_p^L(w_1), \phi_p^U(w_1)], [\phi_p^L(w_2), \phi_p^U(w_2)]\} \\ &= \min\{\tilde{\phi}_p(w_1), \tilde{\phi}_p(w_2)\}; \end{aligned}$$

$$\begin{aligned} \tilde{\psi}_p(w_1 a w_2) &= [\psi_p^L(w_1 a w_2), \psi_p^U(w_1 a w_2)] \\ &\leq [\max\{\psi_p^L(w_1), \psi_p^L(w_2)\}, \max\{\psi_p^U(w_1), \psi_p^U(w_2)\}] \\ &= \max\{[\psi_p^L(w_1), \psi_p^U(w_1)], [\psi_p^L(w_2), \psi_p^U(w_2)]\} \end{aligned}$$

$$= \max\{\tilde{\psi}_p(w_1), \tilde{\psi}_p(w_2)\}.$$

$\tilde{P} = [\tilde{\phi}_p, \tilde{\psi}_p]$ is an interval-valued Pythagorean fuzzy bi-ideal of S .

Theorem 3. If $\{P_i\}_{i \in I}$ is a family of interval-valued Pythagorean fuzzy bi-ideal of a semigroup S . Then $\cap P_i$ is an interval-valued Pythagorean fuzzy bi-ideal of S . Where $\cap P_i = (\cap \tilde{\phi}_{p_i}, \cup \tilde{\psi}_{p_i})$.

$\cap (\tilde{\phi}_{p_i}) = \inf\{(\tilde{\phi}_{p_i})(w_1)/i \in I, w_1 \in S\}$, $\cup (\tilde{\psi}_{p_i}) = \sup\{(\tilde{\psi}_{p_i})(w_1)/i \in I, w_1 \in S\}$ and $i \in I$ is any index set.

Proof. Since $\tilde{P}_i = \langle [\tilde{\phi}_{p_i}, \tilde{\psi}_{p_i}] | i \in I \rangle$ is a family of interval-valued Pythagorean fuzzy bi-ideal of S .

Let $a, w_1, w_2 \in S$.

$$\begin{aligned} \cap \tilde{\phi}_{p_i}(w_1, w_2) &= \inf\{\tilde{\phi}_{p_i}(w_1, w_2)/i \in I, w_1, w_2 \in S\} \\ &\geq \inf\{\min\{\tilde{\phi}_{p_i}(w_1), \tilde{\phi}_{p_i}(w_2)\}\} \\ &= \min\{\inf(\tilde{\phi}_{p_i}(w_1)), \inf(\tilde{\phi}_{p_i}(w_2))\} \\ &= \min\{\cap \tilde{\phi}_{p_i}(w_1), \cap \tilde{\phi}_{p_i}(w_2)\}; \end{aligned}$$

$$\begin{aligned} \cup \tilde{\psi}_{p_i}(w_1 w_2) &= \sup\{\tilde{\psi}_{p_i}(w_1 w_2)/i \in I, w_1, w_2 \in S\} \\ &\leq \sup\{\max\{\tilde{\psi}_{p_i}(w_1), \tilde{\psi}_{p_i}(w_2)\}\} \\ &= \max\{\sup(\tilde{\psi}_{p_i}(w_1)), \sup(\tilde{\psi}_{p_i}(w_2))\} \\ &= \max\{\cup \tilde{\psi}_{p_i}(w_1), \cup \tilde{\psi}_{p_i}(w_2)\}. \end{aligned}$$

Hence, $\cap \tilde{P}_i = (\cap \tilde{\phi}_{p_i}, \cup \tilde{\psi}_{p_i})$ is an interval-valued Pythagorean fuzzy sub-semigroup of S .

$$\begin{aligned} \cap \tilde{\phi}_{p_i}(w_1 a w_2) &= \inf\{\tilde{\phi}_{p_i}(w_1 a w_2)/i \in I, a, w_1, w_2 \in S\} \\ &\geq \inf\{\min\{\tilde{\phi}_{p_i}(w_1), \tilde{\phi}_{p_i}(w_2)\}\} \\ &= \min\{\inf(\tilde{\phi}_{p_i}(w_1)), \inf(\tilde{\phi}_{p_i}(w_2))\} \\ &= \min\{\cap \tilde{\phi}_{p_i}(w_1), \cap \tilde{\phi}_{p_i}(w_2)\}. \end{aligned}$$

$$\begin{aligned} \cup \tilde{\psi}_{p_i}(w_1 a w_2) &= \sup\{\tilde{\psi}_{p_i}(w_1 a w_2)/i \in I, a, w_1, w_2 \in S\} \\ &\leq \sup\{\max\{\tilde{\psi}_{p_i}(w_1), \tilde{\psi}_{p_i}(w_2)\}\} \\ &= \max\{\sup(\tilde{\psi}_{p_i}(w_1)), \sup(\tilde{\psi}_{p_i}(w_2))\} \\ &= \max\{\cup \tilde{\psi}_{p_i}(w_1), \cup \tilde{\psi}_{p_i}(w_2)\}. \end{aligned}$$

Hence, $\cap P_i = (\cap \tilde{\phi}_{p_i}, \cup \tilde{\psi}_{p_i})$ is an interval-valued Pythagorean fuzzy bi-ideals of S .

Theorem 4. Let N be any non-empty subset of a semigroup S . Then N is a bi-ideal of S , if and only if the characteristic interval-valued Pythagorean fuzzy set $\chi_N = [\tilde{\phi}_{p_{\chi_N}}, \tilde{\psi}_{p_{\chi_N}}]$ is IVPFBI of S .

Proof. Assume that N is a bi-ideal of S . Let $a, w_1, w_2 \in S$.

Suppose that $\tilde{\phi}_{p\chi_N}(w_1w_2) < \min\{\tilde{\phi}_{p\chi_N}(w_1), \tilde{\phi}_{p\chi_N}(w_2)\}$ and $\tilde{\psi}_{p\chi_N}(w_1w_2) > \max\{\tilde{\psi}_{p\chi_N}(w_1), \tilde{\psi}_{p\chi_N}(w_2)\}$ it follows that $\tilde{\phi}_{p\chi_N}(w_1w_2) = 0$, $\min\{\tilde{\phi}_{p\chi_N}(w_1), \tilde{\phi}_{p\chi_N}(w_2)\} = 1$

$$\tilde{\psi}_{p\chi_N}(w_1w_2) = 1, \max\{\tilde{\psi}_{p\chi_N}(w_1), \tilde{\psi}_{p\chi_N}(w_2)\} = 0.$$

This implies that $w_1, w_2 \in N$ by $w_1, w_2 \notin N$ a contradiction to N .

$$\text{So } \tilde{\phi}_{p\chi_N}(w_1w_2) \geq \min\{\tilde{\phi}_{p\chi_N}(w_1), \tilde{\phi}_{p\chi_N}(w_2)\}, \tilde{\psi}_{p\chi_N}(w_1w_2) \leq \max\{\tilde{\psi}_{p\chi_N}(w_1), \tilde{\psi}_{p\chi_N}(w_2)\}.$$

Suppose that $\tilde{\phi}_{p\chi_N}(w_1aw_2) < \min\{\tilde{\phi}_{p\chi_N}(w_1), \tilde{\phi}_{p\chi_N}(w_2)\}$ and $\tilde{\psi}_{p\chi_N}(w_1aw_2) > \max\{\tilde{\psi}_{p\chi_N}(w_1), \tilde{\psi}_{p\chi_N}(w_2)\}$ it follows that $\tilde{\phi}_{p\chi_N}(w_1aw_2) = 0$, $\min\{\tilde{\phi}_{p\chi_N}(w_1), \tilde{\phi}_{p\chi_N}(w_2)\} = 1$, $\tilde{\psi}_{p\chi_N}(w_1aw_2) = 1$, $\max\{\tilde{\psi}_{p\chi_N}(w_1), \tilde{\psi}_{p\chi_N}(w_2)\} = 0$.

This implies that $a, w_1, w_2 \in N$ by $a, w_1, w_2 \notin N$ a contradiction to N .

$$\text{So } \tilde{\phi}_{p\chi_N}(w_1aw_2) \geq \min\{\tilde{\phi}_{p\chi_N}(w_1), \tilde{\phi}_{p\chi_N}(w_2)\}, \tilde{\psi}_{p\chi_N}(w_1aw_2) \leq \max\{\tilde{\psi}_{p\chi_N}(w_1), \tilde{\psi}_{p\chi_N}(w_2)\}.$$

This shows that χ_N is an interval-valued Pythagorean fuzzy bi-ideal of S .

Conversely, $\chi_N = [\tilde{\phi}_{p\chi_N}, \tilde{\psi}_{p\chi_N}]$ is an IVPFBI of S for any subset N of S .

Let $w_1, w_2 \in N$ then $\tilde{\phi}_{p\chi_N}(w_1) = \tilde{\phi}_{p\chi_N}(w_2) = \tilde{1}$, $\tilde{\psi}_{p\chi_N}(w_1) = \tilde{\psi}_{p\chi_N}(w_2) = \tilde{0}$, since χ_N is an IVPFBI of S .

$$\tilde{\phi}_{p\chi_N}(w_1w_2) \geq \min\{\tilde{\phi}_{p\chi_N}(w_1), \tilde{\phi}_{p\chi_N}(w_2)\} \geq \min\{\tilde{1}, \tilde{1}\} = \tilde{1}, \tilde{\psi}_{p\chi_N}(w_1w_2) \leq \max\{\tilde{\psi}_{p\chi_N}(w_1), \tilde{\psi}_{p\chi_N}(w_2)\} \leq \max\{\tilde{0}, \tilde{0}\} = \tilde{0}.$$

This implies that $w_1, w_2 \in N$.

$$\text{Let } a, w_1, w_2 \in N \text{ then } \tilde{\phi}_{p\chi_N}(w_1) = \tilde{\phi}_{p\chi_N}(a) = \tilde{\phi}_{p\chi_N}(w_2) = \tilde{1},$$

$$\tilde{\psi}_{p\chi_N}(w_1) = \tilde{\psi}_{p\chi_N}(a) = \tilde{\psi}_{p\chi_N}(w_2) = \tilde{0}, \text{ since } \chi_N \text{ is an IVPFBI of } S.$$

$$\tilde{\phi}_{p\chi_N}(w_1aw_2) \geq \min\{\tilde{\phi}_{p\chi_N}(w_1), \tilde{\phi}_{p\chi_N}(w_2)\} \geq \min\{\tilde{1}, \tilde{1}\} = \tilde{1}, \tilde{\psi}_{p\chi_N}(w_1aw_2) \leq \max\{\tilde{\psi}_{p\chi_N}(w_1), \tilde{\psi}_{p\chi_N}(w_2)\} \leq \max\{\tilde{0}, \tilde{0}\} = \tilde{0}.$$

Which implies that $w_1, w_2 \in N$. Hence N is a bi-ideal of S .

Theorem 5. If $\{\tilde{P}_i\}_{i \in I}$ is a family of interval-valued Pythagorean fuzzy interior ideal of a semigroup S . Then $\cap \tilde{P}_i$ is an interval-valued Pythagorean fuzzy interior ideal (IVPFII) of S .

$$\text{Where } \cap \tilde{P}_i = (\cap \tilde{\phi}_{p_i}, \cup \tilde{\psi}_{p_i});$$

$$\cap (\tilde{\phi}_{p_i}) = \inf\{(\tilde{\phi}_{p_i})(w_1)/i \in I, w_1 \in S\}, \cup (\tilde{\psi}_{p_i}) = \sup\{(\tilde{\psi}_{p_i})(w_1)/i \in I, w_1 \in S\} \text{ and } i \in I \text{ is any index set.}$$

Theorem 6. Let N be any non-empty subset of a semigroup S . Then N is a interior ideal of S , if and only if the characteristic interval-valued Pythagorean fuzzy set $\chi_N = [\tilde{\phi}_{\chi_N}, \tilde{\psi}_{\chi_N}]$ is IVPFII of S .

4. Homomorphism of Interval-Valued Pythagorean Fuzzy Ideals in Semigroups

Let R and T be two non-empty sets of semigroup S . A mapping $f: R \rightarrow T$ is called a homomorphism if $(rt) = f(r)f(t) \forall r, t \in R$.

Definition 9. Let f be a mapping from a set R to a set T and $\tilde{P} = [\tilde{\phi}_p, \tilde{\psi}_p]$ be an interval-valued Pythagorean fuzzy set R the image of R (i.e.) $f(\tilde{P}) = (f(\tilde{\phi}_p), f(\tilde{\psi}_p))$ is an interval-valued Pythagorean fuzzy set of T is defined by

$$f(\tilde{P})(r) = \begin{cases} f(\tilde{\phi}_p)(r) = \begin{cases} \sup_{t \in f^{-1}(r)} (\tilde{\phi}_p)(t), & \text{iff } f^{-1}(r) \neq \emptyset \\ [0, 0] & \text{otherwise} \end{cases} \\ f(\tilde{\psi}_p)(r) = \begin{cases} \inf_{t \in f^{-1}(r)} (\tilde{\psi}_p)(t), & \text{iff } f^{-1}(r) \neq \emptyset \\ [1, 1] & \text{otherwise} \end{cases} \end{cases}$$

Let f be a mapping from a set R to T and $\tilde{P} = [\tilde{\phi}_p, \tilde{\psi}_p]$ be an interval-valued Pythagorean fuzzy set of T then the preimage of T (i.e.) $f^{-1}(\tilde{P}) = \{(f^{-1}(\tilde{\phi}_p), f^{-1}(\tilde{\psi}_p))\}$ is an interval-valued Pythagorean fuzzy set of R is defined as

$$f^{-1}(\tilde{P})(r) = \begin{cases} f^{-1}(\tilde{\phi}_p)(r) = \tilde{\phi}_p(f(r)) \\ f^{-1}(\tilde{\psi}_p)(r) = \tilde{\psi}_p(f(r)) \end{cases}$$

Theorem 7. Let R, T be a semigroups, $f: R \rightarrow T$ be a homomorphism of semigroups.

If $\tilde{P} = [\tilde{\phi}_p, \tilde{\psi}_p]$ is an interval-valued Pythagorean fuzzy sub-semigroup of T the the preimage $f^{-1}(\tilde{P}) = (f^{-1}(\tilde{\phi}_p), f^{-1}(\tilde{\psi}_p))$ is an interval-valued Pythagorean fuzzy sub-semigroup of R .

If $\tilde{P} = [\tilde{\phi}_p, \tilde{\psi}_p]$ is an interval-valued Pythagorean fuzzy left (resp. right) ideal of T the the preimage $f^{-1}(\tilde{P}) = (f^{-1}(\tilde{\phi}_p), f^{-1}(\tilde{\psi}_p))$ is an interval-valued Pythagorean fuzzy left ideal (resp. right ideal) of R .

Proof. Assume that $\tilde{P} = [\tilde{\phi}_p, \tilde{\psi}_p]$ is an interval-valued Pythagorean fuzzy sub-semigroup of T and $r, t \in R$. Then

$$\begin{aligned} f^{-1}(\tilde{\phi}_p)(rt) &= \tilde{\phi}_p(f(rt)) \\ &= \tilde{\phi}_p(f(r)f(t)) \\ &\geq \min\{\tilde{\phi}_p(f(r)), \tilde{\phi}_p(f(t))\} \\ &= \min\{f^{-1}(\tilde{\phi}_p)(r), f^{-1}(\tilde{\phi}_p)(t)\}; \end{aligned}$$

$$f^{-1}(\tilde{\psi}_p)(rt) = \tilde{\psi}_p(f(rt))$$

$$\begin{aligned}
 &= \tilde{\Psi}_p(f(r)f(t)) \\
 &\leq \max\{\tilde{\Psi}_p(f(r)), \tilde{\Psi}_p(f(t))\} \\
 &= \max\{f^{-1}(\tilde{\Psi}_p)(r), f^{-1}(\tilde{\Psi}_p)(f(t))\}.
 \end{aligned}$$

Hence, $f^{-1}(\tilde{P}) = (f^{-1}(\tilde{\Phi}_p), f^{-1}(\tilde{\Psi}_p))$ is an interval-valued Pythagorean fuzzy sub-semigroup of R .

$$\begin{aligned}
 f^{-1}(\tilde{\Phi}_p)(rt) &= \tilde{\Phi}_p(f(rt)) \\
 &= \tilde{\Phi}_p(f(r)f(t)) \\
 &\geq \tilde{\Phi}_p(f(t)) \\
 &= f^{-1}(\tilde{\Phi}_p)(f(t));
 \end{aligned}$$

$$\begin{aligned}
 f^{-1}(\tilde{\Psi}_p)(rt) &= \tilde{\Psi}_p(f(rt)) \\
 &= \tilde{\Psi}_p(f(r)f(t)) \\
 &\leq \tilde{\Psi}_p(f(t)) \\
 &= f^{-1}(\tilde{\Psi}_p)(f(t)).
 \end{aligned}$$

Hence, $f^{-1}(\tilde{P}) = (f^{-1}(\tilde{\Phi}_p), f^{-1}(\tilde{\Psi}_p))$ is an interval-valued Pythagorean fuzzy left (resp.right) ideal of R .

Theorem 8. Let R, T be a semigroups, $f: R \rightarrow T$ be a homomorphism of semigroups. If $\tilde{P} = [\tilde{\Phi}_p, \tilde{\Psi}_p]$ is an interval-valued Pythagorean fuzzy bi-ideal of T the the preimage $f^{-1}(\tilde{P}) = (f^{-1}(\tilde{\Phi}_p), f^{-1}(\tilde{\Psi}_p))$ is an interval-valued Pythagorean fuzzy bi-ideal of R .

Proof. Assume that $\tilde{P} = [\tilde{\Phi}_p, \tilde{\Psi}_p]$ is an interval-valued Pythagorean fuzzy sub-semigroup of T and $a, r, t \in R$. Then

$$\begin{aligned}
 f^{-1}(\tilde{\Phi}_p)(rat) &= \tilde{\Phi}_p(f(rat)) \\
 &= \tilde{\Phi}_p(f(r)f(a)f(t)) \\
 &\geq \min\{\tilde{\Phi}_p(f(r)), \tilde{\Phi}_p(f(t))\} \\
 &= \min\{f^{-1}(\tilde{\Phi}_p)(r), f^{-1}(\tilde{\Phi}_p)(f(t))\};
 \end{aligned}$$

$$\begin{aligned}
 f^{-1}(\tilde{\Psi}_p)(rat) &= \tilde{\Psi}_p(f(rat)) \\
 &= \tilde{\Psi}_p(f(r)f(a)f(t)) \\
 &\leq \max\{\tilde{\Psi}_p(f(r)), \tilde{\Psi}_p(f(t))\} \\
 &= \max\{f^{-1}(\tilde{\Psi}_p)(r), f^{-1}(\tilde{\Psi}_p)(f(t))\}.
 \end{aligned}$$

Hence $f^{-1}(\tilde{P}) = (f^{-1}(\tilde{\Phi}_p), f^{-1}(\tilde{\Psi}_p))$ is an interval-valued Pythagorean fuzzy bi-ideal of R .

Theorem 9. Let R, T be a semigroups, $f: R \rightarrow T$ be a homomorphism of semigroups. If $\tilde{P} = [\tilde{\Phi}_p, \tilde{\Psi}_p]$ is an interval-valued Pythagorean fuzzy interior ideal of T the preimage $f^{-1}(\tilde{P}) = (f^{-1}(\tilde{\Phi}_p), f^{-1}(\tilde{\Psi}_p))$ is an interval-valued Pythagorean fuzzy interior ideal of R .

5. Conclusion

In this paper interval valued Pythagorean fuzzy sub-semigroup, interval valued Pythagorean fuzzy left (resp. right) ideal, interval valued Pythagorean fuzzy ideal, interval valued Pythagorean fuzzy bi-ideal, interval valued Pythagorean fuzzy interior ideal and Homomorphism of interval valued Pythagorean fuzzy ideal in semigroups are studied and investigated some properties with suitable examples.

References

- [1] Akram, M. (2018). *Fuzzy lie algebras*. Springer.
- [2] Atanassov, K. T. (1986). Intuitionistic fuzzy sets. *Fuzzy sets and system*, 20, 87-96.
- [3] Atanassov, K. T., & Gargov, G. (1989). Interval valued intuitionistic fuzzy sets. *Fuzzy sets and systems*, 31, 343-349.
- [4] Chen, T. Y. (2018). An interval valued Pythagorean fuzzy outranking method with a closeness-based assignment model for multiple criteria decision making. *International journal of intelligent systems*, 33(1), 126-168.
- [5] Chen, T. Y. (2018). An outranking approach using a risk attitudinal assignment model involving Pythagorean fuzzy information and its application to financial decision making. *Applied soft computing*, 71(1), 460-487.
- [6] Das, S., & Edalatpanah, S. A. (2016). A general form of fuzzy linear fractional progress with trapezoidal fuzzy numbers. *International journal of data envelopment analysis and operations research*, 2(1), 16-19.
- [7] Edalatpanah, S. A. (2019). A data envelopment analysis model with triangular intuitionistic fuzzy numbers. *International journal of data envelopment analysis*, 2(1), 16-19.
- [8] Garg, H. (2018). A linear programming method based on an improved score function for interval valued Pythagorean to decision making. *International journal of uncertainty, fuzziness and knowledge based systems*, 26(1), 67-80.
- [9] Garg, H. (2018). New exponential operational laws and their aggregation operators for interval valued Pythagorean fuzzy multicriteria decision making. *International journal of intelligent systems*, 33(3), 653-683.
- [10] Hussain, A., Mahmood, T., & Ali, M. I. (2019). Rough Pythagorean fuzzy ideals in semigroups. *Computational and applied mathematics*, 38(2), 67.
- [11] Kumar, R., Edalatpanah, S. A., Jha, S., & Sing, R. (2019). A Pythagorean fuzzy approach to the transportation. *Complex and intelligent systems*, 5(2), 255-263.
- [12] Najafi, H. S., & Edalatpanah, S. A. (2013). Iterative methods with analytical preconditioning technique to linear complementarity problems: application to obstacle problems. *RAIRO-operations research*, 47(1), 59-71.
- [13] Peng, X. (2019). New operations for interval valued Pythagorean fuzzy set. *Scientia iranica*, 26(2), 1049-1076.
- [14] Peng, X., & Yang, Y. (2016). Fundamental properties of interval valued Pythagorean fuzzy aggregation operators. *International journal of intelligent systems*, 31(5), 444-487.
- [15] Thillaigovindan, N., & Chinnadurai, V. (2009). On interval-valued fuzzy quasi-ideals of semigroups. *East Asian mathematical journal*, 25(4), 441-453.
- [16] Thillaigovindan, N., & Chinnadurai, V. (2009). Interval-valued fuzzy generalized bi-ideals. *Proceedings of the national conference on algebra, graph theory and their applications* (pp. 85-98). Department of Mathematics, Manonmaniam Sundaranar University, Narosa.
- [17] Yager, R. R. (2013, June). Pythagorean fuzzy subsets. In *2013 joint IFSA world congress and NAFIPS annual meeting (IFSA/NAFIPS)* (pp. 57-61). IEEE.
- [18] Zadeh, L. A. (1965). Fuzzy sets. *Information and control*, 8, 338-353.
- [19] Zadeh, L. A. (1975). The concept of a linguistic variable and its application to approximate reasoning –I. *Information science*, 8, 199-249.